

MON-14420 Herbicide by Monsanto
WATER SOLUBLE GRANULE
PREMEASURED IN WATER SOLUBLE BAG

For nonselective and selective weed control on industrial sites such as roadsides and associated rights-of-way.


Complete Directions for Use

1988-1

MAP-2982.01

©MONSANTO COMPANY 1988

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167, U.S.A.



Read the entire label before using this product.
Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described herein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit Of Warranty and Liability which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

DANGER!

CAUSES EYE BURNS
HARMFUL IF SWALLOWED.

Do not get in eyes, on skin or on clothing.

Wear goggles or face shield and rubber gloves in mixing and when adjusting, repairing or cleaning equipment.

Wash thoroughly with soap and water after handling.

product 524-424 2/10
FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention. Remove material from skin and clothing.

IF SWALLOWED, this product may cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetland (swamps, bogs, marshes or potholes). Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. Spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl)glycine in the form
of the monoammonium salt..... 74.7%
INERT INGREDIENTS:..... 25.3%

100.0%

PH 20

524 424

11/2/10

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

STORAGE:

Keep the outer bag **TIGHTLY** sealed to prevent moisture from damaging any unused water soluble bags.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Completely use water soluble pouches in application equipment. Then dispose of empty box and outer bag in a sanitary landfill, or by incineration, or, if allowed by state or local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product is a water soluble granule herbicide packaged in a premeasured water soluble bag. This product mixes with water to be applied as a foliar spray for the nonselective or selective control of herbaceous plants in industrial sites such as roadsides and associated rights-of-way. It controls most undesirable annual and perennial grasses, broadleaf weeds, woody brush and vine species. This product is absorbed by the leaves of actively growing vegetation, and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control. Symptoms are a gradual wilting and yellowing of the sprayed plant, followed by deterioration of both shoots and roots. This product has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation.

**APPLICATION PRECAUTIONS
AND WARNINGS**

AVOID DRIFT AND SPRAY CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

The likelihood of injury occurring to adjacent crops from use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when conditions favor spray drift. Apply when there is no hazard from spray drift, since very small quantities of spray which may not be visible may severely injure nearby desirable vegetation.

Reduced performance may occur when grasses or weeds are under poor growing conditions caused by drought stress, disease or insect damage. Reduced results may also occur when treating weeds heavily covered with dust, or if using water contaminated with soil, such as water from ponds or ditches.

Rainfall or watering treated plants within 6 hours after application may reduce effectiveness of this product.

Applications of this product to undesirable vegetation that has been recently mowed, cut, tilled, sliced, cored or mechanically disturbed may result in reduced performance. Allow adequate regrowth of the foliage to ensure good spray coverage. See the "WHEN TO APPLY" section for more information.

This product does not provide residual weed control. For residual weed control, follow an approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

WHEN TO APPLY

Apply to foliage of vigorously growing plants. For best results, apply to annual grasses and broadleaf weeds any time after emergence; apply to perennial grasses and broadleaf weeds after seedheads, flowers or berries appear. On woody brush and vines, apply after leaves have fully expanded. Best results are achieved on most woody plants when applications are made in summer or fall. Apply before fall leaf colors appear or a killing frost. In arid areas, apply to woody brush in the spring when flowers appear and plants have a high moisture content. Applications made at less than the recommended stage of growth may result in reduced control.

Delay mowing, clipping, tilling, planting, or sodding of treated areas for at least 7 days after applications. This allows time for this product to move within the treated foliage and into the roots for the best results. This product does not provide soil residual or herbicidal activity and it will not control new grasses or weeds emerging from seed.

One application will control most grasses and broadleaf weeds listed. Symptoms on certain tough perennials and woody brush and vine species may be slow to develop. If control is not achieved within 4 weeks, allow time for the sprayed vegetation to regrow and make a repeat application. In some cases, control of woody brush and vines treated in the fall may not be fully apparent until the following season.

Ensure uniform and complete spray coverage. Do not spray plants to the point that the spray solution runs off the foliage. Do not spray during windy conditions, as spray drift may cause damage to any vegetation contacted. Coarse sprays are less likely to drift; therefore, use nozzles and/or equipment which do not dispense spray as a fine mist.

MIXING INSTRUCTIONS

Fill the spray tank to about one-half to three-quarters of the desired volume. With the agitator running, add the recommended amount of this product. If a tank mixture is being used, then add the recommended rate of Oust™. Where nonionic surfactant is recommended, add this to the spray tank before completing the filling process. Maintain continuous agitation while completing the mixing procedure.

Remove the hose from the mixing tank immediately after filling to avoid back-siphoning into the spray source. An approved antifoam or defoaming agent can be used to minimize foaming.

™Oust is a trademark of E. I. du Pont de Nemours and Company.

524 424 7 of 12

Only nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. Use 1/2 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe the surfactant cautionary statements and other information appearing on the surfactant label.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

EQUIPMENT, SPRAY VOLUME AND CALIBRATION

Apply uniformly with properly calibrated ground boom and boomless spray equipment. Application should be uniform and of sufficient volume to avoid streaking or skips. Avoid excessive overlap of the spray.

With conventional boom and boomless ground spray equipment, use recommended rates of this product in 10 to 40 gallons of water per acre as a broadcast spray. As density of weeds increases, spray gallonage should be increased within the recommended range to ensure complete coverage.

With hand-held and high-volume spray equipment, use recommended rates of this product and apply to vegetation to be controlled on a spray-to-wet basis. Spray coverage should be uniform and complete, but do not spray to the point of runoff. Use coarse sprays only.

Nonselective Weed Control

When applied as directed, this product will provide control or partial control of the species listed in the "WEEDS CONTROLLED" section of this label. Use the following rates and mixtures for general vegetation control and bareground applications:

	Broadcast Boom, Boomless 10 to 40 GPA lbs./acre	Hand-Held High Volume Spray-to-Wet lbs./100 gal.
Annuals	1.1 to 2.2	4.4
Perennials, Woody Brush and Vines	3.3 to 5.5	4.4 to 8.8

In general, use rates should be increased within the recommended range listed above as the density or height of the infestation increases. Use the higher rates when treating hard-to-control perennials, such as bermudagrass, field bindweed Canada thistle, and woody brush and vine species.

Bareground Weed Control.

When applied as directed on sites where bareground is desired, this product plus Oust provides postemergent and preemergent control of annual weeds listed in the "WEEDS CONTROLLED" sections of the labels for this product and Oust, and control or partial control of the perennial weeds listed below.

Apply 1.1 to 2.2 pounds of this product with 2 to 4 ounces of Oust in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds. For hand-held treatments, use 4.4 to 8.8 pounds of this product plus 1 to 2 ounces of Oust in 100 gallons of spray solution. For CONTROL of annual weeds, use the lower rates of these products. For CONTROL of listed perennial weeds, use the higher rates of both products. For PARTIAL CONTROL, use the lower rates.

Bahiagrass	Johnsongrass**
Bermudagrass*	Poorjoe**
Broomsedge	Quackgrass
Dock, curly	Trumpet creeper*
Dogfennel	Vaseygrass
Fescue, tall	Vervain, blue

*Suppression at the higher rates only.

**Control at the lower rates.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used

**Selective Weed Control
in Bermudagrass and Bahiagrass**

Use the following rates and mixtures only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Dormant (winter) Applications:

Make applications to dormant bermudagrass or bahiagrass for control or suppression of many winter weeds such as little barley, annual bluegrass, Italian ryegrass, common chickweed, henbit, Carolina geranium, common vetch, crimson clover, and tall fescue. This product may be tank-mixed with Oust as recommended for residual control. This tank mixture may delay greenup. To avoid delays in greenup and minimize injury, do not add more than $\frac{1}{2}$ ounce per acre of Oust on bermudagrass or bahiagrass.

The following recommended rates of this product should be applied in 10 to 30 gallons of spray solution. Apply 0.28 to 2.2 pounds of this product alone per acre. When applying the tank mixture, use 0.28 to 0.55 pound of this product with $\frac{1}{4}$ to 1 ounce of Oust per acre. For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage. For the best recommendation for the mixture of weeds within your geographic areas, contact your Monsanto sales representative.

Active (late spring and summer) Applications:

Make applications to only actively growing and well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result. This product may be tank-mixed with Oust as recommended. When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "WEEDS CONTROLLED" section of this and the Oust label, and suppression or partial control of certain perennial weeds.

Use 0.28 to 1.1 pounds of this product alone per acre or in tank mixture with $\frac{1}{2}$ to 2 ounces of Oust per acre. Apply as a broadcast spray in 10 to 25 gallons of spray solution per acre. Use the lower rates for suppression of growth, and the higher rates to provide partial control of the following perennial weeds.

Bahiagrass
 Bluestem, silver
 Broomsedge
 Dock, curly
 Dogfennel
 Fescue, tall

Johnsongrass*
 Poorjoe**
 Trumpetcreeper*
 Vaseygrass
 Vervain, blue

*Suppression at higher rates only.
 **Control at the higher rates.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

**Bahiagrass Seedhead and
 Vegetation Suppression**

When applied as directed, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth.

Apply this product 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 0.28 pound of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 30 gallons of water per acre. Use the 0.5 percent surfactant concentration when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration for those surfactants containing less than 50 percent active ingredients.

A tank mixture of this product plus Oust may be applied only on roadsides for seedhead inhibition and vegetative suppression. Apply 0.28 pound per acre of this product plus ¼ ounce of Oust per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume 1 to 2 weeks following initial spring mowing. When using this product plus Oust for suppression of bahiagrass, make only one application per year.

WEEDS CONTROLLED

This product controls many annual and perennial grasses, broadleaf weeds, woody brush and vines.

Annual Weeds

Balsamapple	Kochia
Barley, little	Lambsquarters
Barnyardgrass	Lettuce, prickly
Bassia, fivehook	Morningglory
Bedstraw, catchweed	Mustard
Bluegrass, annual	Oats, wild
Bluegrass, bulbous	Panicum
Brome	Pennycress, field
Buttercup	Pigweed
Cheat	Ragweed
Chickweed	Rocket, London
Cocklebur	Ryegrass
Crabgrass	Sandbur, field
Dwarf dandelion	Shepherdspurse
Falseflax, smallseed	Signalgrass, broadleaf
Fiddleneck	Smartweed
Flaxleaf fleabane	Sowthistle
Fleabane	Spanishneedles
Foxtail	Stinkgrass
Geranium, Carolina	Thistle, Russian
Groundsel	Velvetleaf
Henbit	Witchgrass
Horseweed/Marestail	

Perennial Weeds

Alfalfa	Kikuyugrass
Bahiagrass	Knapweed
Bermudagrass	Milkweed
Bindweed, field	Muhly, wirestem
Bluegrass, Kentucky	Mullein, common
Blueweed, Texas	Napiergrass
Brackenfern	Nightshade, silverleaf
Broomsedge	Nutsedge (purple, yellow)
Clover	Orchardgrass
Dallisgrass	Pampasgrass
Dandelion	Quackgrass
Dock, curly	Ryegrass, perennial
Dogbane, hemp	Thistle, Canada

Fescue, tall
 Guineagrass
 Horsenettle
 Johnsongrass

Timothy
 Vaseygrass
 Wheatgrass, western

Woody Brush and Vines

Alder
 Ash
 Aspen, quaking
 Bearmat (Bearclover)
 Beech
 Birch
 Blackberry
 Broom (French, Scotch)
 Bracken
 Buckwheat, California
 Cascara
 Catsclaw
 Ceanothus
 Chamise
 Cherry
 Coyote brush
 Creeper, Virginia
 Dewberry
 Elderberry
 Elm
 Eucalyptus, bluegum
 Hasardia
 Hazel
 Holly, Florida
 Honeysuckle

Kudzu
 Locust, black
 Maple
 Monkey flower
 Oak
 Persimmon
 Poison Ivy
 Poison Oak
 Poplar, yellow
 Raspberry
 Rose, multiflora
 Sage, black
 Sagebrush, California
 Salmonberry
 Saltbrush, Sea myrtle
 Sassafras
 Sourwood
 Sumac
 Sweetgum
 Swordfern
 Tallowtree, Chinese
 Thimbleberry
 Tobacco, tree
 Trumpetcreeper
 Waxmyrtle
 Willow

SPRAYER CLEANUP

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

©MONSANTO COMPANY 1988

In case of emergency involving this product,
 Call Collect, day or night, (314) 694-4000.

PM 25

524-424

13 of 18

ACCEPTED

12

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for this pesticide registered under EPA Reg. No. 524-424

MON 14420

Keep out of reach of children
DANGER!

ACTIVE INGREDIENT:

Glyphosate [N-(phosphonomethyl)glycine] in the form of its monoammonium salt	74.7%
INERT INGREDIENTS	25.3%
	100.0%

NET 0.55 lb

MONSANTO COMPANY
ST LOUIS MISSOURI 63167

EPA Reg. No. 524
MAF-2982.01

print

594-424

11/2/78

MON 14420

Keep out of reach of children

DANGER!

ACTIVE INGREDIENT:

Glyphosate [N-(phosphonomethyl)glycine] in the	
form of its monoammonium salt	74.7%
INERT INGREDIENTS	25.3%
	100.0%

NET 1.1 lbs

MONSANTO COMPANY
ST LOUIS MISSOURI 63167

EPA Reg. No. 524
MAF-2982.02

HERBICIDE BY

Monsanto

WATER SOLUBLE GRANULE PREMEASURED IN WATER SOLUBLE BAG

For nonselective and selective weed control on industrial sites
such as roadsides and associated rights-of-way.

Carefully follow detailed instructions in label booklet.

Keep out of reach of children

DANGER!

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY", which appears on
the label booklet, before buying or using. If terms are
not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER!

CAUSES EYE BURNS
HARMFUL IF SWALLOWED.

Do not get in eyes, on skin or on clothing

Wear goggles or face shield and rubber gloves in mixing
and in adjusting, repairing or cleaning equipment.

Wash thoroughly with soap and water after handling

FIRST AID. IF IN EYES, immediately flush with plenty of water
for at least 15 minutes. Get medical attention. Remove
material from skin and clothing.

IF SWALLOWED, this product may cause gastrointestinal tract
irritation. Immediately dilute by swallowing water or
milk. Get medical attention.

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000

Environmental Hazards

Do not apply directly to water or wetland (swamps, bogs,
marshes or potholes). Do not contaminate water by disposal
of equipment washwaters

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and
applied only in stainless steel, aluminum, fiberglass,
plastic and plastic-lined steel containers

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF
THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT
STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or
spray solutions of this product react with such containers
and tanks to produce hydrogen gas which may form a highly
combustible gas mixture. This gas mixture could flash or
explode, causing serious personal injury, if ignited by open
flame, spark, welder's torch, lighted cigarette or other
ignition source

ACTIVE INGREDIENT

Glyphosate [N-(phosphonomethyl)glycine] in the
form of its potassium salt

INERT INGREDIENTS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any
manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food, or feed by storage or
disposal

STORAGE

Keep the outer bag **TIGHTLY** sealed to prevent moisture from
damaging any unused water soluble bags.

DISPOSAL

Wastes resulting from the use of this product that cannot be
used or chemically reprocessed should be disposed of in a
landfill approved for pesticide disposal or in accordance
with applicable Federal, state or local procedures.

Completely use water soluble pouches in application equipment
Then dispose of empty box and outer bag in a sanitary landfill,
or by incineration, or, if allowed by state or local authorities,
by burning. If burned, stay out of smoke

(See label booklet for Complete Directions for Use)

Product protected by U.S. Pat. Nos. 2,949,764 and
4,439,531. Other patents pending. No license granted
under any non-U.S. patent.

MON

14420

HERBICIDE BY **Monsanto**

WATER SOLUBLE GRANULE PREMEASURED IN WATER SOLUBLE BAG
for nonselective and selective weed control on industrial sites
such as roadsides and associated rights-of-way.

Carefully follow detailed instructions in label booklet

Keep out of reach of children

DANGER! Use precautions
on back panel

Read the entire label before using this product.
Use only according to label instructions

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or
using. If terms are not acceptable, return at once unopened

REFORMULATION OR REPACKAGING IS PROHIBITED

ACTIVE INGREDIENT

Glyphosate [N-(phosphonomethyl)glycine] in the
form of its monoammonium salt

INERT INGREDIENTS

74.7%

25.3%

100.0%



MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

NET 44 LBS

MON 14420

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER!

CAUSES EYE BURNS
HARMFUL IF SWALLOWED

Do not get in eyes, on skin or on clothing

Wear goggles or face shield and rubber gloves in mixing and when adjusting, repairing or cleaning equipment

Wash thoroughly with soap and water after handling

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention. Remove material from skin and clothing

IF SWALLOWED, this product may cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000

Environmental Hazards

Do not apply directly to water or wetland (swamps, bogs, marshes or potholes). Do not contaminate water by disposal of equipment washwaters

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal

STORAGE

Keep the outer bag TIGHTLY sealed to prevent moisture from damaging any unused water soluble bags

DISPOSAL

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures

Completely use water soluble pouches in application equipment. Then dispose of empty box and outer bag in a sanitary landfill, or by incineration, or, if allowed by state or local authorities, by burning. If burned, stay out of smoke

See label booklet for Complete Directions for Use

MON 14420

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described herein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well

as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit of Warranty and Liability which may not be varied by any verbal or written agreement.

® Registered trademark of Monsanto Company
MONSANTO COMPANY, 1988
EPA Reg. No. 524

Product protected by U.S. Pat. Nos. 3,799,758 and 4,405,531. Other patents pending. No license granted under any non U.S. patent.