

MON-8750

WATER SOLUBLE GRANULE
Herbicide by Monsanto

Complete Directions for Use

EPA Reg. No. 524-427

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

This product has been approved for use in California except as stated otherwise on pages XXX.

1991-1

A-MAP-2985

Read each of these sections of this label for essential product performance information.

USE THESE CONVENIENT SECTION MARKERS.

GENERAL INFORMATION

MIXING, ADDITIVES, AND APPLICATION
INSTRUCTIONS


WEEDS CONTROLLED

NONCROP USES

CROPPING SYSTEMS

CONSERVATION, MINIMUM AND NO-TILL SYSTEMS

TREE AND VINE CROPS



Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

SPINACH
SQUASH*** (Summer,
Winter)
SUGAR APPLE
TAMARIND
TOMATILO***
TOMATOES***†
TURNIP
WATERMELON***
WATERMELON SEEDS***
WHEAT
YAMS

- * Spot treatments may be applied in these crops.
- ** Do not treat rice fields ~~as leaves~~ when the fields contain flood water.
- *** Apply only prior to planting. Allow at least 3 days between application and planting.
- **** Do not feed or graze treated pineapple forage following application.
- † Use is restricted to direct seeded crops only.

When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 1/2 inch natural rainfall or by applying water via a sprinkler irrigation system.

NOTE: This product is not registered in California for use on atemoya, breadfruit, canistel, carambola, jaboticaba, jackfruit, longan, lychee, parsnips, passion fruit, sapodilla, sapote, sugar apples, tamarind, and yams.

Spot Treatment (Only those crops with "*" can be spot treated.) Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn, or boll opening on cotton.

For forage grasses and forage legumes see "Spot Treatment" in the "Pastures" section of "Cropping Systems" in this label.

For dilution and rates of application using boom^{....} or hand-held equipment, see "Mixing, Additives and Application Instructions" and "Weeds Controlled" sections of this label.

NOTE: FOR FORAGE GRASSES AND FORAGE LEGUMES, NO MORE THAN ONE-TENTH OF ANY ACRE SHOULD BE TREATED AT ONE TIME. FOR ALL OTHER CROPS, DO NOT TREAT MORE THAN 10 PERCENT OF THE TOTAL FIELD AREA TO BE HARVESTED.