

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup and this supplement.

ROUNDUP® Herbicide by Monsanto

EPA Reg. No. 524-445

FOR DISTRIBUTION AND USE ONLY WITHIN TEXAS AND CHAVES, DONA ANA, EDDY AND LUNA COUNTIES IN NEW MEXICO

ACCEPTED

Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, for the pesticide Roundup, EPA Reg. No. 524-445

FOR PREHARVEST APPLICATIONS TO COTTON.

Keep out of reach of children.

WARNING!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

*Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup for essential product performance information.

RECOMMENDATIONS

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label booklet for Roundup prior to the harvest of cotton.

This product may be applied using either aerial or ground spray equipment. For aerial applications, refer to the "APPLICATION EQUIPMENT AND TECHNIQUES" and "AERIAL EQUIPMENT" sections of the label booklet.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT IN EAST AND CENTRAL TEXAS AND IN NEW MEXICO OR 2 QUARTS PER ACRE IN WEST TEXAS¹ WHEN MAKING APPLICATIONS BY AIR.

NOTE: DO NOT APPLY TO CROPS GROWN FOR SEED.

Allow a minimum of 7 days between application and harvest. Do not feed or graze treated cotton forage or hay following preharvest applications.

WEED CONTROL

For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of the label booklet.

TANK MIXTURES

This product may be tank-mixed with the following harvest-aid products or combination of harvest-aid products:

**DEF[™] 6
Prep[™]**

**Folex[™]
Dropp[™]**

For application guidelines, precautions and use rates, refer to the DEF, Folex, Prep, and Dropp labels.

HARVEST AID

At rates of 24 ounces to 2 quarts per acre and when tank-mixed with Def, Folex or Dropp, this product may provide enhancement of cotton leaf drop and regrowth inhibition. The degree of leaf drop or regrowth inhibition achieved is strongly affected by plant and weather factors.

TIMING OF APPLICATIONS

Apply this product after 60 percent of the cotton bolls have opened, or when there are sufficient mature bolls to produce the desired yields. Care must be taken not to apply this product before a sufficient number of bolls has matured since further boll maturation and development will most likely NOT occur after application.

A boll is matured when it is too hard to be dented when squeezed between thumb and finger, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.

In the Texas counties listed below, apply this product at any time after October 1 since the boll maturation process has essentially stopped. Prior to October 1, follow the application instructions listed above. For optimum control of whiteweed, make applications two weeks prior to frost.

Armstrong, Bailey, Borden, Briscoe, Castro, Childress, Cochran, Collingsworth, Crosbey, Dawson, Deaf Smith, Dickens, Donley, Floyd, Gaines, Garza, Hale, Hall, Hockley, Kent, Lamb, Lubbock, Lynn, Motley, Parmer, Randall, Scurry, Swisher, Terry and Yoakum.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

† "West Texas" as used in this label is defined as the area bordered by Hwy. 86 on the north, Hwy. 287 on the northwest, Hwy. 283 on the east, Hwy. 190 on the south, Hwy. 285 on the southwest, and the Texas-New Mexico border on the west.

*Def is a trademark of Mobay Chemical Corporation.

*Folex and Prep are trademarks of Rhone-Poulenc, Inc.

*Dropp is a trademark of NOR-AM Chemical Company.

©Monsanto Company 1992

MAP

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup and this supplement.

ROUNDUP®

Herbicide by Monsanto

EPA Reg. No. 524-445

FOR NONSELECTIVE WEED CONTROL IN CHERIMOYA, COCOA BEANS, GENIP AND POMEGRANATES

Keep out of reach of children.

WARNING!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

*Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup for essential product performance information.

ACCEPTED



BEST AVAILABLE COPY

RECOMMENDATIONS

This product is recommended for weed control in established groves or orchards, or for site preparation prior to transplanting cherimoya, cocoa beans, genip or pomegranates. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, or orchard guns, or with wiper applicator equipment. See the "Application Equipment and Techniques" section of the label booklet for Roundup for specific information on the use of equipment.

When applying this product, refer to the "Weeds Controlled" and the "Tree and Vine Crops" sections of the label booklet for rates to be used.

NOTE: Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product. Do not apply more than 10.6 quarts of this product per acre per year.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING CUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Allow a minimum of 14 days between the last application and harvest.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©Monsanto Company 1992

MAP

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup and this supplement.

ROUNDUP® Herbicide by Monsanto

EPA Reg. No. 524-445



FOR USE IN WIPER APPLICATORS

Keep out of reach of children.

WARNING!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

*Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup herbicide for essential product performance information.

RECOMMENDATIONS

This product may be applied through a wiper applicator in any noncrop site and only when specifically recommended in cropping systems as stated in the label booklet for Roundup herbicide.

AVOID CONTACT WITH DESIRABLE VEGETATION.

Equipment must be designed, maintained, and operated to prevent the herbicide solution from contacting desirable vegetation.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

APPLICATION RATES - Use solutions consisting of 33 to 100 percent of this product. Limit solutions to 33 percent when using a standard wick-type applicator. Use the more concentrated solutions only with equipment that can handle a thick solution; such as the porous plastic-type applicators.

Add 1 quart of a nonionic surfactant for every 4 gallons of solution.

Read the "Limit of Warranty and Liability" in the label booklet for Honcho herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©Monsanto Company 1992

MAP

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167