

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR ROUNDUP® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup herbicide and this supplement.

Roundup
Herbicide by Monsanto

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-445

EPA Reg. No. 524-445

**FOR JOHNSONGRASS CONTROL POSTHARVEST TO CORN,
SORGHUM OR WHEAT IN TEXAS AND OKLAHOMA.**

Keep out of reach of children.

WARNING!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Roundup® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

See the "General Information" and "Mixing, Additives and Application Instructions" sections of the label booklet for essential product performance information.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from seed. Except as otherwise specified on the label, repeat treatments must be made before crop emergence in accordance with the instructions in the label.

Do not plant subsequent crops other than those listed in the label booklet for this product for 30 days following application.

CROPPING SYSTEMS

When applied as directed, under the condi-

tions described, this product controls the weed listed in the following cropping systems following the harvest of the crops:

**CORN (ALL)
SORGHUM (MILO)
WHEAT**

WEEDS CONTROLLED

When applied as recommended, under the conditions described, this product WILL CONTROL:

**Johnsongrass
*Sorghum halepense***

Apply 1.5 pints of this product plus 0.5 to 1 percent nonionic surfactant, plus 2 percent ammonium sulfate (17 pounds per 100 gallons of spray solution) in a total spray volume of 3 to 10 gallons of water per acre. Use 0.5 percent surfactant concentration when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration for those surfactants containing less than 70 percent active ingredient.

Applications may be made using ground or aerial application equipment. When applying by air, apply 3 to 5 gallons of water per acre. Flat fan nozzle tips (8002-8004) on 20 to 30-inch spacings are recommended for ground equipment. Ground applications should be made in 5 to 10 gallons of water per acre.

Ammonium sulfate should be predissolved and then poured through a 50-mesh strainer

when adding to the spray tank. If ammonium sulfate is added directly to the tank, add slowly and with good agitation. Ensure that ammonium sulfate is completely dissolved in the tank prior to the addition of this product and surfactant. Spray grade ammonium sulfate is much easier to dissolve than fertilizer grade material.

Following harvest, the field must be disked to cut up rhizomes and force new shoots. This product should be applied when johnsongrass is in the boot-to-head stage. Plants must be actively growing and not under drought stress. If heavy dew exists prior to spraying, delay application until the dew dries. Application should be made at least 7 days prior to frost. Allow 7 or more days after application before tillage.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

MAP-3079.01

© MONSANTO COMPANY 1992

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

