

524-445


06/13/2007

1/4

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR HONCHO HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Honcho herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



EPA Reg. No. 524-445

Roundup Ready and Honcho are trademarks of Monsanto Technology LLC

FOR APPLICATION TO CORN HYBRIDS WITH ROUNDUP READY® 2 TECHNOLOGY (INCLUDING ROUNDUP READY CORN 2 AND PRODUCTS DISPLAYING THE ROUNDUP READY 2 TECHNOLOGY LOGO)

Keep out of reach of children

CAUTION!

In case of emergency involving this product, call collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

These instructions are for use with corn hybrids with Roundup Ready 2 Technology, including Roundup Ready Corn 2 and products displaying the Roundup Ready 2 Technology logo, alone or in combination with other traits. Corn hybrids with Roundup Ready 2 Technology contain a patented glyphosate tolerance gene.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED IN THIS SUPPLEMENTAL LABEL), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION", "MIXING", "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" of the label booklet for Honcho herbicide for essential product performance information.

Corn Hybrids with Roundup Ready 2 Technology

Corn hybrids with Roundup Ready 2 Technology include Roundup Ready Corn 2 and products displaying the Roundup Ready 2 Technology logo.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (In-crop), Spot treatment, Preharvest, Post-Harvest

Annual Maximum Application Rates	
Combined total per year for all applications	8 quarts per acre
Preplant, At-Planting, Preemergence applications	5 quarts per acre
Total In-crop applications from emergence through 48 inches	3 quarts per acre
Maximum preharvest application (See PRECAUTIONS, RESTRICTIONS section for Preharvest)	1 quart per acre

GENERAL PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of the product label for general precautionary instructions for use of this product in Roundup Ready crops. See the "GENERAL INFORMATION" section of the product label for more information on "Annual Maximum Application Rates." Refer to individual tank mixture product label for restrictions and precautions, use according to the most restrictive precautionary statements for each product in the tank mixture.

Preplant, At-planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting.

TANK MIXTURES: This product may be tank mixed with the products listed below. Ensure that the specific product used is labeled for corn application. Read and follow label directions for all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to corn: 2,4-D, acetochlor, alachlor, atrazine, carfentrazone-ethyl, clopyralid, dicamba, diflufenzopyr, dimethenamid, dimethenamid-p, flufenacet, flumetsulam, flumiclorac pentyl ester, isoxaflutole, linuron, metolachlor, s-metolachlor, metribuzin, pendimethalin, rimsulfuron.

Aim, Axiom, Balance, Balance PRO, Banvel, Bicep MAGNUM, Bicep II MAGNUM, Bicep Lite II MAGNUM, Bullet, Cinch, Cinch ATZ, Clarity, Define, Degree, Degree Xtra, Distinct, Dual II MAGNUM, Epic, Frontier, FulTime, Guardsman MAX, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Keystone, Keystone LA, Lariat, Linex, Lorox, Marksman, Me-Too-Lachlor II, Micro-Tech, Prowl, Python, Python II, Radius, Resolve, Resource, Stalwart, Stalwart Xtra, Surpass, TopNotch]

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

USE INSTRUCTIONS: This product may be applied alone or in tank mixtures postemergence (in-crop) to corn hybrids with Roundup Ready 2 Technology, including Roundup Ready Corn 2 and products displaying the Roundup Ready 2 Technology logo. When applied as directed, this product controls annual grass and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. A postemergence application of this product should be made before the weeds reach a height of 4 inches or before they become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre should be made before the weeds reach a height of 4 inches.

This product may be broadcast over-the-top or applied with drop nozzles to corn hybrids with Roundup Ready 2 Technology from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first. Drop nozzles are recommended for optimum

spray coverage and weed control when corn height is 24 to 30 inches. For corn heights 30 to 48 inches (free standing), apply this product only by ground application using drop nozzles aligned to avoid spraying into the whorls of the corn plants.

TANK MIXTURES: This product may be tank-mixed with the following products. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to corn. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to corn: atrazine, acetochlor, alachlor, clopyralid, dicamba, diflufenzopyr, flumetsulam, flumiclorac pentyl ester, foramsulfuron, halosulfuron-methyl, iodosulfuron-methyl-sodium, mesotrione, nicosulfuron, rimsulfuron, thifensulfuron methyl.

Banvel, Basis, Basis Gold, Bullet, Callisto, Clarity, Degree, Degree Xtra, Distinct, Equip, Harness, Harness Xtra, Harness Xtra 5.6L, Hornet, Marksman, Micro-Tech, Option, Resolve, Resource, YUKON]

Tank Mix Partner	Maximum Height Of Corn For Application
Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6L	11 inches
Bullet* Micro-Tech*	5 inches
Atrazine	12 inches

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet® or Micro-Tech® herbicides.

PRECAUTIONS, RESTRICTIONS: Single in-crop applications of this product should not exceed 3 pints per acre. Sequential in-crop applications of this product from emergence through 48 inches in height must not exceed a total of 3 quarts per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product in-crop and harvest of corn forage or grain. (For applications at preharvest timing, see **Preharvest** section of this label). The use of additional surfactants and other additives, including fertilizers and micronutrients, in the spray solution are not recommended with this product and may result in increased potential for crop injury or reduced yield, and is not recommended for over-the-top applications, unless otherwise directed in supplemental labeling of fact sheets published separately by Monsanto.

Preharvest

USE INSTRUCTIONS: A single preharvest application of up to 32 fluid ounces per acre of this product may be applied for annual and perennial weed control prior to crop harvest. Make application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

PRECAUTIONS, RESTRICTIONS: Do not make a preharvest application of this product if more than a combined total of 64 fluid ounces of this product has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

Post-Harvest

USE INSTRUCTIONS: This product may be applied for weed control after corn harvest. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2, 4-D or dicamba may be used.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Honcho herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2007 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167