EXPEDITE●

GRASS & WEED PLUS RESIDUAL HERBICIDE

by Monsanto

11 -1 -- -- -- -- -- 1

FOR NONSELECTIVE POSTEMERGENCE AND RESIDUAL CONTROL

Contains glyphosate, the active ingredient in Roundup herbicide.

[The Roundup Logo]

Specially formulated for ExpediteTM Application Equipment.

Complete Directions for Use

EPA Reg. No. 524-449


Covers from 2,100 to 2,800 square feet.

©TMExpedite is a registered trademark of, and used under license from, Nomix Plc.

©Roundup is a registered trademark of Monsanto Company.

1993-1

875.15-400.05/53



OMONSANTO COMPANY 1993

1

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.
Read the entire label before using this product.

(COMPACT.LBL)

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED. This container may be refilled with this product om 1 gallon containers specifically labeled for refilling. Follow refilling instructions on the side panel of the 1 gallon container.

PRECAUTIONARY STATEMEN'S

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF INHALED.
MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.

Remove contaminated clothing and wash before reuse.

FIRST AID: IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of ana human health emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such

(COMPACT.LBL)



containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine

Contains petroleum distillates.

*Contains 125 grams per litre or 1.04 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt(Equivalent to 93 grams per litre or 0.77 pounds per U.S. gallon of the acid, glyphosate) and 120 grams per litre or 1.0 pounds per U.S. gallon of the active ingredient, oryzalin.

This product is protected by U.S. Patent No. 4,405,531. Other patents are pending. No license granted under any non-U.S. patent(s).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container

When container is completely emptied through Expedite app rection equipment or can no longer be refilled, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(COMPACT.LBL)

(10/30/90)

5,42.

1. 7 -10

General Information

This product is a mixture of a nonselective postemergence and a selective preemergence weed killer. Visible symptoms of postemergence control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

The preemergence portion of this product will remain stable on the soil surface up to 21 days. In the absence of timely rainfall for activation, irrigation can be used to move this product into the soil surface. If this product is not activated by rainfall or irrigation within 21 days of application, erratic preemergence weed control may result.

This product is recommended for commercial, industrial, noncropland and residential vegetation management in areas such as airports, christmas tree plantations, ditchbanks, farmsteads, fencerows, greenhouses, golf courses, highways and roadsides, industrial plant sites, institutional sites, municipalities, lawns, lumberyards, nurseries, nonbearing trees and vines, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites.

This product may be used as a directed application to control undesirable vegetation growing around established ornamentals, trees or shrubs. To avoid injury on desirable plants, prevent spray or drift contact with foliage, stems or green (immature) tree bark.

Do not apply this product where the likelihood of runoff onto lawn areas containing dichondra or cool-season turfgrass species exist as severe injury may occur.

Reseeding of desirable species should be delayed by 120 days following an application of this product.

Rainfall or irrigation within 2 hours after application may wash the product off plant foliage resulting in reduced post-emergent activity.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AYOID SPRAY CONTACT WITH THE FOLIAGE OF DESIRABLE TURF, CENAMENTALS, SHRUBS OR TREES. DO NOT APPLY OVER THE TOP OF DESIRABLE PLANTS.

DEST AVAILABLE COPT

ーッチンし

How to Apply

Apply this product as a uniform spray over the foliage of undesirable vegetation. Avoid applications during windy conditions, as spray drift or swath displacement may cause damage to any vegetation contacted. Use only Expedite application equipment when applying this product.

To select general flow rate settings for applying this product, use the Expedite Set-N-Spray settings listed in the following application guide. The letters followed by a number (B2A1...G+5) represent equipment settings for flow rates within the application range recommended for this product. If calibration is desired to obtain e-uipment settings, see the "HOW TO CALIBRATE" section of this laber.

A range of application rates are represented in the following application guide. Use the higher rate in the range on established perennial weeds or where longer residual control is needed. For best results, make applications before weeds are allowed to form a dense canopy.

Best results are obtained when the spinner is producing a uniform spray pattern. This is obtained by holding the lance head as close as practical to the top of the weed canopy(less than 4 inches) and increasing the spinner speed to obtain the correct swath width. The square spinner is recommended for applying this product.

EXPEDITE GRASS & WEED PLUS RESIDUAL HERBICIDE APPLICATION GUIDE

SET-N-SPRAY SETTING

	SPRAY			
SPINNER	SWATH	WALKING SPEED		
TYPE	WIDTH	1.5 MPH	2.25 MPH	
	<6"	B2 to B4	B5 to C2	
Spiral Spinner	6"	B5 to C3	C4• to 03	
	12"	D3 to E4	F1 to G3	
Square Spinner	18"	F1 to G3	G+ to *	

BEST AVAILABLE COPY


ر ۔۔ لزردن

The recommended rate range for this product represents 6 to 8 pounds total active ingredient or 3 to 4 gallons of product per acre.

*The flow rate through the equipment is not adequate to apply the recommended rate at this application rate.

How to Calibrate

- 1. Determine the proper Multi Flow units required fro the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for a 1.5 mph walking speed and a 12" swath width, 22 to 29 Multi Flow units are recommended.
- 2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held slightly lower than the product container.
- 3. Remove the spray disc and attach the Expedit calibration flow cup.
- 4. For the initial calibration check, set the equipment within the range of Set-N-Spray settings listed in the application guide in the "HOW TO APPLY" section. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
- 5. To remove the product from the calibration cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
- 6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.
- 7. When the desired head setting is obtained, remove the calibration cup and reattach the square spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.



(COMPACT.LBL)

(10/30/90)

下げつら

EXPEDITE® GRASS & WEED PLUS RESIDUAL HERBICIDE CALIBRATION GUIDE

30 SECOND MULTI FLOWTH MEASUREMENT*

SPINNER TYPE	SPRAY SWATE WIDTE	WALKING SPEED 1.5 MPH 2.25 MPH		
TIPE		1.5 FM 1		
	<6 *	7 to 10	11 to 14	
Spiral Spinner	6**	11 to 14	15 to 22	
	1.2"	22 to 29	32 to 43	
	18"	32 to 43	48 to 65**	
Square Spinner	24"	43 to 57**	64 to 86**	

The recommended rate range for this product represents 6 to 8 pounds total active ingredient or 3 to 4 gallons of product per acre.

*One Multi Flow equals 0.8 Milliliter.

**The flow rate through the equipment may not be adequate to apply the recommended rate at these walking speed and swath width combinations.

TMMulti Flow is a trademark of Nomix, Plc.

Cleaning of Equipment

The Expedite application equipment must be cleaned after each use and prior to storage. Use Expedite cleaner to remove residues of this product. To clean the lance, connect the . nce to the cleaner container, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner container and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. With extended use of this product, periodic cleaning of the lance head cap is recommended to prevent product buildup. Follow the procedure outlined in the equipment operating instructions for removing and replacing the Failure to properly clean the equipment muy result, in head cap. equipment failure or injury to subsequently treated desirable vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Empedite application equipment.

The disample tees

(1); -'>

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides POSTEMERGENCE control of many grasses and broadleaf weeds, including the following:

Bahiagrass Barnyardgrass Bentgrass Bermudagrass* Bindweed, field* Bluegrass, annual Brome, Downy Bromegrass, smooth Cheat Chickweed Clover: red, white Crabgrass Dallisgrass Dandelion Dock, curly Foxtail Geranium, Carolina Goosegrass Lambsquarters, common

Fescues Johnsongrass Lettuce, prickly Nutsedge: purple, yellow Mustard, tansy Orchardgrass Quackgrass Pennycress, field Pigweed: redroot, smooth Ragweed, common Ryegrass: annual, perennial Sandbur, field Shepherdspurse Sowthistle Stinkgrass Thistle, Canada Timothy Vaseygrass

*Partial control

When applied as recommended under the conditions described, this product provides PREEMERGENCE control of may broadleaf and grass weeds, including the following:

Barley, little
Barnyardgrass
Bittercress
Bluegrass, annual
Brachiaria
Carpetgrass
Chickweed, common
Crabgrass: large, smooth
Crowfootgrass
Cupgrass, southwestern
Fiddleneck, coast
Filaree: redstem, whitestem
Foxtail: bottlegrass, bristlegrass,

Morningglory*
Mustard: black, wild*
Nightshade, black*
Oat, wild
Panicum: fall; Texas
Pigweed: prostrate,
redroot; Tuugh;
spring, tumble
Punturevine
Purslane, common
Pusley, Florida
Ragweed, common*
Rocket, London

DEST AVAILABLE COPY

90:06

giant, green, robust, yellow
Groundsel, common
Henbit
Horseweed*
Johnsongrass(seedling only)
Junglerice
Knotweed, prostrate
Ladysthumb*
Lambsquarter
Lettuce, prickly*
Lovegrass: Mexican, orcutt
Mallow, common*
Milkweed, climbing*

Rockpurslane, desert
Ryegrass, annual
Sandbur, field
Shepherdspurse
Smartweed*
Sowthistle*
Sprangletop, red
Spurge, prostrate
Spurge, spotted*
Teaweed(Prickly sida)*
Velvetleaf*
Wheat, volunteer
Witchgrass
Woodsorrel, yellow

*This product provides suppression of these species.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURFJSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USFR OR BUYER, AND THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OF DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE, THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR

BEST AVAILABLE COPY

(COMPACT.LBL)

100-06

ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

875.15-400.05/53

EPA Reg. No. 524-449

)

CMONSANTO COMPANY 1993

|In case of ana human health emergency involving this product, Call Collect, | |day or night, (314) 694-4000.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(COMPACT.LBL)

10

(10/30/90)

ACCEPTED APR - 1 1993 Tidenci An or on it by Colored by



Keep out of reach of children.

CAUTION!

Please the entire laber before using this product

Use any according to take improcess

Read "LIMIT OF WARRANTY AND LIMBE/TY " before buying or using if learns are not acceptable, return at once uncomed

REFORMULATION IS MACHINITED

The container may be refilled with this product from 1 gaton containers solectically interest for refilling. Follow refilling instruction. In the sale femal of the 1 gaton container.

to case of an interpretty sheaking this product. Call Correct, call or hight (314) 694-4000.

AMOSANTO COMPANY
AMOSANTO COMPANY
AMOSANTO COMPANY
ST. COUS MASSIM SCHITTO ST.
ST. COUS MASSIM SCHITTO ST.
ST. COUS MASSIM SCHITTO ST.
ST. COUS MASSIM
ST. MASSIM ST. ST.
ST. MASSIM ST.
S

MONSANTO COMPANY 1991

EPA Pag No 574-449 EPA EU 343/377-001

Covers from 2,100 to 2,800 square feet

ACTIVE MIGREDIENT

Chambaile in Iphocutanomethyli glycine in the form of the soprocylamine sufficiently sometime in the form common in the formation of the soprocylamidation of the source o

75 e--

Contains personal assesses

Contains 125 grains per life (in 10 dounds per U.S. gallon of the active expresses, grains per life in the torm of its intomospharine sat (Equivalent to 90 grains per life or 0.77 pound per U.S. Callon of the active or 0.75 pound per U.S. Callon of the active act act 20 grains per little or 1.0 lbs. per U.S. gallon of the active indirected on popular.

The graph rule component of the product is protected by if S. Patrin No. 4.406 531 Other parents pending. No License 6 grands under any non-U.S. parents).

Monahaka is 5 registered trace— an of Monahaka Company

#ar: 1 of 3 875 15 400 14 53m*

NET 750 ML



PRECAUTIONARY STATEMENTS

is to Humans and Demants Arimais

Keep out of reach of children. CAUTION

CALI FLANT
HARMFUL IF SHIMLED,
MAY CAUSE RESPIRATORY TRACT IRRITATION,
Antid breathing vapor or spotly mist.
With these poly with state paid with the bandling.
Rathers continued classify and wish before more.
FIRST AID: F BONLED, approve believed to tresh air. Soutpredict direction if breathing difficulty devotes.

In case of a human health americanly invalving this product, Call Colect, day or reght, (\$14) disk-4600.

Environmental Hezarda

Do not apply directly to water, or to usest where surface water is present, or to extended areas paties the cross-legit water proof. On not constructed water when disposing all appropriat classing

Physical or Chanacal Hazards

Apply the product ONLY through Expedite" application reput most many will be as use .00 MOT STORE OR APPLY THIS PRODUCT IN MAY OTHER TYPE OF CONTAMER, PARTICULARLY GALYMAZE, STEEL OR LABOUR STEEL CONTAMERS OR STRAY TAMES THE I makes will such such assumers and trains as produce her-super gas which way seen a happy combacility gas receive. "Just gas microse could fluct, or outside, crassing service parasital largy of great by span fluxes, parts, values"s tools, bytest opposite or other syndrom stance.

CARECTIONS FOR USE

it is a weighter of Equipment that the special states are response accomplished with the importunity.

Storage or Disposal

Do not containwate water, laughthifts, field or sould by storage or femoral.

STOPAGE. Since the product between 12°F and 100°F in its original container and it a secured storage area.

Waster resident from the size of this product that cannot be used or chemically represented should be abspaced of in a tensible abspraced for postcode disposal of in accordance with applicable Federal, state or local proteodorus

Emplied detailer retains veptr and project residue. Chooses all liabilist adisquests seed operated to digitaryes.

When container to compatibly emplied grouph Expedite application and depress of its accessor tended, or by purchase depressed its accessor tended, or by individualism, or, if allowed by case and load authorities, by burney. If burned, stop out of annuls.

Since their place of the Countries to burney and their accessors and their contries of their countries.

(See taked beganing for Computers Securious for Alexa

LIBET OF WARRANTY AND LIABILITY

LIBIT OF WARRANTY AND LIABILITY

This Company various that their product gentures to the chaincal description on the lade and in naturality in the the proposes set
forth in the Company Conceptus (in the lade pooling ("Doctors")
what made in accordance with the time Decisions senter the conductors
described themes. NO OTHER EXPRESS SUMPRANTY OF MATLED
WARRANTY OF FITNESS FOR PARTICULAR PURPOSE ON MERCHANTARIE.TY IS MAKE. They wasterly is given subject to the coditions and invitames signed harms.

Beyor and all more study promptly notify this Company of any clume schedule beated in carefract, requirement, study lightly, other part or

they and all evers as responsible for all bass or damage trans one or handley which ear as term conditions beyond the carried of the Correlated by recompanishing with products which their flowers is forth on the Correlates, application to it contacts with described or yearney, understined the conditions which are earlier or recompanied flowers and the service or carried contacted flowers and the service or carried when the product as application pies and for the one. — "of when the product as application pages withing con." However, which we conditions earlied the machine capage specified in the Correlations, application in any informer and explicitly set forth with the Correlation, application in any informer and explicitly set forth on the Correlations, and the products of products other than these sets forth in the Correlations and on the sets, crop of twested originations.

VIBRORIAN.

THE ENCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIART OF THE LIARDLTY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, BULINES OR DOMANCES RESIALTING FROM THE USE OR HINDLEDGO THIS PRODUCT RIFLELY OF THE TORTH OR OTHERWISE) SHALL BE THE PLANCULTY, OTHER TORTH OR OTHERWISE) SHALL BE THE PLANCULTY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER LIER THE REPLACEMENT OF SUCH CUMMITTY, OR, F. HOT ACCUMENT SHALL THIS COMPANY OR ANY OTHER SELLER THE REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER THE REPLACEMENT.

Buyer and all wars are destruct to have accepted the terms of this LEMT OF WARRANTY AND LIABILITY which may not be varied by any varied or embar agreement.

875 15-400 (B/53



EXPEDITE® GRASS & WEED PLUS RESIDUAL HERBICIDE by Monsanto

FOR NONSELECTIVE POSTEMERGENCE AND RESIDUAL CONTROL

Contains glyphosate, the active ingredient in Roundup® herbicide.

[The Roundup Logo]

Specially formulated for ExpediteTM Application Equipment.

Complete Directions for Use

EPA Req. No. 524-449


Covers from 14,000 to 19,00010,600 to 14,100 square feet.

[®]TMExpedite is a registered trademark of, and used under license from, Nomix Plc.
[®]Roundup is a registered trademark of Monsanto Company.

1993-1

875.15.400.06/52

15 126



MONSANTO COMPANY 1993

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(SUPERPRO.LBL)

(10/22/90)

1

Read the entire label before using this product

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED. This package may be used to refill 750 ml containers that are specifically labeled for refilling with this product. Follow refilling instructions on side panel of this container.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF INHALED.
MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.

Remove contaminated clothing and wash before reuse.

FIRST AID: IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of ana human health emergency involving this product, Call Collect, day or night, (314) 694 1000.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertigal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STZEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such

(SUPERPRO.LBL)

(10/22/90)



containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

Contains petroleum distillates.

*Contains 125 grams per litre or 1.04 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt(Equivalent to 93 grams per litre or 0.77 pounds per U.S. gallon of the acid, glyphosate) and 120 grams per litre or 1.0 pounds per U.S. gallon of the active ingredient, oryzalin.

This product is protected by U.S. Patent No. 4,405,531. Other patents are pending. No license granted under any non-U.S. patent(s).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emptied through Expedite application equipment, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(SUPERPRO.LBL)

BEST AVAILABLE COPY

General Information

This product is a mixture of a nonselective postemergence and a selective preemergence weed killer. Visible symptoms of postemergence control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

The preemergence portion of this product will remain stable on the soil surface up to 21 days. In the absence of timely rainfall for activation, irrigation can be used to move this product into the soil surface. If this product is not activated by rainfall or irrigation within 21 days of application, erratic preemergence weed control may result.

This product is recommended for commercial, industrial, noncropland and residential vegetation management in areas such as airports, christmas tree plantations, ditchbanks, farmsteads, fencerows, greenhouses, golf courses, highways and roadsides, industrial plant sites, institutional sites, municipalities, lawns, lumberyards, nurseries, nonbearing trees and vines, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites.

This product may be used as a directed application to control undesirable vegetation growing around established ornamentals, trees or shrubs. To avoid injury on desirable plants, prevent spray or drift contact with foliage, stems or green (immature) tree bark.

Do not apply this product where the likelihood of runoff onto lawn areas containing dichondra or cool-season turfgrass species exist as severe injury may occur.

Reseeding of desirable species should be delayed by 120 days following an application of this product.

Rainfall or irrigation within 2 hours after application may wash the product off plant foliage resulting in reduced postemergence activity.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY CONTACT WITH THE FOLIAGE OF DESIRABLE TURF; URNAMENTALS, SHRUBS OR TREES. DO NOT APPLY OVER THE TOP OF DESIRABLE PLANTS.

(10/22/90)

(SUPERPRO.LBL)

How to Apply

Apply this product as a uniform spray over the foliage of undesirable vegetation. Avoid applications during windy conditions, as spray drift or swath displacement may cause damage to any vegetation contacted. Use only Expedite application equipment when applying this product.

To select general flow rate settings for applying this product, use the Expedite Set-N-Spray settings listed in the following application guide. The letters followed by a number (B2...G+) (A1...G5) represent equipment settings for flow rates within the application range recommended for this product. If calibration is desired to obtain equipment settings, see the "HOW TO CALIBRATE" section of this label.

A range of application rates are represented in the following application guide. Use the higher rate in the range on established perennial weeds or where longer residual control is needed. For best results, make applications before weeds are allowed to form a dense canopy.

When the desired head setting is obtained, install the spiner as indicated in the following guide. Best results are obtained when the spinner is producing a uniform spray pattern. This is obtained by holding the lance head as close as practical to the top of the weed canopy(less than 4 inches) and increasing the spinner speed to obtain the correct swath width. The square spinner is recommended for applying this product.

EXPEDITE GRASS & WEED PLUS RESIDUAL HERBICIDE APPLICATION GUIDE

SET-N-SPRAY SETTINGS

SPINNER TYPE	SPRÄY SWATH WALKING SPEE WIDTH 1.5 MPH 2.25 MPH	<u>D</u>	
Spiral Spinner	6" B2 to B4 B5 to C3		
• • • • • • • • • • • • • • • • • • •	12" C3 to D2 D3 to F2	•	
Square Spinner	18" D3 to P2 G4 to *	***	•

IEST AVARABLE COPT

1807631

*The flow rate through the equipment is not adequate to apply the recommended application rates at these walking speed and swath width combinations.

How to Calibrate

- 1. Determine the proper Multi Flow units required from the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for a 1.5 mph walking speed and a 12" swath width, 22 to 29 Multi Flow units are recommended.
- 2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held above the handle but below the level of the product container.
- 3. Remove the spray disc spinner and attach the Expedite calibration flow cup.
- 4. For the initial calibration check, set the equipment within the range of Set-N-Spray settings listed in the application guide in the "HOW TO APPLY" section. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
- 5. To remove the product from the calibration cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
- 6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.
- 7. When the desired head setting is obtained, remove the calibration cup and reattach the desired square spinner and follow the instructions in the "HOW TO APPLY section on how to spray this product.



19-1205

EXPEDITE GRASS & WEED PLUS RESIDUAL HERBICIDE CALIBRATION GUIDE

30 SECOND MULTI FLOWTH MEASUREMENT*

SPINNER TYPE

SPRAY SWATH

WALKING SPEED

6" 11 to 14 16 to 22

Spiral

Spinner 22 to 29 32 to 43

Square

32 to 43 48 to 65**

Spinner

The recommended rate range for this product represents 6 to 8 pounds total active ingredient or 3 to 4 gallons of product per

*One Multi Flow equals 0.8 Milliliter.

**The flow rate through the equipment may not be adequate to apply the recommended rates at these walking speed and swath width combinations.

TMMulti Flow is a trademark of Nomix, Plc.

Cleaning of Equipment

The Expedite application equipment must be cleaned after each use and prior to storage. Use Expedite cleaner to remove residues of this product. To clean the lance, connect the lance to the cleaner box, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner box and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. With extended use of this product, periodic cleaning of the lance head cap is recommended to prevent product buildup. Follow the procedure outlined in the equipment operating instructions for removing and replacing the head cap. Failure to properly clean the equipment may result in equipment failure or injury to subsequently treated desirable vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Expedite application equipment.

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides POSTEMERGENCE control of many grasses and broadleaf weeds, including the following:

Bahiagrass Barnyardgrass **Bentgrass** Bermudagrass* Bindweed, field* Bluegrass, annual Brome, Downy Bromegrass, smooth Cheat Chickweed Clover: red, white Crabgrass Dallisgrass Dandelion Dock, curly Fortail Geranium, Carolina Goosegrass Lambsquarters, common

Fescues Johnsongrass Lettuce, prickly Nutsedge: purple, yellow Mustard, tansy Orchardgrass Quackgrass Pennycress, field Pigweed: redroot, smooth Ragweed, common Ryegrass: annual, perennial Sandbur, field Shepherdspurse Sowthistle ... Stinkgrass: :: Thistle, Canada Timothy Vaseygrass

*Partial control

LEST AMAILABLE CONS

(SUPERPRO.LBL)

(10/22/90)

2000

210101

When applied as recommended under the conditions described, this product provides PREEMERGENCE control of many broadleaf and grass weeds, including the following:

Barley, little Barnyardgrass Bittercress Bluegrass, annual Brachiaria Carpetgrass Chickweed, common Crabgrass: large, smooth Crowfootgrass Cupgrass, southwestern Fiddleneck, coast Filaree: redstem, whitestem Foxtail: bottlegrass, bristlegrass, giant, green, robust, yellow Groundsel, common Henbit Horseweed* Johnsongrass(seedling only) Junglerice Knotweed, prostrate Ladysthumb* Lambsquarter Lettuce, prickly* Lovegrass: Mexican, orcutt Mallow, common* Milkweed climbing*

Morningglory* Mustard: black, wild* Nightshad:, black* Oat, wild Panicum: fall, Texas Pigweed: prostrate, redroot, rough, spring, tumble Puncturevine Purslane, common Pusley, Florida Ragweed, common* Rocket, London Rockpurslane, desert Ryegrass, annual Sandbur, field Shepherdspurse Smartweed* Sowthistle* Sprangletop, red Spurge, prostrate Spurge, spotted* Teaweed(Prickly sida) * Velvetleaf* Wheat, volunteer Witchgrass Woodsorrel, yellow

*This product provides suppression of these species.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

875.15-400.06/53



スー・ノディーク

J- 200

EPA Reg. No. 524-449

ONONSANTO COMPANY 1993

|In case of ana numer health emergency involving this product, Call Collect, | |day or night, (314) 694-4000.

MONSANTO COMPANY AGRICULTURAL PRODUCTS ST. LOUIS, MISSOURI 63167 U.S.A.

(10/72/90)

ACCEPTED

APR - 1 1993

Underthe federal imperious, furnities and Rodenticide Act, as smerciaed, for the professe registered order Elia Pegallia 524-449



RESIDUAL HERBICIDE

by Monsanto

FOR NONSELECTIVE A STEMERGENCE AND RESIDUAL CONTROL

Specially formulated for Expedite® Application Equipment

Covers from 10,600 to 14,100 square feet

Keep out of reach of children.

CAUTION! (Sur precadent on back party Read the entire label before using this product.

Use only according to label instructions.


Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened

REFORMALATION IS PROHIBITED.

This package may be used to refit 750 ml containers that are specifically lebeted for refilling with this product. Follow refilling instructions on side panel of this container.

In case of a human health emergency involving this product, Call Collect, day or night, (314) 694-4000.

MONSANTO COMPANY AGNICULTURAL PRODUCTS ST. LOUIS, MISSOURI 63167 U.S.A. A PEXPENSE IS A Processory of, and used under Scenes from, Nomes Ptc. © MONSANTO COMPANY 1991 EPA F89 No 524-449 EPA Est 34313-TX-001



Glyphosate, N-(phosphonomethyl) glycine in the form of its isopropylamine salt 12.4%

Contains petroleum distillate:

"Contains 125 grams per litre or). 04 pounds per U.S. gallon of the active ingredient, gyphosate, in the form of its isopropylamine sait. (Equivalent to 93 grams per litre or 0.77 pound per U.S. gallon of the acid, glyphosate) and 120 grams per litre or 1.0 lbs. per U.S. gallon of the active ingredient, oryzalin.

The glyphosate component of this product is protected by U.S. Patent No. 4,405.531. Other patents pending. No License is granted under any nr., LU.S. patent(s).

This product contains physhosale, the active ingradient in Roundup* herbicide. Afficiently is a registered tradement of Moneanto Company.

Part 1 of 2 875 15-400.14/53HT

GALLON NET 1

iest available copy

GRASS & WEED PLUS RESIDUAL HERBICIDE by Monsanto

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF INHALED.
MAY CAUSE RESPIRATORY TRACT IRRITATION.
Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.
Remove contaminated clothing and wash before reuse.
FIRST AID: IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of a liuman health emergency involving this product, Call Collect, day or night, (314) 894-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product Or*LY through Expedite** application equipment designed for its \(\text{se.} DO \) NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR (INLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers \(\text{...} \) at tanks to produce hydrogen gas which may form a highly combustible gamedure. This gas modure could flash or explode, causing senous personal injury if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIO'IS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage or Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°C \Rightarrow its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a tandilli approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emplied, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(See label booldet for Complete Directions for Use)

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACCUMED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.


Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

Part 2 of 2


875.15-400.08/53




AS A BACKPACK



Insert Nomix Link-Up cassette into the Compact




Insert Big Five into backpack, Connect to herbicide pack by aligning groove of pipe to ridges of Big Five and twist through 90° clockwise




Raise Compact above backback.

Depress Compact trigger to drain nerbicide through lance and into Big Five.




Unlock by twisbing anti-clockwise through 90° and disconnect.


TO REFILL




Align grooves of the Compact cartridge to ridges of the Big Five refill pack.



Twist the cartridge clockwise through 90° to lock the two containers and open valves



Invert, ensuring the Big Five pack is uppermost and allow the cartridge to refill.



Unlock the packs by twisting anti-clockwise through 90° and disconnect.

Keep a cloth handy to write the nozzle of the refilled cartridge.

NO. Empty cartridges smooth only be ratified for one season.

Enough optimizing any only rating over the same numberals as arrangely appears.