10 25




Specially terminoled for ExpediteTM Application Equipment

2, 900 205, 700 Covers from 14,000 to 19,000 square feet

Keep out of reach of children. CAUTION!

PRECAUTIONARY STATEMENTS



NET 750 ML

ACCEPTED

Under the Federal Insecticide, Fungicide, and Redenticide Act, as one made, for the posticide required under EPA Reg. No. 500 - 1/5 60

BEST AVAILABLE COPY

(Cordinate from four pared)

Fortune of Paris

Do not apply directly to reduce or resident (presence, being, maryers, and particular, Do not correspond excess when

Provided or Chambridge Habitation

Apply this product ORLY through Expedite application objections designate the sun, DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GRAVANIZED STEEL OR UNLINED STEEL, CONTAINER ON SPREY, TANKE, This product will reset with such containers and tarks to produce in product any most which may larm a rapidy communities gas medium. This gas within may larm a rapidy communities gas medium this gas within may larm a reprint order to gas medium through the or expects, making survival parameter view, if sprides the question, sperit, mostor's larm, digital disparate or other medium termine.

DIRECTIONS FOR USE

a to a variation of Federal law to use this product in May

Storage and Disposal

Co not contempe with feetballs, had to load by Parago or Philippal.

STORAGE:

Store this product between 20° F and 100° F in its original coreases are in a secured storage area.

UISPOSAL

Analog resulting from the sale of this product that cannot be used or chemically reprocursed should be disposed of this sample approved by posterior disposed of the carried appear of the contribution will analogue features that or local structures.

Emphod contigues intomo vapos and product residue.

Claserus all lacerus safeguines until container is destroyed.

Co net reuse container.

when contains a completely emptine Triming Expends application southering, purchase and Majorie of this settling unique or by increasing, or 5 allowed by likes and food

(See year boolest for Company Direction for Use)

CARE OF WARRANTY AND CARE.

This Company various that this product ancients to the purposes on facts to the Company Constantly III for the purposes on facts to the Company Constants for the shell bested (Circultura) when send to accordance with Peac Directions sends the sendings described therein, and OTHES DIVERSE MINISTERY OR IMPLIED WARRANTY OF PRINCIPLE FOR PRINCIPLES WARRANTY OR IMPLIED WARRANTY OF PRINCIPLE FOR PRINCIPLES WARRANTY OR IMPLIED WARRANTY OF PRINCIPLE SACCE, This receively is also excluded in the considera and

Buyer and all seven shall promptly notify this Company of any department based in content, maphymen, and basely, other half or otherwise.

Buyer and all soons are instantiable for all loss or disregaligations or humiling which results from considering buyers. The extend of Jun Company, including, but not britted its, humilingshiftly with products other than those on facts in the Developing, applicables to be original with developing hypothesis, urusual vession, vession completely with any autoria the urusual vession, vession completely and autoria the time partial select the product is applicables and on the fine lines partial velocity to product is applicables active and are time partially selected by applicables recipies on the large in the (Developing, receiving constitutes autoria the matches report parallel on the Circumstant, or the processes of produces other than these out tops in the Chemistrus to or an

THE EXCLUSIVE REMEDY OF THE WIER OR BLYER, AND THE LIBIT OF THE LIMBLITY OF THIS COMMANY OR ANY OTHER SELER FOR MANY AND ALL LOBBER, RULBERS OR CHANGES OR CHANGES OR CHANGES RESULTING FROM THE USE ON HANDLING OF THIS PRODUCT (RECLUSIONS CLAMES BASED IN CONTRACT, HE GUIGH NCE, STRICT LIMBLITY, OTHER TOTAL OR OTHER HANDLING FROM THE USER OR GUYER FOR THE GUANTITY OF THIS BYOUNCE FOR THE COMMANY OR MAY OTHER SELER THE REPLACEMENT OF SUCH QUANTITY, OF IF MOY ACCUMED BY PARCHASE, REPLACEMENT OF SUCH QUANTITY IN MO EYE HT SHALL THIS COMMANY OR ANY OTHER SELER.

Super and all waters are decrease to find accepted the forms or part (and OF MANPANIET AND LIABILITY which may not be consisted as a second to the consistency of the

HONSANTO COMPANY AGRICULTURAL PRODUCTS

EPA Est. 34313 TH-491

Part 2 of 2

BEST AVAILABLE COPY





Specially formulated for Expedite Tal Application Equipment

Keep out of reach of children. CAUTION!

Read the entire label before using this product. Use or ly according to tabel instructions.
Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using, if terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTIONI

CAUSES EYE IRRITATION. HARMFUL IF INHALED. MAY CAUSE RESPIRATORY TRACT IRRITATION. Avoid contact with eyes, sidn or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling.

FIRST AID. IF IN EYES, immediately flush with plenty of water. Get medical attention. IF INHALED, remove individual to fresh air. Seek medical attention If breathing difficulty develops.

(Continued on back panel)

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000,

Covers from 19,000 to 38,000 square feet

ACTIVE INGREDIENT: Glyphosale, N-(phosphonomethyl)glycine in the form of its leopropylamine

84 2% INERT INGREDIENTS:..... 100.0%

Contains petroleum distillates.

*Contains 162 grams per litre or 1.35 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 120 grams per litre or 1.0 pound per U.S. gallon of the acid, glyphosale.

This product is protected by U.S. Patents No. 3,799,758, No. 4,405,531 and No. 4,840,650, Other patents pending. No Hoense granted under any non-U.S. patent(s).

 $^{\mbox{\scriptsize TM}}$ Expedite is a trademark of, and used under license from, Nombs Pfc.

This product contains glyphosale, the active ingredient in Roundup herbicide.

Roundup is a registered trademark of Monsanio Companys

MONSANTO COMPANY 1990

• EBA Beg. No. 584-XXXX

Part 1 062 875.85-300.02/53HP

•••••

NET 1.32 GALLONS (1 GAL., 1 QT., 9 FL.OZ.

MCCEPIED tom Tark Colonia Tark Tark Colonia

BEST AVAILABLE COPY

Environmental Hazards

Do not apply directly to water or watend (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and fanite to produce hydrogen gas which may form a highly combustible gas middure. This gas middure could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's forch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

Il is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodsluffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32° F and 100° F in its original container and in a secured storage area.

DISPOSAL

Wastes resulting from the use of this product that cannot be i sed or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container relains vapor and product residue.

Observe all labeled safeguards until container is destroyed.

Do not reuse container.

When container is completely emptied through Expedite application equipment, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning, it burned, stay out of smoke.

(See label booklet for Complete Direction for Use.)

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for use label booldst ("Directions") when used in accordance with those Directions under the conditions describes therein. NO OTHER EXPRESS WARRANTY OR INFLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all uters are responsible for all loss or demage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions witch are outside the range considered normer at the application site and for the time period when the product is application alle and for the time period when the product is application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LLABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OF IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SCILLER BE LIABLE FOR ANY INCIDENTAL OR SPECIAL DAMAGES.

Buyer and all users are deemedselecture accepted the terms of this LIMIT OF WARRANTY AND LUBBILITY which may not be varied by any verbal or writen-agreement.

MONSANTO COMPANY, AGRICULTURAL PRODUCTS. ST. LOUIS, MISSOURI 63167 USA

EPA Est. 524-34313-TX-0012

875.13-367.03/53HT

....

GRASS & WEED II HERBICIDE by Monsanto

For Non-Selective Postemergence Control of Unwanted Vegetation

Contains glyphosate, the active ingredient in Roundup herbicide.

[The Roundup Logo]

Specially formulated for Expedite Application Equipment.

Complete Directions for Use

EPA Reg. No. 524-XXX

Covers from 19,000 to 38,000 square feet.

^{® IM}Expedite is a registered trademark of, and used under license from, Nomix Plc.
[®]Roundup is a registered trademark of Monsanto Company.

1990-1

xxx.xx.xxx.xx/xx

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 450

©MONSANTO COMPANY 1990

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(SUPERPRO.LBL)

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

HARMFUL IF INHALED.

MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid contact with eyes, skin or clothing.

Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water or wetland (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine

Contains petroleum distillates.

*Contains 162 grams per litre or 1.35 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 120 grams per litre or 1.0 pound per U.S. gallon of the acid, glyphosate).

This product is protected by U.S. Patent No. 3,799,758, U.S. Patent No. 4,405,531 and U.S. Patent No. 4,840,650. Other patents are pending.

No license granted under any non-U.S. patent(s).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emptied through Expedite application equipment, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a nonselective weed killer that is applied, using Expedite application equipment, to the foliage of actively growing weeds. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms of control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

This product is recommended for commercial, forestry, industrial, noncropland and residential vegetation management in areas such as airports, ditchbanks, farmsteads, fencerows, forestry and Christmas plantations, greenhouses, golf courses, highways industrial roadsides, plant sites, institutional sites. municipalities, lawns, lumberyards, nurseries, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites. This product is also recommended for weed control in agricultural crops as specified in the "CROPPING SYSTEMS" section of this label.

When this product is used as a directed application to control undesirable vegetation growing around established ornamentals, trees, shrubs or vines, avoid spray or drift contact with foliage, stems or green(immature) tree bark.

Rainfall or irrigation within 2 hours after application may wash the product off plant foliage resulting in reduced weed control. Delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application to allow product to move within the plant for the most effective control.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY CONTACT WITH THE FOLIAGE OF DESIRABLE TURF, ORNAMENTALS, SHRUBS OR TREES. DO NOT APPLY OVER THE TOP OF DESIRABLE PLANTS.

How to Apply

Apply this product as a uniform spray over the foliage of undesirable vegetation. Avoid applications during windy conditions, as spray drift or swath displacement may cause damage to any vegetation contacted. Use only Expedite application equipment when applying this product.

When us a in greenhouses, turn off air circulation fans while spraying print of this product.

BY SPRAY PRIFT OF THIS PRODUCT.

To select general flow rate settings for applying this product, use the Expedite Set-N-Spray settings listed in the following application guide. The letters followed by a number (Al...F1) represent equipment settings for flow rates within the application range recommended for this product. If calibration is desired to obtain equipment settings, see the "HOW TO CALIBRATE" section of this label. Calibration is recommended where more precise equipment calibration is desired, or when the application rate needs to be customized for the type of weeds present.

When the desired head setting is obtained, install the spinner as specified in the following guide. For best results, make applications before weeds are allowed to form a dense canopy. Best results are obtained when the spinner is producing a uniform spray pattern. For spray widths less than 15 inches, this is obtained by holding the lance head as close as practical to the top of the weed canopy(less than 4 inches) and increasing the spinner speed to obtain the correct swath width.

EXPEDITE GRASS & WEED II HERBICIDE APPLICATION GUIDE

SET-N-SPRAY SETTING

SPINNER TYPE	SPRAY SWATH WIDTH	1.5 MPH	WALKIN 2.0 MPH	NG SPEED 2.5 MPH	3.0 MPH
	<6 "	A1	A1	A1	A2
	6"	A1	A1	A2	А3
	9"	A2	A3	A4	A 5
	12"	А3	A5	B1	В3

Square						_
Spinner	15"	A4	В1	В3	B4	
	18"	A5	В3	B4	C1	
	21"	B1	B4	C1	СЗ	_
Round	24"	В3	B 5	С3	C5	_ _ _
Spinner	36"	C1	C5	D5	F1	

The recommended rate in this guide represents 2.25 pounds acid equivalent glyphosate or 2.25 gallons of product per acre.

How to Calibrate

- 1. Determine the proper Multi Flow units required from the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for a 1.5 mph walking speed and a 12" swath width, 11 to 22 Multi Flow units are recommended. Use the lowest setting for easy-to-control weeds such as small annuals, and the highest setting for difficult-to-control weeds such as bermudagrass and field bindweed.
- 2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held above the handle but below the level of the product container.
- 3. Remove the spray disc and attach the Expedite calibration flow cup.
- 4. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
- 5. To remove the product from the calibration cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
- 6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.

7. When the desired head setting is obtained, remove the calibration cup and reattach the desired spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.

EXPEDITE GRASS & WEED II HERBICIDE CALIBRATION GUIDE

30 SECOND MULTI PLOWTH MEASUREMENT*

SPINNER Type	SPRAY SWATH						
	WIDTH	1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH		
Square Spinner	<6"	4 to 7	5 to 10	6 to 12	7 to 14		
	6"	5 to 11	7 to 14	9 to 18	11 to 22		
	9"	8 to 16	11 to 22	13 to 27	16 to 32		
	12"	11 to 22	14 to 28	18 to 36	22 to 43		
	15"	13 to 27	18 to 36	22 to 45	27 to 54		
	18"	16 to 32	22 to 43	27 to 54	32 to 65		
	21"	19 to 38	25 to 50	31 to 63	38 to 75		
Round	24#	22 to 43	29 to 57	36 to 72	43 to 86		
Spinner	36"	32 to 65	43 to 86	54 to 108	65 to 129		

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre. *One Multi Flow equals 0.8 milliliter.

IMMulti Flow is a trademark of Nomix, Plc.

CROPPING SYSTEMS

For crops listed in this section of this label, this product may be used as directed prior to the emergence of direct seeded crops, prior to transplanting of crops, in established groves, vineyards or orchards and for spot treatments as specified. For crops not listed, planting must be delayed for 30 days following an application.

This product does not provide residual weed control. For subsequent weed control, follow a program using residual herbicides or us repeat applications of this product. When using repeat applications of this product, do not exceed 2 applications per year in cropping systems or 3 applications per year in established groves, vineyards or orchards.

Unless otherwise directed in this label, do not harvest or feed treated crops for 8 weeks after application of this product. Allow 14 days following spot treatments before grazing domestic livestock or harvesting forage grasses and legumes.

When using this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 1/2 inch of rainfall or irrigation.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURE BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

FIELD CROPS

This product can be used prior to the emergence of direct seeded crops or prior to transplanting of crops in the following list.

ALFALFA*
ARTICHOKE,
JERUSALEM
ASPARAGUS
ATEMOYA
BARLEY*
BEANS(ALL)
BEET GREENS
BEETS(RED, SUGAR)
BLACKBERRY
BLUEBERRY
BOYSENBERRY

KALE
LENTILS
LETTUCE
LOGANBERRY
MELONS***
MUSTARD GREENS
OATS*
OKRA
OLALLIEBERRY
ONION
PARSNIPS
PEANUTS

11 7 15

BROCCOLI CABBAGE CARAMBOLA CARROT CAULIFLOWER CELERY CHICORY CORN(ALL) * COTTON* CRANBERRY CUCUMBER*** CURRANT LEWBERRY **ELDERBERRY** FORAGE GRASSES* FORAGE LEGUMES* GARLIC*** GOOSEBERRY

GOURDS***

HORSERADISH

HUCKLEBERRY

PEAS(ALL) PEPPER*** PINEAPPLE**** POTATO(IRISH, SWEET) PUMPKIN*** RADISH RASPBERRY (BLACK, RED) RICE** RUTABAGA SORGHUM (MILO) * SOYBEANS* SPINACH SQUASH***(SUMMER, WINTER) SUGAR APPLE TOMATILO*** TOMATOES***+ TURNIPS WATERMELONS*** WATERCRESS*** WHEAT* YAMS

*Spot treatments may be applied in these crops and must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn and boll opening on cotton

**Do not treat rice fields or levees when the fields contain flood water.

***Apply only prior to planting. Allow at least 3 days between application and planting.

****Do not feed or graze treated pineapple forage following application.

+Use is restricted to direct seeded crops only.

NOTE: This product is not registered in California for use on atemoya, carambola, sugar apples, yams and parsnips.

TREE CROPS

This product is recommended for weed control in established groves, vineyards, or orchards, or for site preparation prior to transplanting these crops as directed below.

CITRUS*: citron, grapefruit, kumquat, lemon, lime, orange, tangelo, tangerine.

NUTS**: almond, filbert, macadamia, pecan, pistachio, walnut.

POME FRUIT*: apple, pear.

STONE FRUIT***: apricots, cherries, nectarines, olives, peaches, plums/prunes. Remove suckers and low hanging limbs at least 10 days prior to application. Avoid application near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. Cherries may be treated with this product in Apricots, nectarines, peaches and plums/prunes all states. may only be treated in the states of Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington. If a spray shield is used and the application is made no later than 90 days after first bloom, this product may be used on peaches in the states of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee. EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED BY THIS PRODUCT.

TROPICAL FRUIT: acerola, atemoya*, avocado*, banana(plantains), carambola*, coffee****, figs*, guava, mango*, papaya, sugar apple*, tea. Allow a minimum of 1 day between last application and harvest of guava and papaya. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

- *Allow a minimum of 14 days between last application and harvest.
- **Allow a minimum of 21 days between last application and harvest.
- ***Allow a minimum of 17 days between last application and harvest.
- ****Allow a minimum of 28 days between last application and harvest.

VINE CROPS

KIWI FRUIT*

GRAPES:* Any variety of table, wine, or raisin grape may be treated with this product. In the Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

*Applications should not be made when green shoots, canes, or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest.

Cleaning of Equipment

The Expedite application equipment must be cleaned after each use and prior to storage. Use Expedite cleaner to remove residues of this product. To clean the lance, connect the lance to the cleaner box, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner box and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. Failure to properly clean the equipment may result in equipment failure or injury to subsequently treated desirable vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Expedite application equipment.

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides control of many grasses and broadleaf weeds, including the following:

Bahiagrass Barnyardgrass Bentgrass Bermudagrass* Bindweed, field* Bluegrass, annual Brome, downy Bromegrass, smooth Cheat Chickweed Clover: red, white Crabgrass Dallisgrass Dandelion Dock, curly Foxtail Geranium, Carolina Goosegrass Guineagrass Lambsquarters, common

Fescues Johnsongrass Lettuce, prickly Mustard, tansy Nutsedge: purple, yellow Orchardgrass Quackgrass Pennycress, field Pigweed: redroot, smooth Ragweed, common Ryegrass: annual, perennial Sandbur, field Shepherdspurse Sowthistle Stinkgrass Thistle, Canada Timothy Vaseygrass

^{*}Partial control

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH CUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

xxx.xx-xxx.xx/xx

EPA Reg. No. 524-XXX

ONONSANTO COMPANY 1990

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4070.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.