

PM25 524-450

10432

**EXPEDITE®
GRASS & WEED II HERBICIDE
by Monsanto**



**For Non-Selective Postemergence Control of Unwanted Vegetation
Contains glyphosate, the active ingredient in Roundup® herbicide.**

[The Roundup Logo]

Specially formulated for Expedite™ Superpro Application Equipment.

Complete Directions for Use

EPA Reg. No. 524-450

Covers from 14,000 to 29,000 square feet.

®™ Expedite is a registered trademark of, and used under license from, Nomix Plc.

® Roundup is a registered trademark of Monsanto Company.

1992-13-1

875.15.300.xx/53

ACCEPTED
APR - 1 1993
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 524-450

©MONSANTO COMPANY 1992

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(SUPERPRO.LBL)

1

(11/21/90)

20732

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

~~REFORMULATION OR REPACKAGING IS PROHIBITED.~~

~~RECALL INFORMATION: CONTACT THE MANUFACTURER AT THE FOLLOWING ADDRESS: [REDACTED]~~

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

HARMFUL IF INHALED.

MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid contact with eyes, skin or clothing.

BEST AVAILABLE COPY

Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of ~~an~~ human health emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

20732

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine
in the form of its isopropylamine salt15.8 %

INERT INGREDIENTS:.....84.2 %
100.0 %

Contains petroleum distillates.

*Contains 162 grams per litre or 1.35 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 120 grams per litre or 1.0 pound per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531. Other patents are pending.

No license granted under any non-U.S. patent(s).

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emptied through ~~expedite application equipment~~, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a nonselective weed killer that is applied, using Expedite application equipment, to the foliage of actively growing weeds. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms of control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

This product is recommended for commercial, forestry, industrial, noncropland and residential vegetation management in areas such as airports, ditchbanks, farmsteads, fencerows, forestry and Christmas tree plantations, greenhouses, golf courses, highways and roadsides, industrial plant sites, institutional sites, municipalities, lawns, lumberyards, nurseries, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites. This product is also recommended for weed control in agricultural crops as specified in the "CROPPING SYSTEMS" section of this label.

When this product is used as a directed application to control undesirable vegetation growing around established ornamentals, trees, shrubs or vines, avoid spray or drift contact with foliage, stems or green (immature) tree bark.

Rainfall or irrigation within 2 hours after application may wash the product off plant foliage resulting in reduced weed control. Delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application to allow product to move within the plant for the most effective control.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY CONTACT WITH THE FOLIAGE OF DESIRABLE TURF, ORNAMENTALS, SHRUBS OR TREES. DO NOT APPLY OVER THE TOP OF DESIRABLE PLANTS.



52/32

How to Apply

Apply this product as a uniform spray over the foliage of undesirable vegetation. Avoid applications during windy conditions, as spray drift or swath displacement may cause damage to any vegetation contacted. Use only Expedite application equipment when applying this product.

When used in greenhouses, turn off air circulation fans while spraying. **DESIRABLE GREENHOUSE PLANTS WILL BE INJURED IF CONTACTED BY SPRAY OR DRIFT OF THIS PRODUCT.**

To select general flow rate settings for applying this product, use the Expedite Set-N-Spray settings listed in the following application guide. The letters followed by a number (A1...F1) represent equipment settings for flow rates within the application range recommended for this product. If calibration is desired to obtain equipment settings, see the "HOW TO CALIBRATE" section of this label. Calibration is recommended where more precise equipment calibration is desired, or when the application rate needs to be customized for the type of weeds present.

When the desired head setting is obtained, install the spinner as specified in the following guide. For best results, make applications before weeds are allowed to form a dense canopy. Best results are obtained when the spinner is producing a uniform spray pattern. For spray widths less than 15 inches, this is obtained by holding the lance head as close as practical to the top of the weed canopy (less than 4 inches) and increasing the spinner speed to obtain the correct swath width.

EXPEDITE® GRASS & WEED II HERBICIDE APPLICATION GUIDE

SET-N-SPRAY SETTING

SPINNER TYPE	SPRAY SWATH WIDTH				WALKING SPEED			
	1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH	1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH
<6"	A1	A1	A1	A2				
6"	A1	A1	A2	A3				
9"	A2	A3	A4	A5				
12"	A3	A5	B1	B3				

BEST AVAILABLE COPY

Square Spinner

15"	A4	B1	B3	B4
18"	A5	B3	B4	G1
21"	B1	B4	G1	G3
24"	B3	B5	G3	G5

Round Spinner

36"	C1	C5	D5	F1
-----	----	----	----	----

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED
		1.5 MPH
		2.25 MPH

Spiral Spinner	6"	A1	A2
-----------------------	----	----	----

12"	A3	B1
-----	----	----

Square Spinner

18"	D1	B4
-----	----	----


24"	B3	C2
-----	----	----

Serrated Spinner

30"	B5	C5
-----	----	----


36"	C2	D3
-----	----	----

The recommended rate in this guide represents 2.25 pounds acid equivalent glyphosate or 2.25 gallons of product per acre.



How to Calibrate

1. Determine the proper Multi Flow™ units required from the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for a 1.5 mph walking speed and a 12" swath width, 11 to 22 Multi Flow units are recommended. Use the lowest setting for easy-to-control weeds such as small annuals, and the highest setting for difficult-to-control weeds such as bermudagrass and field bindweed.
2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held above the handle but below the level of the product container.
3. Remove the spray—disc spinner and attach the Expedite calibration flow cup.
4. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
5. To remove the product from the calibration cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.
7. When the desired head setting is obtained, remove the calibration cup and reattach the desired spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.



80432

EXPEDITE® GRASS & WEED II HERBICIDE CALIBRATION GUIDE

30 SECOND MULTI FLOW™ MEASUREMENT*

SPINNER TYPE	SPRAY	WALKING SPEED			
	SWATH WIDTH	1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH
Square Spinner	6"	4 to 7	5 to 10	6 to 12	7 to 14
	9"	5 to 11	7 to 14	9 to 18	11 to 22
	12"	8 to 16	11 to 22	13 to 27	16 to 32
	15"	11 to 22	14 to 28	18 to 36	22 to 43
	18"	13 to 27	18 to 36	22 to 45	27 to 54
	21"	16 to 32	22 to 43	27 to 54	32 to 65
	24"	19 to 38	25 to 50	31 to 63	38 to 75
	36"	22 to 43	29 to 57	36 to 72	43 to 86
Round Spinner	36"	32 to 65	43 to 86	54 to 108	65 to 129

SPINNER TYPE	SPRAY	WALKING SPEED	
	SWATH WIDTH	1.5 MPH	2.25 MPH
Spiral Spinner	6"	5 to 11	8 to 16
	12"	11 to 21	16 to 32
Square Spinner	18"	16 to 32	24 to 48
	24"	21 to 42	32 to 64
Round Spinner	30"	26 to 52	38 to 76
	36"	32 to 64	48 to 96

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.
*One Multi Flow equals 0.8 milliliter.
TMMulti Flow is a trademark of Nonix, Plc.

CROPPING SYSTEMS

For crops listed in this section of this label, this product may be used as directed prior to the emergence of direct seeded crops, prior to transplanting of crops, in established groves, vineyards or orchards and for spot treatments as specified. For crops not listed, planting must be delayed for 30 days following an application.

This product does not provide residual weed control. For subsequent weed control, follow a program using residual herbicides or use repeat applications of this product. When using repeat applications of this product, do not exceed 2 applications per year in cropping systems or 3 applications per year in established groves, vineyards or orchards.

Unless otherwise directed in this label, do not harvest or feed treated crops for 8 weeks after application of this product. Allow 14 days following spot treatments before grazing domestic livestock or harvesting forage grasses and legumes.

When using this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 1/2 inch of rainfall or irrigation.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURE BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

FIELD CROPS

BEST AVAILABLE COPY

This product can be used prior to the emergence of direct seeded crops or prior to transplanting of crops in the following list.

ALFALFA*
ARTICHOKE,
JERUSALEM
ASPARAGUS
ATEMOYA

KALE
LENTILS
LETTUCE
LOGANBERRY
MELONS***

BARLEY*
 BEANS (ALL)
 BEET GREENS
 BEETS (RED, SUGAR)
 BLACKBERRY
 BLUEBERRY
 BOYSENBERRY
 BROCCOLI
 CABBAGE
 CARAMBOLA
 CARROT
 CAULIFLOWER
 CELERY
 CHICORY
 CORN (ALL) *
 COTTON*
 CRANBERRY
 CUCUMBER***
 CURRANT
 DEWBERRY
 ELDERBERRY
 FORAGE GRASSES*
 FORAGE LEGUMES*
 GARLIC***
 GOOSEBERRY
 GOURDS***
 HORSERADISH
 HUCKLEBERRY

MUSTARD GREENS
 OATS*
 OKRA
 OLALLIEBERRY
 ONION
 PARSNIPS
 PEANUTS
 PEAS (ALL)
 PEPPER***
 PINEAPPLE****
 POTATO (IRISH, SWEET)
 PUMPKIN***
 RADISH
 RASPBERRY (BLACK, RED)
 RICE**
 RUTABAGA
 SORGHUM (MILO) *
 SOYBEANS*
 SPINACH
 SQUASH*** (SUMMER, WINTER)
 SUGAR APPLE
 TOMATO***
 TOMATOES****+
 TURNIPS
 WATERMELONS***
 WATERCRESS***
 WHEAT*
 YAMS

*Spot treatments may be applied in these crops and must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn and boll opening on cotton
 **Do not treat rice fields or levees when the fields contain flood water.
 ***Apply only prior to planting. Allow at least 3 days between application and planting.
 ****Do not feed or graze treated pineapple forage following application.
 +Use is restricted to direct seeded crops only.

NOTE: This product is not registered in California for use on atemoya, carambola, sugar apples, yams and parsnips.

TREE CROPS

BEST AVAILABLE COPY

This product is recommended for weed control in established groves, vineyards, or orchards, or for site preparation prior to transplanting these crops as directed below.

CITRUS*: citron, grapefruit, kumquat, lemon, lime, orange, tangelo, tangerine.

NUTS**: almond, filbert, macadamia, pecan, pistachio, walnut.

POME FRUIT*: apple, pear.

STONE FRUIT***: apricots, cherries, nectarines, olives, peaches, plums/prunes. Remove suckers and low hanging limbs at least 10 days prior to application. Avoid application near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. Cherries may be treated with this product in all states. Apricots, nectarines, peaches and plums/prunes may only be treated in the states of Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington. If a spray shield is used and the application is made no later than 90 days after first bloom, this product may be used on peaches in the states of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee. EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED BY THIS PRODUCT.

TROPICAL FRUIT: acerola, atemoya*, avocado*, banana(plantains), carambola*, coffee****, figs*, guava, mango*, papaya, sugar apple*, tea. Allow a minimum of 1 day between last application and harvest of guava and papaya. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

*Allow a minimum of 14 days between last application and harvest.

**Allow a minimum of 21 days between last application and harvest.

***Allow a minimum of 17 days between last application and harvest.

****Allow a minimum of 28 days between last application and harvest.

VINE CROPS

BEST AVAILABLE COPY

KIWI FRUIT*

GRAPES:* Any variety of table, wine, or raisin grape may be treated with this product. In the Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

*Applications should not be made when green shoots, canes, or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest.

10/2/92

Cleaning of Equipment

The Expedite application equipment must be cleaned after each use and prior to storage. Use Expedite cleaner to remove residues of this product. To clean the lance, connect the lance to the cleaner box, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner box and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. Failure to properly clean the equipment may result in equipment failure or injury to subsequently treated desirable vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Expedite application equipment.

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides control of many grasses and broadleaf weeds, including the following:

Bahiagrass
Barnyardgrass
Bentgrass
Bermudagrass*
Bindweed, field*
Bluegrass, annual
Brome, downy
Bromegrass, smooth
Cheat
Chickweed
Clover: red, white
Crabgrass
Dallisgrass
Dandelion
Dock, curly
Foxtail
Geranium, Carolina
Goosegrass
Guineagrass
Lambsquarters, common

Fescues
Johnsongrass
Lettuce, prickly
Mustard, tansy
Nutsedge: purple, yellow
Orchardgrass
Quackgrass
Pennycress, field
Pigweed: redroot, smooth
Ragweed, common
Ryegrass: annual, perennial
Sandbur, field
Shepherdspurse
Sowthistle
Stinkgrass
Thistle, Canada
Timothy
Vaseygrass

*Partial control

BEST AVAILABLE COPY

130732

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description or the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

BEST AVAILABLE COPY

875.05-300.xx/53

140132

EPA Reg. No. 524-450

©MONSANTO COMPANY 1992

In case of an emergency involving this product, Call
Collect,
day or night, (314) 694-4000.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

11/21/90

15 p 732



GRASS & WEED II HERBICIDE

by Monsanto

HIGH VISIBILITY - NO MARK
FOR NON-SELECTIVE POSTEMERGENT CONTROL OF UNWANTED VEGETATION
Specially formulated for Expedite® Application Equipment

Keep out of reach of children.
CAUTION!

Covers from 14,000 to 25,000 square feet

Read the entire label before using this product.
Use only according to label instructions.
Read "LIMIT OF WARRANTY AND LIABILITY" before buying
or using. If terms are not acceptable, return at once
unopened.
REFORMULATION IS PROHIBITED.

ACTIVE INGREDIENT:
*Glyphosate, N-(phosphonomethyl)glycine
in the form of its isopropylamine
salt _____ 15.8%
INERT INGREDIENTS: _____ 84.2%
100.0%

This package may be used to refill 750 ml containers that are
specifically labeled for refilling with this product. Follow refilling
instructions on side panel of this container.

Contains petroleum distillates.
*Contains 162 grams per liter or 1.35 pounds per U.S. gallon
of the active ingredient, glyphosate, in the form of its
isopropylamine salt. Equivalent to 120 grams per liter or 1.0
pound per U.S. gallon of the acid, glyphosate.

ACCEPTED
APR - 1 1993
EPA Reg. No. 524-450


PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION!
CAUSES EYE IRRITATION.
HARMFUL IF INHALED.
MAY CAUSE RESPIRATORY TRACT IRRITATION.
Avoid contact with eyes, skin or clothing.
Avoid breathing vapor or spray mist.
Wash thoroughly with soap and water after handling.
FIRST AID: IF IN EYES, immediately flush with plenty of
water. Get medical attention.
IF INHALED, remove individual to
fresh air. Seek medical attention
if breathing difficulty develops.

This product is protected by U.S. Patent No. 4,405,531. Other
patents pending. No license granted under any non-U.S.
patents.
TM Expedite is a trademark of, and used under license from,
Monsanto.

This product contains glyphosate, the active
ingredient in Roundup® herbicide.
* Roundup is a registered
trademark of Monsanto Company.
© MONSANTO COMPANY 1990
EPA Reg. No. 624-480

(Continued on back panel)
In case of a human health emergency involving this product,
Call Collect, day or night, (314) 634-4000.



NET 1 GALLON
Part 1 of 2
875.15-308.02/534P

BEST AVAILABLE COPY

160732

EXPEDITE **GRASS & WEED II HERBICIDE** by Monsanto

(Continued from front panel)

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32° F and 100° F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emptied, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(See label booklet for Complete Direction for Use.)

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based on contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OF IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.


MONSANTO COMPANY, AGRICULTURAL PRODUCTS,
ST. LOUIS, MISSOURI 63167 U.S.A.

EPA Est. 34313-TX-001


Part 2 of 2
878.18-300.03/03/NT

BEST AVAILABLE COPY


AS A BACKPACK




Insert Nomix Link-Up cassette into the Compact



Insert Big Five into backpack. Connect to herbicide pack by aligning groove of pipe to ridges of Big Five and twist through 90° clockwise.




Raise Compact above backpack. Depress Compact trigger to drain herbicide through lance and into Big Five.




Unlock by twisting anti-clockwise through 90° and disconnect.

TO REFILL


170432




Align grooves of the Compact cartridge to ridges of the Big Five refill pack.



Twist the cartridge clockwise through 90° to lock the two containers and open valves.



Invert, ensuring the Big Five pack is uppermost and allow the cartridge to refill.



Unlock the packs by twisting anti-clockwise through 90° and disconnect.

Keep a cloth handy to wipe the nozzle of the refilled cartridge.

10. Empty cartridges should only be refilled for one session. Ensure cartridges are only refilled with the same herbicide as originally located.



EXPEDITE

GRASS & WEED II HERBICIDE

by Mow-into

HIGH VISIBILITY - NO MARK
FOR NON-SELECTIVE POSTEMERGENT CONTROL OF UNWANTED VEGETATION
Specially Formulated for Broadcast Application Equipment

Keep out of reach of children.

Covers from 2,900 to 5,700 square feet.

CAUTION!

Read the entire label before using this product.
Use only according to label instructions.
Read "LIMIT OF WARRANTY AND LIABILITY" before buying
or using. If terms are not acceptable, return of your
purchase.

REFORMULATION IS PROHIBITED.

This container may be filled with this product from 1 gallon
containers specifically labeled for refilling. Follow refilling
instructions on the side panel of the 1 gallon container.

PRECAUTIONARY STATEMENTS

Hazard to Fishes and Aquatic Animals

CAUTION

CAUSES EYE IRRITATION.
IRRITATE, IF SWALLOWED,
MAY CAUSE RESPIRATORY TRACT IRRITATION.
Avoid contact with eyes, nose or clothing.
Avoid breathing vapor of spray mist.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, Immediately flush with plenty of

water. Get medical attention.

IF SWALLOWED, Remove individual or

if breathing difficulty develops.

Contains an acid stain

In case of a human health emergency involving this product,
Call Calson, day or night, (714) 884-4500.

ACTIVE INGREDIENT:

*Glyphosate, is-glyphosate-monomethylammonium
in the form of its isopropylamide

100%
100.00%
100.00%

Contains petroleum distillates.

*Contains 100 grams per liter or 1.25 pounds per U.S. gallon
of the active ingredient, glyphosate, in the form of its
isopropylamide salt. Equivalent to 120 grams per liter or 1.5
pounds per U.S. gallon of the salt, glyphosate.

This product is registered by U.S. Patent No. 4,498,521. Other
patents pending. No claims granted under any non-U.S.
patents.

TM Expires in a maximum of 600 hours unless otherwise stated.
Roundup

This product contains glyphosate, the active
ingredient in Roundup® herbicide.

*Roundup is a registered
trademark of Monsanto Company.

A MONSANTO COMPANY 1990
EPA Reg. No. 536-020

Roundup

NET 750 ML

Part 1 of 2
079.15-000.00/0207

ACCEPTED
APR - 1 1993
524-450

BEST AVAILABLE COPY

190732

EXPEDITE[®] GRASS & WEED^{II} HERBICIDE by Monsanto

(Continued from last page)

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to irrigation areas below the mean high water mark. Do not discharge water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite applicator equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED IRON CONTAINERS OR SPRAY TANKS. This product will react with such containers and tends to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, light & cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32° F and 100° F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a method approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

When container is completely emptied through Expedite application equipment, or can no longer be refilled, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(See label booklet for Complete Direction for Use)

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with these Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, unusual conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application range set forth in the Directions, application in any manner not exactly set forth in the Directions, moisture conditions not in the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT PURCHASE PRICE OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

MONSANTO COMPANY, AGRICULTURAL PRODUCTS,
ST. LOUIS, MISSOURI 63167 U.S.A.

EPA Est. 34515-TX-601

Page 2 of 2
876 16-388-28E247

BEST AVAILABLE COPY

500432

**EXPEDITE®
GRASS & WEED II HERBICIDE
by Monsanto**

High Visibility - No Mark

**For Non-Selective Postemergence Control of Unwanted Vegetation
Contains glyphosate, the active ingredient in Roundup® herbicide.**

[The Roundup Logo]

Specially formulated for Expedite™ Application Equipment.

Complete Directions for Use

EPA Reg. No. 524-450

Covers from 2,900 to 5,700 square feet.

®™ Expedite is a registered trademark of, and used under license from, Nomix Plc.

® Roundup is a registered trademark of Monsanto Company.

1993-1

XXX.XX.XXX.XX/XX

ACCEPTED
APR - 1 1993
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 524-450

©MONSANTO COMPANY 1993

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(COMPACT.LBL)

(11/9/90)

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED. [REDACTED]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

HARMFUL IF INHALED.

MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid contact with eyes, skin or clothing.

Avoid breathing vapor or spray mist.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of an **ana human health** emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to interstitial areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.



204-2

Physical or Chemical Hazards

Apply this product ONLY through sprayer application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonorethyl) glycine
in the form of its isopropylamine salt15.9 %

INERT INGREDIENTS:.....84.1 %
100.0 %

Contains petroleum distillates.

*Contains 162 grams per litre or 1.35 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 120 grams per litre or 1.0 pound per U.S. gallon of the acid, glyphosate).

This product is protected by U.S. Patent No. 4,405,531. Other patents are pending. No license granted under any non-U.S. patent(s).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

BEST AVAILABLE COPY

22/12/90

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. ~~Do not reuse container.~~

When container is completely emptied through Expedite application equipment, ~~it can no longer be refilled~~ puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a nonselective weed killer that is applied, using Expedite application equipment, to the foliage of actively growing weeds. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms of control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

This product is recommended for commercial, forestry, industrial, noncropland and residential vegetation management in areas such as airports, ditchbanks, farmsteads, fencerows, forestry and Christmas tree plantations, greenhouses, golf courses, highways and roadsides, industrial plant sites, institutional sites, municipalities, lawns, lumberyards, nurseries, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites. This product is also recommended for weed control in agricultural crops as specified in the "CROPPING SYSTEMS" section of this label.

When this product is used as a directed application to control undesirable vegetation growing around established ornamentals, trees, shrubs or vines, avoid spray or drift contact with foliage, stems or green (immature) tree bark.

Rainfall or irrigation within 2 hours after application may wash the product off plant foliage resulting in reduced weed control. Delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application to allow product to move within the plant for the most effective control.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY CONTACT WITH THE FOLIAGE OF DESIRABLE TURF, ORNAMENTALS, SHRUBS OR TREES. DO NOT APPLY OVER THE TOP OF DESIRABLE PLANTS.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

BEST AVAILABLE COPY

240732

How to Apply

Apply this product as a uniform spray over the foliage of undesirable vegetation. Avoid applications during windy conditions, as spray drift or swath displacement may cause damage to any vegetation contacted. When used in greenhouses, turn off air circulation fans while spraying. Use only Expedite application equipment when applying this product.

When used in greenhouses, turn off air circulation fans while spraying. **DESIRABLE GREENHOUSE PLANTS WILL BE INJURED IF CONTACTED BY SPRAY OR DRIFT OF THIS PRODUCT.**


To select general flow rate settings for applying this product, use the Expedite Set-N-Spray settings listed in the following application guide. The letters followed by a number (A2...E3G5) represent equipment settings for flow rates within the application range recommended for this product. If calibration is desired to obtain equipment settings, see the "HOW TO CALIBRATE" section of this label. Calibration is recommended where more precise equipment calibration is desired, or when the application rate needs to be customized for the type of weeds present.

When the desired head setting is obtained, install the spinner as specified in the following guide. For best results, make applications before weeds are allowed to form a dense canopy. Best results are obtained when the spinner is producing a uniform spray pattern. For spray widths less than 15 inches, this is obtained by holding the lance head as close as practical to the top of the weed canopy (less than 4 inches) and increasing the spinner speed to obtain the correct swath width.

EXPEDITE® GRASS & WEED II HERBICIDE APPLICATION GUIDE

SET-N-SPRAY SETTING

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED	
		1.5 MPH	2.25 MPH
	<6"	A2	A3
Spiral Spinner	6"	A3	A5
	9"	A5	B2



	12"	B1	B4
Square Spinner	15"	B3	C1
	18"	B4	C3
	21"	B5	C4
Serrated Spinner	24"	C1	D1
	36"	D1	E3

The recommended rate in this guide represents 2.25 pounds acid equivalent glyphosate or 2.25 gallons of product per acre.

How to Calibrate

1. Determine the proper Multi Flow™ units required from the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for a 1.5 mph walking speed and a 12" swath width, 11 to 22 Multi Flow units are recommended. Use the lowest setting for easy-to-control weeds such as small annuals, and the highest setting for difficult-to-control weeds such as bermudagrass and field bindweed.
2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held below the level of the product container.
3. Remove the spray disc and attach the Expedite calibration flow cup.
4. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
5. To remove the product from the calibration cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.



260032

7. When the desired head setting is obtained, remove the calibration cup and reattach the desired spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.

**EXPEDITE® GRASS & WEED II HERBICIDE
CALIBRATION GUIDE**

30 SECOND MULTI FLOW™ MEASUREMENT*

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED	
		1.5 MPH	2.25 MPH
Spiral Spinner	<6"	4 to 7	5 to 11
	6"	5 to 11	8 to 16
	9"	8 to 16	12 to 24
	12"	11 to 22	16 to 32
	15"	13 to 27	20 to 41
	18"	16 to 32	24 to 49
Square Spinner	21"	19 to 38	29 to 56
	24"	22 to 43	32 to 65
Serrated Spinner	36"	32 to 65	49 to 97

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.
*One Multi Flow equals 0.8 milliliter.

™Multi Flow is a trademark of Nomix, Plc.

BEST AVAILABLE COPY

CROPPING SYSTEMS

For crops listed in this section of this label, this product may be used as directed prior to the emergence of direct seeded crops, prior to transplanting of crops, in established groves, vineyards or orchards and for spot treatments as specified. For crops not listed, planting must be delayed for 30 days following an application.

This product does not provide residual weed control. For subsequent weed control, follow a program using residual herbicides or us repeat applications of this product. When using repeat applications of this product, do not exceed 2 applications per year in cropping systems or 3 applications per year in established groves, vineyards or orchards.

Unless otherwise directed in this label, do not harvest or feed treated crops for 8 weeks after application of this product. Allow 14 days following spot treatments before grazing domestic livestock or harvesting forage grasses and legumes.

When using this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 1/2 inch of rainfall or irrigation.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURE BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

FIELD CROPS

This product can be used prior to the emergence of direct seeded crops or prior to transplanting of crops in the following list.

- | | |
|--------------------|----------------|
| ALFALFA* | KALE |
| ARTICHOKE, | LENTILS |
| JERUSALEM | LETTUCE |
| ASPARAGUS | LOGANBERRY |
| ATEMOYA | MELCNS*** |
| BARLEY* | MUSTARD GREENS |
| BEANS (ALL) | O.TS* |
| BEET GREENS | OKRA |
| BEETS (RED, SUGAR) | OLALLIEBERRY |
| BLACKBERRY | ONION |
| BLUEBERRY | PARSNIPS |
| BOYSENBERRY | PEANUTS |
| BROCCOLI | PEAS (ALL) |
| CABBAGE | PEPPER*** |

CARAMBOLA
 CARROT
 CAULIFLOWER
 CELERY
 CHICORY
 CORN (ALL) *
 COTTON*
 CRANBERRY
 CUCUMBER***
 CURRANT
 DEWBERRY
 ELDERBERRY
 FORAGE GRASSES*
 FORAGE LEGUMES*
 GARLIC***
 GOOSEBERRY
 GOURDS***
 HORSERADISH
 HUCKLEBERRY

PINEAPPLE****
 POTATO (IRISH, SWEET)
 PUMPKIN***
 RADISH
 RASPBERRY (BLACK, RED)
 RICE**
 RUTABAGA
 SORGHUM (MILO) *
 SOYBEANS*
 SPINACH
 SQUASH*** (SUMMER, WINTER)
 SUGAR APPLE
 TOMATILLO***
 TOMATOES****+
 TURNIPS
 WATERMELONS***
 WATERCRESS***
 WHEAT*
 YAMS

*Spot treatments may be applied in these crops and must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn and boll opening on cotton

**Do not treat rice fields or levees when the fields contain flood water.

***Apply only prior to planting. Allow at least 3 days between application and planting.

****Do not feed or graze treated pineapple forage following application.

+Use is restricted to direct seeded crops only.

NOTE: This product is not registered in California for use on atemoya, carambola, sugar apples, yams and parsnips.

TREE CROPS

This product is recommended for weed control in established groves, vineyards, or orchards, or for site preparation prior to transplanting these crops as directed below.

CITRUS*: citron, grapefruit, kumquat, lemon, lime, orange, tangelo, tangerine.

NUTS**: almond, filbert, macadamia, pecan, pistachio, walnut.

POME FRUIT*: apple, pear.

STONE FRUIT*:** apric , cherries, nectarines, olives, peaches, plums/prunes. Remove suckers and low hanging limbs at least 10 days prior to application. Avoid application near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. Cherries may be treated with this product in all states. Apricots, nectarines, peaches and plums/prunes may only be treated in the states of Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington. If a spray shield is used and the application is made no later than 90 days after first bloom, this product may be used on peaches in the states of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED BY THIS PRODUCT.**

TROPICAL FRUIT: acerola, atemoya*, avocado*, banana(plantains), carambola*, coffee:****, figs*, guava, mango*, papaya, sugar apple*, tea. Allow a minimum of 1 day between last application and harvest of guava and papaya. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

- *Allow a minimum of 14 days between last application and harvest.
- **Allow a minimum of 21 days between last application and harvest.
- ***Allow a minimum of 17 days between last application and harvest.
- ****Allow a minimum of 28 days between last application and harvest.

VINE CROPS

KIWI FRUIT*

GRAPES:* Any variety of table, wine, or raisin grape may be treated with this product. In the Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

*Applications should not be made when green shoots, canes, or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest.

BEST AVAILABLE COPY

Cleaning of Equipment

The Expedite application equipment must be cleaned after each use and prior to storage. Use Expedite cleaner to remove residues of this product. To clean the lance, connect the lance to the cleaner box, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner box and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. Failure to properly clean the equipment may result in equipment failure or injury to subsequently treated desirable vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Expedite application equipment.

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides control of many grasses and broadleaf weeds, including the following:

- Bahiagrass
- Barnyardgrass
- Bentgrass
- Bermudagrass*
- Bindweed, field*
- Bluegrass, annual
- Brome, do
- Bromegrass, smooth
- Cheat
- Chickweed
- Clover: red, white
- Crabgrass
- Dallisgrass
- Dandelion
- Dock, curly
- Fescues
- Foxtail
- Geranium, Carolina
- Goosegrass
- Guineagrass

- Johnsongrass
- Lambsquarters, common
- Lettuce, prickly
- Mustard, t nsy
- Nutsedge:
- Nutsedge: purple, yellow
- Orchardgrass
- Pennycress, field
- Pigweed: redroot, smooth
- Quackgrass
- Ragweed, common
- Ryegrass: annual, perennial
- Sandbur, field
- Shepherdspurse
- Sowthistle
- Stinkgrass
- Thistle, Canada
- Timothy
- Vaseygrass

*Partial control

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

xxx.xx-xxx.xx/xx

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

220732

©MONSANTO COMPANY 1993

In case of a human health emergency involving this product, Call
Collect, |
|day or night, (314) 694-4000. |

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

5 5