

PM-25

Reg # 524-450

19 20

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 1-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
235677

Application for Pesticide - Section I

1. Company/Product Number Monsanto / 524-450	2. EPA Product Manager Robert J. Taylor	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Expedite Grass & Weed II Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Monsanto Company 700 14th Street, N.W., #1100 Washington, D.C. 20005 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Submit revised labels to change chart and calibration settings for Expedite & Nomix lances.

NOTIFICATION

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container 1.32 gal; 1 gal; 750 ml			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 1.32 gal; 1 gal; 750 ml		5. Location of Label Directions <input type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Russell P. Schneider, Ph.D.	Title Agricultural Regulation Director	Telephone No. (Include Area Code) (202) 783-2460
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager	
4. Typed Name Lydia A. Suba	5. Date Nov 8/96	

29 20
8


EXPEDITE® GRASS & WEED II HERBICIDE
EPA Reg. No. 524-450

LABEL for 1.32 GALLON SIZE PACK

Revised from stamped label March 24, 1994

NOTIFICATION

MONSANTO COMPANY
St. Louis, Missouri 63167
November, 1996



3920

**EXPEDITE®
GRASS & WEED II HERBICIDE
by Monsanto**

High Visibility - No Mark

For Non-Selective Postemergence Control of Unwanted Vegetation

Contains glyphosate, the active ingredient in Roundup® herbicide.

[The Roundup Logo]

Specially formulated for Expedite® and Nomix™ Application Equipment.

Complete Directions for Use

EPA Reg. No. 524-450

Covers from 19,000 to 38,000 square feet.


*Roundup is a registered trademark of Monsanto Company.

1993-1

875.15.300.xx/53

©MONSANTO COMPANY 1993

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



4920

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

HARMFUL IF INHALED.

MAY CAUSE RESPIRATORY TRACT IRRITATION.

Avoid contact with eyes, skin or clothing.


Avoid breathing vapor or spray mist.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



User Safety Recommendation:
 User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of an emergency involving this product,
 Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment cleaning solutions.

Physical or Chemical Hazards

Apply this product ONLY through Expedite and Nomix application equipment designed for its use. DO NOT STORE OR APPLY THIS PRODUCT IN ANY OTHER TYPE OF CONTAINER, PARTICULARLY GALVANIZED STEEL OR UNLINED STEEL CONTAINERS OR SPRAY TANKS. This product will react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

* Glyphosate, N-(phosphonomethyl) glycine	
in the form of its isopropylamine salt	15.8 %

INERT INGREDIENTS:	84.2 %
	100.0 %

Contains petroleum distillates.

6920

- * Contains 162 grams per litre or 1.35 pounds per U.S. gallon of the active ingredient, glyphosate, in the form of its isopropylamine salt. Equivalent to 120 grams per litre or 1.0 pound per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531.
Other patents are pending.

No license granted under any non-U.S. patent(s).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

- coveralls
- waterproof gloves
- shoes plus socks

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product between 32°F and 100°F in its original container and in a secured storage area.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container.

When container is completely emptied, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a nonselective weed killer that is applied, using Expedite and Nomix application equipment, to the foliage of actively growing weeds. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms of control may require a week or more to appear, with complete browning usually occurring in 1 to 4 weeks. Symptoms are a gradual wilting and yellowing of the treated plant foliage followed by browning and deterioration of both shoots and roots.

THIS PRODUCT IS FOR NON-RESIDENTIAL USE ONLY such as commercial, forestry, industrial, noncropland vegetation management in areas such as airports, ditchbanks, farmsteads, fencerows, forestry and Christmas tree plantations, greenhouses, golf courses, highways and roadsides, industrial plant sites, institutional sites, municipalities, lawns, lumberyards, nurseries, parking areas, parks, playgrounds, petroleum tank farms, pipeline, power, telephone and utility rights-of-way, railroads, schools, storage areas and similar sites. This product is also recommended for weed control in agricultural crops as specified in the "CROPPING SYSTEMS" section of this label.

When this product is used as a directed application to control undesirable vegetation growing around established ornamentals, trees, shrubs or vines, avoid spray or drift contact with foliage, stems or green (immature) tree bark.

8920

INSTRUCTIONS FOR SETTING UP AND CALIBRATION OF
COMPACT MARK I AND EXPEDITE ~~SUPERPRO~~

EXPEDITE® GRASS & WEED II HERBICIDE
EXPEDITE ~~SUPERPRO~~ APPLICATION GUIDE

SET-N-SPRAY SETTING

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED			
		1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH
Spiral Spinner	6"	A1	A1 to A2	A1 to A4	A1 to A5
	12"	A1 to A5	A2 to B2	A4 to B3	A5 to B5
Square Spinner	18"	A3 to B3	A5 to B5	B1 to C2	B3 to C4
	24"	A5 to B5	B2 to C3	B3 to C5	B5 to D4
Serrated Spinner	30"	B1 to C2	B3 to C5	B5 to D5	C2 to F2
	36"	B3 to C4	B5 to D4	C2 to F2	C4 to G5

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.

9920


EXPEDITE® GRASS & WEED II HERBICIDE
COMPACT MARK I APPLICATION GUIDE

SET-N-SPRAY SETTINGS

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED 2.25 MPH
Spiral Spinner	6"	A1 to A4
	12"	A4 to B3
Square Spinner	18"	B2 to C3
	24"	B3 to D2
Serrated Spinner	30"	B5 to E1
	36"	C3 to F5

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.

CALIBRATION INSTRUCTIONS FOR THE EXPEDITE AND COMPACT MARK I



107 20

How to Calibrate

1. Determine the proper Multi Flow[®] units required from the following calibration guide. The number of Multi Flow units required is where the swath width row intersects the walking speed column. For example, for the Expedite at a 1.5 mph walking speed and a 12" swath width, 11 to 22 Multi Flow units are recommended. Use the lowest setting for easy-to-control weeds such as small annuals, and the highest setting for difficult-to-control weeds such as bermudagrass and field bindweed.
2. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the head setting to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held above the handle but below the level of the product container.
3. Remove the spinner and attach the Calibration Flow Cup.
4. Lower the lance head to the spraying position, depress the trigger for 30 seconds and compare the measured volume to the desired Multi Flow units.
5. To remove the product from the Calibration Flow Cup, raise the lance above head level and depress the trigger long enough to allow the product to flow back into the lance. Do not allow air to reenter the lance. If air enters the lance, it must be reprimed.
6. If required, adjust the head setting and repeat the process until the desired Multi Flow units are obtained.
7. When the desired head setting is obtained, remove the Calibration Flow Cup and reattach the desired spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.

119 20

EXPEDITE® GRASS & WEED II HERBICIDE
EXPEDITE/SUPERPRO CALIBRATION GUIDE

30 SECOND MULTI FLOW™ MEASUREMENT*

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED			
		1.5 MPH	2.0 MPH	2.5 MPH	3.0 MPH
Spiral Spinner	6"	5 to 11	7 to 14	9 to 18	11 to 22
	12"	11 to 22	14 to 28	18 to 36	22 to 43
Square Spinner	18"	16 to 32	22 to 43	27 to 54	32 to 65
	24"	22 to 43	29 to 57	36 to 72	43 to 86
Serrated Spinner	30"	27 to 54	36 to 72	45 to 90	54 to 108
	36"	32 to 65	43 to 86	54 to 108	65 to 129

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.

*One Multi Flow equals 0.8 milliliter.

12720

EXPEDITE® GRASS & WEED II HERBICIDE
COMPACT MARK I CALIBRATION GUIDE

30 SECOND MULTI FLOW® MEASUREMENT*

SPINNER TYPE	SPRAY SWATH WIDTH	WALKING SPEED 2.25 MPH
--------------	-------------------	---------------------------

Spiral Spinner	6"	8 to 16
----------------	----	---------

	12"	16 to 32
--	-----	----------

Square Spinner	18"	24 to 48
----------------	-----	----------

	24"	32 to 63
--	-----	----------

Serrated Spinner	30"	40 to 79
------------------	-----	----------

	36"	48 to 95
--	-----	----------

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.

*One Multi Flow equals 0.8 milliliter.

13720

INSTRUCTIONS FOR THE SETTING UP AND CALIBRATION
OF THE ~~NOMEX 700 AND~~ COMPACT MARK II

EXPEDITE® GRASS & WEED II HERBICIDE
~~NOMEX 700 AND~~ COMPACT MARK II
SET-N-SPRAY SETTINGS AND CALIBRATION GUIDE

WALKING SPEED ~~1.5 YARDS/SEC~~ (2.25 MPH)

SPINNER TYPE	SPRAY SWATH WIDTH	CALIBRATION TIME
		21 (secs) 11

Spiral Spinner	6"	A2 to B3 A3 to C2
----------------	----	------------------------------

	12"	B3 to D3 C1 to D4
--	-----	------------------------------

Square Spinner	18"	C3 to E4 C5 to F4
----------------	-----	------------------------------

	24"	D3 to E4 D4 to G1
--	-----	------------------------------

Serrated Spinner	30"	D5 to G4 E2 to G4
------------------	-----	------------------------------

	36"	E4 to F2 to *
--	-----	--------------------------

The rate range in this guide represents 1.5 to 3.0 pounds acid equivalent glyphosate or 1.5 to 3.0 gallons of product per acre.

* The flow rate through the equipment may not be adequate to apply the recommended rate at these walking speed and swath width combinations.

14 g 20

CALIBRATION INSTRUCTIONS FOR ~~NOMEX 700~~ AND COMPACT MARK II

HOW TO CALIBRATE

1. Determine the proper flow rate setting required from the above Calibration guide. The flow rate setting required is where the swath width row intersects the Set-N-Spray column. For example, for a 12" swath width with the ~~Nomex 700~~ Compact Mark II, settings between ~~B3 to D3~~ B5 to D2 are recommended. Use the lowest setting ~~(B3)~~ (B5) for easy-to-control weeds such as small annuals, and the highest setting ~~(D3)~~ (D2) for difficult-to-control weeds such as bermudagrass and field bindweed.
2. Note the time required for calibration as shown above the required Flow Rate setting column. For example, for applying at the maximum rate, a calibration time of 11 seconds is required.
3. Before calibrating, the lance must be primed to remove trapped air. To prime the lance, set the Flow Rate control to G+ and depress the trigger until all air is removed from the lance. Priming must be done with the head of the lance held lower than the product container.
4. Remove the spinner and attach the Calibration Flow Cup.
5. Prime the Calibration Flow Cup by lowering the lance into the spraying position, depressing the trigger for 5 seconds. Raise the spray head to head level and depress the trigger to allow the product to flow back to the zero line of the Calibration Flow Cup.
6. Lower the lance head to the spraying position, depress the trigger for the required calibration time and check the measured volume matches the desired spray swath width line on the Calibration Flow Cup.
7. If the measured volume does not match the line, adjust the Flow Rate control up or down and repeat the process until the desired volume is obtained.
8. To remove the product from the Calibration Flow Cup, raise the spray head to head height and depress the trigger to allow the Calibration Flow Cup to empty. Do not allow air to reenter the lance. If air enters the lance, ie. the Flow Rate window empties, the lance must be reprimed.
9. When the desired flow rate is obtained, remove the Calibration Flow Cup and reattach the desired spinner and follow the instructions in the "HOW TO APPLY" section on how to spray this product.

15720

CROPPING SYSTEMS

For crops listed in this section of this label, this product may be used as directed prior to the emergence of direct seeded crops, prior to transplanting of crops, in established groves, vineyards or orchards and for spot treatments as specified. For crops not listed, planting must be delayed for 30 days following an application.

This product does not provide residual weed control. For subsequent weed control, follow a program using residual herbicides or us repeat applications of this product. When using repeat applications of this product, do not exceed 2 applications per year in cropping systems or 3 applications per year in established groves, vineyards or orchards.

Unless otherwise directed in this label, do not harvest or feed treated crops for 8 weeks after application of this product. Allow 14 days following spot treatments before grazing domestic livestock or harvesting forage grasses and legumes.

When using this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 1/2 inch of rainfall or irrigation.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURE BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

FIELD CROPS

This product can be used prior to the emergence of direct seeded crops or prior to transplanting of crops in the following list.

ALFALFA*	KALE
ARTICHOKE	LENTILS
JERUSALEM	LETTUCE
ASPARAGUS	LOGANBERRY
ATEMOYA	MELONS***
BARLEY*	MUSTARD GREENS
BEANS(ALL)	OATS*
BEET GREENS	OKRA
BEETS(RED, SUGAR)	OLALLIEBERRY
BLACKBERRY	ONION
BLUEBERRY	PARSNIPS
BOYSENBERRY	PEANUTS
BROCCOLI	PEAS(ALL)
CABBAGE	PEPPER***
CARAMBOLA	PINEAPPLE****
CARROT	POTATO(IRISH, SWEET)
CAULIFLOWER	PUMPKIN***
CELERY	RADISH
CHICORY	RASPBERRY(BLACK, RED)
CORN(ALL)*	RICE**
COTTON*	RUTABAGA
CRANBERRY	SORGHUM(MILO)*
CUCUMBER***	SOYBEANS*
CURRANT	SPINACH
DEWBERRY	SQUASH***(SUMMER, WINTER)
ELDERBERRY	SUGAR APPLE
FORAGE GRASSES*	TOMATILO***
FORAGE LEGUMES*	TOMATOES***+
GARLIC***	TURNIPS
GOOSEBERRY	WATERMELONS***
GOURDS***	WATERCRESS***
HORSERADISH	WHEAT*
HUCKLEBERRY	YAMS

* Spot treatments may be applied in these crops and must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn and boll opening on cotton.

** Do not treat rice fields or levees when the fields contain flood water.

TROPICAL FRUIT:

acerola, atemoya*, avocado*, banana plantains), carambola*, coffee****, figs*, guava, mango*, papaya, sugar apple*, tea. Allow a minimum of 1 day between last application and harvest of guava and papaya. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

- * Allow a minimum of 14 days between last application and harvest.
- ** Allow a minimum of 21 days between last application and harvest.
- *** Allow a minimum of 17 days between last application and harvest.
- **** Allow a minimum of 28 days between last application and harvest.

VINE CROPS

KIWI FRUIT*

GRAPES:

- * Any variety of table, wine, or raisin grape may be treated with this product. In the Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.
- * Applications should not be made when green shoots, canes, or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest.

Cleaning of Equipment

The Expedite and Nomix application equipment must be cleaned after each use and prior to storage. ~~Using~~ Use Expedite or Nomix cleaner to remove residues of this product. To clean the lance, connect the lance to the cleaner box, set the head setting to G+ and let the cleaner flow until it runs clear. Disconnect the lance from the cleaner box and remove the remaining cleaner from the lance. Do not drain cleaner back into the box. Failure to properly clean the equipment may result in equipment failure or injury to subsequently treated, desirable, vegetation. After thorough cleaning, equipment may be used to apply other products, especially formulated for use in Expedite and Nomix application equipment.

19720

Wastes resulting from the use or cleaning of equipment that cannot be used or chemically reprocessed should be disposed of in accordance with applicable Federal, state or local procedures. See the "Storage and Disposal" section of this label for more information.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product provides control of many grasses and broadleaf weeds, including the following:

- | | |
|-----------------------|-----------------------------|
| Bahiagrass | Fescues |
| Barnyardgrass | Johnsongrass |
| Bentgrass | Lettuce, prickly |
| Bermudagrass* | Mustard, tansy |
| Bindweed, field* | Nutsedge: purple, yellow |
| Bluegrass, annual | Orchardgrass |
| Brome, downy | Quackgrass |
| Bromegrass, smooth | Pennycress, field |
| Cheat | Pigweed: redroot, smooth |
| Chickweed | Ragweed, common |
| Clover: red, white | Ryegrass: annual, perennial |
| Crabgrass | Sandbur, field |
| Dallisgrass | Shepherdspurse |
| Dandelion | Sowthistle |
| Dock, curly | Stinkgrass |
| Foxtail | Thistle, Canada |
| Geranium, Carolina | Timothy |
| Goosegrass | Vaseygrass |
| Guineagrass | |
| Lambsquarters, common | |

*Partial control

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

20820

Buyers and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

® Expedite and Multi Flow are registered trademarks of Nomix Chipman Limited, UK.

~~NOMIX and SUPERPRO are trademarks of Nomix Chipman Limited, UK~~

875.15-300.xx/53

EPA Reg. No. 524-450

©MONSANTO COMPANY 1993

In case of a human health emergency involving this product,
Call Collect, day or night, (314) 694-4000.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.