

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR MANAGE® TURF HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Manage Turf herbicide and this supplement.

MANAGE®

TURF HERBICIDE BY MONSANTO

EPA Reg. No. 524-465

MANAGE TURF HERBICIDE FOR USE AROUND ESTABLISHED WOODY ORNAMENTALS, FALLOW LAND AND FOR HORSETAIL CONTROL.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Manage® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

ESTABLISHED WOODY ORNAMENTALS

This product may be applied at recommended rates as a post directed spray around any established woody ornamental species. See the label booklet for application instructions, rate recommendations, weeds controlled and proper weed growth stage.

For transplanted woody ornamentals, allow 3 months after transplanting before applying this product.

NOTE: Applications of this product should not contract leaves of desirable plants since foliar injury, discoloration or death may result.

ACCEPTED
MAY 16 1996
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-465

FALLOW TREATMENTS IN TURFGRASS AND ORNAMENTAL PRODUCTION AREAS

This product may be used on fallow land prior to establishing turfgrass or ornamental plants. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage. Allow 4 weeks between application and seeding or transplanting.

ROADSIDES

This product may be applied at recommended rates as a post-emergence spray on roadsides. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage.

CONTROL OF HORSETAIL (*Equisetum sp.*)

For post-emergence control of horsetail (*Equisetum sp.*), apply 1 1/3 ounces of this product per acre (0.062 pound active ingredient per acre) after horsetail has leafed out.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Manage before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1996

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

