

PP 25 524-465 11/3/98 page 18 3

SUPPLEMENTAL LABELING


READ AND FOLLOW THE ENTIRE LABEL FOR MANAGE® TURF HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Manage Turf herbicide and this supplement.

MANAGE®

TURFTURF HERBICIDE BY MONSANTO

EPA Reg. No. 524-465



MANAGE TURFTURF HERBICIDE FOR USE ON ROADSIDES AND OTHER INDUSTRIAL SITES AROUND ESTABLISHED WOODY ORNAMENTALS FALLOW LAND AND FOR HORSETAIL CONTROL.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Manage® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

~~ESTABLISHED WOODY ORNAMENTALS~~

~~This product may be applied at recommended rates as a post directed spray around any established woody ornamental species. See the label booklet for application instructions, rate recommendations, weed controlled and proper weed growth stage.~~

Manage supplemental label revised 8/28/98 NEJ

2 9 3

~~For transplanted woody ornamentals, allow 3 months after transplanting before applying this product.~~

NOTE: Applications of this product should not contact leaves of desirable plants since foliar injury, discoloration or death may result.

FALLOW TREATMENTS IN TURFGRASS AND ORNAMENTAL PRODUCTION AREAS

~~This product may be used on fallow land prior to establishing turfgrass or ornamental plants. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage. Allow 4 weeks between application and seeding or transplanting.~~

ROADSIDES AND OTHER INDUSTRIAL SITES

This product may be applied at recommended-recommended rates as a post-emergence spray on roadsides and other industrial sites. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage.

When applied postemergence, this product will suppress the following weeds:

Cocklebur, common sunflower, common ragweed, giant ragweed, pigweed, and velvetleaf.

This product may be tank mixed with Roundup Pro™ herbicide.

CONTROL OF HORSETAIL (*Equisetum arvense* sp.)

For post-emergence control of horsetail (*Equisetum arvense* sp.), apply 2 2/3 + 1/3 ounces of this product per acre, or 1.8 grams of this product per 1000 sq.ft. (0.125 0.062 pound active ingredient per acre) after horsetail has leafed out. This rate of product will control horsetail that is less than 6" tall and suppress horsetail that is greater than 6" tall.

Herbicide symptoms are likely to show within 2 weeks as a necrotic ring at the base of the plant, even though the leaves and stems remain green and a deep leathery green in color.

Roundup Pro is a trademark of Monsanto Company

3 7 3

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Manage before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 19986 Expires 12/31/00

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

Manage supplemental label revised 8/28/98 NEW
3

