

471025 50...

104...

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

FOR WEED CONTROL IN BANANAS FOR UP TO ONE DAY PRIOR TO HARVEST

Keep out of the reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.


This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used for weed control in established plantations of bananas. Applications may be made with boom equipment, shielded sprayers, hand-held and high-volume wands, lances and orchard guns, CDA equipment or wiper application equipment.



For specific rates of application and instructions, see the Weed Control Tables and the Tree and Vine Crops sections of the label booklet for Roundup Ultra herbicide.

Allow a minimum of 1 day between last application and harvest.

FOR DIRECT APPLICATION TO BANANAS (BANANACIDE), REMOVE FRUIT PRIOR TO TREATMENT.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

- 4f -

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup ULTRA herbicide and this supplement.

ROUNDUP® Ultra

HERBICIDE BY MONSANTO

EPA Reg. No. 524-475

SHIELDED APPLICATOR USE IN GRASSES GROWN FOR SEED IN IDAHO, OREGON AND WASHINGTON ONLY.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

When applied using shielded applicator equipment designed to prevent direct contact, movement of spray droplets, or mist onto desirable grasses grown for seed production, this product may be used to control labeled weeds. Use of low spray pressure through low pressure nozzles will minimize the potential of spray drift.

Apply 1 to 3 quarts of this product as a broadcast spray in 10 to 20 gallons of total spray volume per acre.

Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.



1002

**GROWER ASSUMES ALL RESPONSIBILITY FOR CROP LOSSES FROM
MISAPPLICATION.**

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Ultra before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1996

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

5201

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® PRO HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Pro herbicide and this supplement.

ROUNDUP PRO

HERBICIDE BY MONSANTO

EPA Reg. No. 524-475

ROUNDUP PRO HERBICIDE AND TANK MIXTURES FOR NONCROP AREAS, INCLUDING RAILROAD RIGHTS-OF-WAY, UTILITY SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES.

Keep out of reach of children.

CAUTIC

In case of emergency involving this product, call Collect, day or night, 314-694-4000.

®Roundup is a registered trademark of Monsanto Company.


Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the Roundup Pro label booklet for rate recommendations. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quarts per acre of this product may be used for improved results.



(10/1/77)

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product. Any recommended rate of this product may be used in a tank mixture with these products.

<u>Tank Mix Product</u>	<u>Rate per Acre</u>
Arsenal™	0.5 to 4 pints
Banvel	1 to 4 pints
2,4-D	0.5 to 1 pound
Garlon™ 3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar™ X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar™ I DF	4 to 6 pounds
Krovar I DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar I DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar I DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust™	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike™ 80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

*Arsenal is not approved for use in the state of California.

Refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of

7/20/96
all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank mix products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution which can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Pro before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

™Arsenal is a trademark of American Cyanamid Company.

™Banvel is a trademark of Sandoz Crop Protection Corporation.

™Garlon and Spike are trademarks of DowElanco Products Company.

™Hyvar, Krovar and Oust are trademarks of E. I duPont de Nemours and Company.

MONSANTO COMPANY 1996

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

10/10

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra herbicide and this supplement.

ROUNDUP® ULTRA

HERBICIDE BY MONSANTO

EPA Reg. No. 524-475

ROUNDUP ULTRA HERBICIDE FOR USE IN POPLAR (*Populus spp.*) PRODUCTION.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

See the Roundup Ultra herbicide label booklet for essential product performance information.


PREPLANT

This product is recommended for use prior to planting *Populus spp.* This includes, but is not limited to, hybrid poplars and hybrid cottonwoods.

See the Roundup Ultra herbicide label booklet for specific rates for the weeds being controlled.

IN CROP USES

1



9/24

DIRECTED SPRAYS

Use a 2 percent spray solution as a spray-to-wet application for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution. Avoid contact of spray, drift, or mist with foliage, green bark or non-woody surface roots of the *Populus* spp.

WIPERS

This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed on the label.

For wick applicators, mix 1 gallon of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force fed systems, a 33 to 100% Roundup Ultra solution may be used.

For best results ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.

AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION. Desirable vegetation contacted by the herbicide solution may be injured or controlled. This includes foliage, fruit, or green stems.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Ultra before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1996

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

10921

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® PRO HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Pro herbicide and this supplement.

ROUNDUP® PRO

HERBICIDE BY MONSANTO

EPA Reg. No. 524-475

ROUNDUP PRO HERBICIDE FOR USE IN POPLAR (*Populus spp.*) PRODUCTION.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup® is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

See the Roundup Pro herbicide label booklet for essential product performance information.

PREPLANT

This product is recommended for use prior to planting *Populus spp.* This includes, but is not limited to, hybrid poplars and hybrid cottonwoods.

See the Roundup Pro herbicide label booklet for specific rates for the weeds being controlled.

IN CROP USES

ACCEPTED
APR - 1 1996
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered with EPA Reg. No. 524-475

11/2/91

DIRECTED SPRAYS

Use a 2 percent spray solution as a spray-to-wet application for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution. Avoid contact of spray, drift, or mist with foliage, green bark or non-woody surface roots of the *Populus* spp.

WIPERS

This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed on the label.

For wick applicators, mix 1 gallon of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force fed systems, a 33 to 100% Roundup Pro solution may be used.

For best results ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.

AVOID HERBICIDE CONTACT WITH DESIRABLE VEGETATION. Desirable vegetation contacted by the herbicide solution may be injured or controlled. This includes foliage, fruit, or green stems.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Pro before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1996

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

Dup

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® PRO HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Pro herbicide and this supplement.

ROUNDUP PRO
HERBICIDE BY MONSANTO

EPA Reg. No. 524-475


FOR AERIAL APPLICATION IN CALIFORNIA ONLY.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, call Collect, day or night, 314-694-4000.

Roundup® is a registered trademark of Monsanto Company.



Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

See the label booklet for specific recommendations on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION, SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

When applied as recommended under the conditions described, this product controls annual and perennial weeds and woody brush and trees listed in the label booklet. See the Weeds Controlled section of the label booklet for rate recommendations.

Aerial applications of this product are allowed in the following situations:

1. Prior to the emergence or transplanting of labeled crops-
2. Aid to burning for establishment and maintenance of fuel breaks
3. Establishing fire perimeters and black lines
4. Aid to prescribed burning
5. Along fire roads
6. Range conversion
7. Habitat restoration and management
8. Wildlife food plots

~~DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR.~~

AERIAL APPLICATIONS

Apply 1 to 5 quarts of this product in 5 to 15 gallons of water per acre using aerial (helicopter only) applications.

To broaden the spectrum of control, Garlon 4 may be tank mixed with this product at the rate of 0.5 to 2 quarts per acre. The rate of Garlon should not exceed 1/2 the rate of this product (e.g. 1 quart of Garlon to 2 quarts of this product) for best results.

~~Use the recommended rates of this product in 3 to 15 gallons of water per acre.~~

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and

1/10/75

before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of ~~all~~ any desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within a minimum of 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.
5. APPLY BY AIR ONLY TO NONRESIDENTIAL AREAS.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application - To avoid streaking, uneven, or overlapped application, use appropriate marking devices.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear are most susceptible.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Pro herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Garlon™ is a trademark of DowElanco

MAP-

MONSANTO COMPANY 1995

MONSANTO COMPANY
AGRICULTURAL PRODUCTS

105201


ST. LOUIS, MISSOURI 63167

10/2/96

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.



ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

FOR DISTRIBUTION AND USE ONLY WITHIN ARKANSAS,
LOUISIANA, MISSISSIPPI, MISSOURI AND TENNESSEE

FOR CONTROL OF EMERGED WEEDS PRIOR TO THE PLANTING OR EMERGENCE OF CORN, COTTON, RICE, SORGHUM AND SOYBEANS; PRIOR TO THE HARVEST OF COTTON AND SOYBEANS; AND FOLLOWING THE HARVEST OF ANY CROP* IN THE FALL VIA AERIAL APPLICATIONS

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

- ® Roundup is a registered trademark of Monsanto Company.
- * Do not plant subsequent crops other than those on the label for 30 days following application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

11/2/81

This product controls annual and perennial weeds listed in the label booklet for Roundup Ultra prior to planting or emergence of corn, cotton, rice, sorghum and soybeans; prior to the harvest of cotton and soybeans; and following the harvest of any crop in the fall via aerial applications.

Aerial applications of this product may be made in fallow systems and conventional, reduced and zero tillage systems.

For applications via aerial equipment, use the recommended rates of this product in 3 to 10 gallons of water per acre. Do not exceed a rate of 3 quarts per acre.

AVOID DRIFT - DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser velocities, will allow spray drift to occur.

Read the "AERIAL EQUIPMENT" section of the Roundup Ultra label booklet for additional precautions and instructions.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475



FOR USE IN DORMANT RANGELAND

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product will control or suppress many weeds, including downy brome, cheat grass, cereal rye, medusahead rye and jointed goatgrass in dormant rangeland.

17/2/01

Apply 8 to 16 fluid ounces of this product per acre in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass, are still truly dormant.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

Do not use ammonium sulfate when spraying dormant rangeland grasses with this product.

Do not make more than one application per year.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

3/24/96

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED
APR - 1 1996
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

FOR RED RICE CONTROL IN TEXAS ONLY

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used to control red rice, prior to planting rice. Apply 1 1/2 quarts of this product in 5 to 10 gallons of water per acre.

Flush fields prior to applications to obtain uniform germination and stand of red rice. Make applications when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2

true leaves may only be partially controlled. Do not treat rice fields or levees when the fields contain flood water.

Avoid spraying during low humidity conditions, as reduced control may result.

Do not re-flood treated fields for 8 days following application.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167