

# Roundup ULTRA

Herbicide by **Monsanto**

**AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

Carefully follow detailed instructions in label booklet.

Read the entire label before using this product. Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY," which appears in the label booklet, before buying or using. If terms are not acceptable, return at once unopened.

The contents of this container may not be REFORMULATED, nor, except in accordance with the Monsanto Repackaging Agreement, REPACKAGED.

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

### CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

**FIRST AID: IF IN EYES,** flush with plenty of water. Get medical attention if irritation persists.

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep

and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

### Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**DISPOSAL:** Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

Do not reuse this container, except for refill in accordance with a valid Monsanto Repackaging or Toll Repackaging Agreement. If not refilled or returned to an authorized repackaging facility, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(See label booklet for Complete Directions for Use)

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt . . . . . 41.0%  
INERT INGREDIENTS: . . . . . 59.0%  
100.0%

\*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

For product information or assistance in using this product, Call toll-free, 1-800-332-3111.

\*Roundup is a registered trademark of Monsanto Company  
©MONSANTO COMPANY 1996

**M** MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167 U.S.A.

EPA Reg. No. 524-475

21137U2-7/53-1

**ACCEPTED**  
OCT 23 1996  
Under the Federal Insecticide, Fungicide, and Rodenticide Act.  
as amended by the pesticide  
regulatory reform  
Act Reg. No. 524-475

PLACE  
Complete Directions For Use  
Booklet  
HERE

'96 SEP -6 A9:22

RECD EPA/DP/DPD1

NET CONTENTS

EPA Est.

6-109


524-475

25  
7M 26 WK

# SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra, container labels and this supplement.



96 AUG 23 P12:25

RECD EPA/OPP/DPD1

EPA Reg. No. 524-475

## FOR POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE

### ROUNDUP READY® GENE

Keep out of reach of children.

### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Roundup® and Roundup Ready® Gene are registered trademarks of Monsanto Company.  
Roundup Ready™ is a trademark of Monsanto Company.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup® Ultra herbicide for essential product performance information.

**NOTE: Do not feed or graze treated soybean forage.**

#### General Information

FOR POSTEMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES WHICH HAVE THE ROUNDUP READY® GENE.

•Applying this product to soybean varieties which are not designated as Roundup Ready™ will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene, since severe injury or destruction will result.

•Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Monsanto when this product is used in conjunction with "brown bag" or "bin run" soybean seed saved from previous year's production and replanted.

•The Roundup Ready designation indicates that the soybean contains a patented gene which provides tolerance to Monsanto's Roundup brand herbicides. Information on Roundup Ready soybeans may be obtained from your seed supplier or Monsanto representative.

#### Application Instructions

This product may be applied postemergence to Roundup Ready Soybeans from the cracking stage through the full flowering stage.

Allow a minimum of 14 days between application and harvest of soybeans.

#### Maximum Allowable Yearly Rates

- |  |            |
|--|------------|
| 1. Combined total per year for all applications                                | 8 quarts/A |
| 2. Preplant, Preemergence applications   | 5 quarts/A |
| 3. Total of single or multiple in-crop applications from cracking to flowering | 2 quarts/A |
| 4. Maximum preharvest application rate   | 1 quart/A  |

When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready Soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product.

There are no rotational crop restrictions following applications of this product.

**For ground applications:** Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**For aerial applications:** Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of this product per acre. DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

AERIAL APPLICATIONS ON ROUNDUP READY SOYBEANS, MAY BE MADE ONLY IN THE FOLLOWING STATES: ALABAMA, ARKANSAS, FLORIDA, GEORGIA, LOUISIANA, MISSISSIPPI, MISSOURI (BOOTHEEL ONLY), NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE AND TEXAS.

**ANNUAL WEED RATE TABLES**

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the label booklet for rate recommendations for specific annual weeds.

Tank mixtures with other herbicides are not recommended due to potential for crop injury and/or weed antagonism and to rotational crop restrictions of the tank mix partner.

This product may be used up to 64 fluid ounces per acre for control of annual weeds, where heavy weed densities exist.

**NOTE:** In no-till and stale seedbed systems, always burn down existing weeds before soybeans emerge. Apply a preplant burndown treatment of 16-48 fluid ounces per acre of this product.

**MIDWEST RECOMMENDATIONS**

**Narrow row or drilled soybeans:** A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fl oz/A)
0-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16-24 fluid ounces per acre may be necessary to control late flushes of weeds.

**Wide row soybeans:** A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

<u>Initial Treatment</u>	
Weed Height (inches)	Rate (fl oz/A)
0-6	24
6-12	32
12-18	48

<u>Sequential Application*</u>	
Weed Height (inches)	Rate (fl oz/A)
0-3	16
3-6	24

\*Total application not to exceed 64 fluid ounces per acre.

For morningglory, black nightshade, groundcherry apply:

Weed Height (inches)	Rate (fl oz/A)
0-3	24
3-6	32
6-12	48

**Giant ragweed:** Apply 32 fl oz/A when the weed is 8-12" tall to avoid the need for sequential application.

Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, and giant ragweed, with multiple germination times may require a sequen-

tial application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments (not including preharvest) must not exceed 64 fluid ounces per acre.

**MID-ATLANTIC/SOUTHEAST RECOMMENDATIONS**

**Narrow row or drilled soybeans:** A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fl oz/A)
0-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16-24 fluid ounces per acre may be necessary to control late flushes of weeds.

**Wide row soybeans:** A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

<u>Initial Treatment</u>	
Weed Height (inches)	Rate (fl oz/A)
0-6	24
6-12	32
12-18	48

<u>Sequential Application*</u>	
Weed Height (inches)	Rate (fl oz/A)
0-3	16
3-6	24

\*Total application not to exceed 64 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

For morningglory, black nightshade, groundcherry apply:

Weed Height (inches)	Rate (fl oz/A)
0-3	24
3-6	32
6-12	48

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum and sicklepod, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments (not including preharvest) must not exceed 64 fluid ounces per acre.

**DELTA/MID-SOUTH RECOMMENDATIONS**

**Narrow row or drilled soybeans:** An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds.

**Wide row soybeans:** An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. Cultivation, as needed, may be necessary to control late flushes of weeds.

Initial Treatment

Weed Height (inches)	Rate (fl oz/A)
0-3	24
3-6	32
6-12	48

Sequential Application\*

Weed Height (inches)	Rate (fl oz/A)
0-3	16
3-6	24

\*Total application not to exceed 64 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

**PERENNIAL WEEDS RATE RECOMMENDATIONS**

A 32 to 64 fluid ounces per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horse-nettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for Roundup Ultra. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©Monsanto Company 1996

MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167

21137U2-12/53

7/31/96

4089

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

**ACCEPTED**  
OCT 23 1996  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra, container labels and this supplement.

Roundup Ultra Herbicide

EPA Reg. No. 524-475

REC'D EPA/OPP/DPD1  
96 SEP 19 P2:49

FOR POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE

ROUNDUP READY® GENE

Keep out of reach of children.

**CAUTION!**

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Roundup® and Roundup Ready® Gene are registered trademarks of Monsanto Company.  
Roundup Ready™ is a trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application. See "GENERAL INFORMATION" and "APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup® Ultra herbicide for essential product performance information.

~~NOTE: Do not feed or graze treated soybean forage.~~

General Information

FOR POSTEMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES WHICH HAVE THE ROUNDUP READY® GENE.

- Applying this product to soybean varieties which are not designated as Roundup Ready™ will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene, since severe injury or destruction will result.
- Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Monsanto when this product is used in conjunction with "brown bag" or "bin run" soybean seed saved from previous year's production and replanted.
- The Roundup Ready designation indicates that the soybean contains a patented gene which provides tolerance to Monsanto's Roundup brand herbicides. Information on Roundup Ready soybeans may be obtained from your seed supplier or Monsanto representative.

**Application Instructions**

This product may be applied postemergence to Roundup Ready soybeans from the cracking stage through the full flowering stage.

Allow a minimum of 14 days between application and harvest of soybeans.

**Maximum Allowable Yearly Rates**

- Preplant: Maximum amount of this product which can be applied prior to crop emergence is 5 quarts/A.
- In-crop: Maximum combined total of single or multiple in-crop applications from cracking to flowering is 3 quarts/A and the maximum rate for any single in-crop application is 2 quarts/A.
- Preharvest: Maximum amount of this product that can be applied after loss of green color in soybean pods until 14 days before harvest is 1 quart/A. The combined total of preharvest and in-crop applications may not exceed 3 quarts/A.
- Cropping Season: Combined total per year for all applications may not exceed 8 quarts/A.

**Maximum Allowable Yearly Rates**

1. Combined total per year for all applications	8 quarts/A
2. Preplant, Preemergence applications	5 quarts/A
3. Total of single or multiple in-crop applications from cracking to flowering	2 quarts/A
4. Maximum preharvest application rate	1 quart/A

When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product.

There are no rotational crop restrictions following applications of this product.

**For ground applications:** Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles which provide a flat fan pattern~~flat fan nozzles~~. Check for even distribution of spray droplets.

**For aerial applications:** Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of this product per acre. DO NOT APPLY DURING LOW LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

AERIAL APPLICATIONS ON ROUNDUP READY SOYBEANS, MAY BE MADE ONLY IN THE FOLLOWING STATES: ALABAMA, ARKANSAS, FLORIDA, GEORGIA, KANSAS, LOUISIANA, MISSISSIPPI, MISSOURI (BOOT-HEEL ONLY), NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE AND TEXAS.

**ANNUAL WEED RATE TABLES**

The following rate recommendations will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the label booklet for rate recommendations for specific annual weeds.

Tank mixtures with other herbicides are not recommended due to potential for crop injury and/or weed antagonism and to rotational crop restrictions of the tank mix partner.

This product may be used up to 64 fluid ounces per acre in any single application for control of annual weeds, where heavy weed densities exist.

**NOTE:** In no-till and stale seedbed systems, always burn down existing weeds before soybeans emerge. Apply a preplant burn-down treatment of 16-6448 fluid ounces per acre of this product.

**MIDWEST RECOMMENDATIONS**

**Narrow row or drilled soybeans:** A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fl oz/A)
3-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soy-bean stand that slows or delays canopy closure, a sequential application of this product at 16-3224 fluid ounces per acre may be necessary to control late flushes of weeds.

**Wide row soybeans:** A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

Initial Treatment

Weed Height (inches)	Rate (fl oz/A)
3-6	24
6-12	32
12-18	48

Sequential Application\*

Weed Height (inches)	Rate (fl oz/A)
1-3	16
3-6	24
6-12	32

\*Combined total application in-crop not to exceed 9664 fluid ounces per acre.

For morningglory, black nightshade, groundcherry, and Pennsylvania smartweed apply:

Weed Height (inches)	Rate (fl oz/A)
1-3	24
3-6	32
6-12	48

Giant ragweed: Apply 32 fl oz/A when the weed is 8-12" tall to avoid the need for sequential application.

Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments (not including preharvest)-must not exceed 9664 fluid ounces per acre.

MID-ATLANTIC/SOUTHEAST RECOMMENDATIONS

Narrow row or drilled soybeans: A single in-crop application of this product will provide effective control of labeled weeds.

Weed Height (inches)	Rate (fl oz/A)
3-6	24
6-12	32
12-18	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this product at 16-3224 fluid ounces per acre may be necessary to control late flushes of weeds.

Wide row soybeans: A single in-crop application of this product will provide effective control of initial stand of labeled weeds. If new flushes of weeds occur, they can be controlled by sequential applications of this product.

Initial Treatment

Weed Height (inches)	Rate (fl oz/A)
3-6	24
6-12	32
12-18	48

Sequential Application\*

Weed Height (inches)	Rate (fl oz/A)
1-3	16
3-6	24
6-12	32

\*Combined total application in-crop not to exceed 9664 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

For morningglory, black nightshade, groundcherry, and Pennsylvania smartweed apply:

Weed Height (inches)	Rate (fl oz/A)
10-3	24
3-6	32
6-12	48

Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of this product. Suppressed or stunted weeds may also require sequential applications. Sequential applications should be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of this product per acre for sequential applications. The combined total of all in-crop postemergence treatments (not including preharvest) must not exceed 9664 fluid ounces per acre.

### DELTA/MID-SOUTH RECOMMENDATIONS

**Narrow row or drilled soybeans:** An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds.

**Wide row soybeans:** An in-crop application of this product will provide effective control of initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. Cultivation, as needed, may be necessary to control late flushes of weeds.

#### Initial Treatment

Weed Height (inches)	Rate (fl oz/A)
10-3	24
3-6	32
6-12	48

#### Sequential Application\*

Weed Height (inches)	Rate (fl oz/A)
10-3	16
3-6	24
6-12	32

\*Combined Total application in-crop not to exceed 9664 fluid ounces per acre.

Hemp sesbania: Apply 24 fl oz/A at 0-2" tall; 32 fl oz/A at 2-4" tall; 40 fl oz/A at 4-6" tall; 48 fl oz/A at 6-8" tall.

### PERENNIAL WEEDS RATE RECOMMENDATIONS

A 32 to 64 fluid ounces per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for

9099

Roundup Ultra. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©Monsanto Company 1996  
MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167

9/16/96 |