

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra, container labels and this supplement.

ROUNDUP® ULTRA

Herbicide by Monsanto

EPA Reg. No. 524-475

FOR POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company. Roundup Ready® is a registered trademark of Monsanto Company.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

~~NOTE: THIS PRODUCT MAY BE USED ONLY OVER THE TOP OF IMPROVED CORN VARIETIES THAT ARE DESIGNATED AS CORN WITH THE ROUNDUP READY® GENE. SEVERE INJURY OR DEATH OF CORN WILL RESULT IF ANY CORN VARIETIES NOT DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES OTHER THAN CROPS WITH THE ROUNDUP READY GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.~~

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.



GENERAL INFORMATION

MONSANTO COMPANY RECOMMENDS USE OF THIS PRODUCT ONLY ON CORN SEED DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

• Applying this product to corn varieties which are not designated as Roundup Ready® will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready gene since severe injury or destruction will result.

• Roundup Ready varieties must be purchased from an authorized seed supplier. Crop safety and weed control performance are not warranted by Monsanto when this product is used in conjunction with "brown bag" or "bin run" corn seed saved from previous year's production and replanted.

• The Roundup Ready designation indicates that the corn contains a patented gene which provides tolerance to Monsanto's Roundup brand herbicides. Information on Roundup Ready corn may be obtained from your seed supplier or Monsanto representative.

Application Instructions

This product may be applied postemergence to Roundup Ready corn during the period beginning at corn emergence and continuing through the 12-leaf stage or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of this product are not to exceed 1 quart per acre. Sequential in-crop applications of this product should not exceed 2 quarts per growing season. Total Roundup Ultra use should not exceed 8 quarts per acre per year.

Allow a minimum of 50 days between application of this product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 14 days between in-crop applications of this product. THE USE OF ADDITIVES FOR IN-CROP APPLICATIONS TO ROUNDUP READY CORN IS PROHIBITED.

Maximum Yearly Rates Allowed

Preplant / Preemergence (Maximum)	5 quarts/A
Total in-crop applications from emergence to	
12-leaf stage or 30 inches	2 quarts/A
Maximum preharvest rate.	1 quart/A
<hr/>	
Combined total per year for all applications.	8 quarts/A

When applied as directed, this product controls annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. Applications should be made to actively growing weeds before they reach the maximum size listed in the "WEEDS CONTROLLED" section of the label booklet. Refer to the "MIXING" section of the label booklet for proper use instructions.

There are no rotational crop restrictions following applications of this product.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto other desirable vegetation since minute quantities of this product can cause severe damage or destruction to crop plants in non-target areas. The likelihood of plant injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions which allow spray drift to occur such as combinations of pressure and nozzle type that will result in fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR SPRAY PRESSURE.**

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. See "WEEDS CONTROLLED" section of this label for specific recommended rates. Carefully select proper nozzles and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart per acre. See "WEEDS CONTROLLED" section on this label. **AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN**

4/2/12

WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Weeds controlled

For specific rates of application and instructions for control of various annual and perennial weeds, refer to the label booklet. Roundup Ultra applied at up to 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: nutsedge, rhizome johnsongrass, quackgrass, Canada thistle, wirestem muhly.

Sequential Applications: Some weeds with multiple germination times or suppressed (stunted) weeds may required a second application of this product. The second application should be made after some regrowth has occurred.

Tank mixtures

A tank mixture of Roundup Ultra plus Micro-Tech or Partner may be used for postemergence and residual control of annual weeds in corn. These tank mixtures may be made during the period beginning at corn emergence and continuing until corn height reaches 5 inches.

A Roundup Ultra tank mixture with atrazine, Banvel, Clarity, Permit or 2,4-D may be used for postemergence control of additional annual weeds in corn. A Roundup Ultra tank mixture with atrazine may be made during the period beginning at corn emergence and continuing until corn height reaches 12 inches. A Roundup Ultra tank mixture with Banvel or Clarity at 0.125 to 0.25 lb per acre may be made during the period beginning at corn emergence and continuing until corn height reaches 30 inches. A Roundup Ultra tank mixture with Permit may be made during the period beginning at corn emergence and continuing until corn height reaches 30 inches. A Roundup Ultra tank mixture with 2,4-D at 0.125 to 0.25 lb per acre may be made during the period beginning at corn emergence and continuing until corn is at the five leaf stage or corn height reaches 8 inches, whichever comes first.

Refer to the specific product label and observe all precautions, mixing and application instructions for all products used in tank mixtures.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997/1996

MAP

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

50/12

'96 JUN 26 P12-22

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA RT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra RT and this supplement.

ROUNDUP ® ULTRA RT

Herbicide by Monsanto

EPA Reg. No. 524-475

<p>ACCEPTED MAR 28 1997</p>	<p>Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. <u>524-475</u></p>
--	--

PREHARVEST APPLICATIONS IN CORN

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra RT herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of corn to control annual and perennial weeds listed on the label booklet for Roundup Ultra RT herbicide.

For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 1 quart of this product per acre.

60112

Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). Allow a minimum of 7 days between application and harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that corn grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra RT herbicide before using. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

7012

RECD EPA/017/DPD2

'96 JUN 26 P12:14 **SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

PREHARVEST APPLICATIONS IN CORN

ACCEPTED
MAR 28 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLLAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of corn to control annual and perennial weeds listed on the label booklet for Roundup Ultra herbicide.

For ground applications, apply up to 3 quarts of this product per acre. For aerial applications, apply up to 1 quarts of this product per acre.

8942

Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). Allow a minimum of 7 days between application and harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that corn grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

90112

RECD EPA/OPP/DPD2

SUPPLEMENTAL LABELING

'96 JUN 26 P12 :09

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA

Herbicide by Monsanto

EPA Reg. No. 524-475

PREHARVEST APPLICATIONS IN GRAIN SORGHUM (MILO)

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of grain sorghum (milo) to control annual and perennial weeds listed on the label booklet for Roundup Ultra herbicide.

Apply up to 2 quarts of this product per acre.

ACCEPTED
MAR 28 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

10/12

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that grain sorghum grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

110112

RECD EPA/OTR/DPD2

SUPPLEMENTAL LABELING

'96 JUN 26 P12 :25

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA RT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra RT and this supplement.

ROUNDUP ® ULTRA RT
Herbicide by Monsanto

EPA Reg. No. 524-475

PREHARVEST APPLICATIONS IN GRAIN SORGHUM (MILO)

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra RT herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of grain sorghum (milo) to control annual and perennial weeds listed on the label booklet for Roundup Ultra RT herbicide.

Apply up to 2 quarts of this product per acre.

ACCEPTED
MAR 28 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

Make applications at 30% grain moisture or less and at least 7 days prior to harvest.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for instructions.

Note: It is not recommended that grain sorghum grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra RT herbicide before using. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167