

PM 25

524-475

8/26/97

page 1 of 2

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

FOR USE IN PASTURES FOR HAWAII ONLY

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used in Hawaiian pastures, including kikuya grass, pangola grass and guineagrass pastures. Spot treatment, wiper application, preplant, preemergence and pasture renovation applications may be made.

ACCEPTED

AUG 26 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 524-475

Spot treatment

This product may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals. For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

Preplant, preemergence and pasture renovation

When applied prior to planting or emergence of forage grasses or as a direct application to the listed perennial pasture species, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

See the Weed Control Tables in the label booklet for Roundup Ultra for information about rates and timings for specific weeds.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

MAP


MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP® ULTRA
Herbicide by Monsanto
EPA Reg. No. 524-475



USE ONLY FOR SEED PRODUCTION OF CANOLA WITH THE ROUNDUP READY® GENE

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup and Roundup Ready are registered trademarks of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN CANOLA WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT CANOLA IN PRODUCTION FIELDS OF CANOLA CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF CANOLA VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE RECOMMENDATIONS

This product will control non-glyphosate tolerant canola in seed production fields of canola containing the Roundup Ready gene. This product may be applied using ground spray equipment only. Apply 1 pint of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application of 1 pint per acre may be applied, if needed to control non-glyphosate tolerant canola plants.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART OF THIS PRODUCT PER ACRE PER SEASON.

Application timing -- This product can be applied to Roundup Ready canola from emergence to the pre-flower (early bolting) stage.

Treated canola may not be used for food or feed. Do not feed or graze treated canola. Do not process treated canola for food or feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED
AUG 26 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

ROUNDUP® ULTRA HERBICIDE FOR USE IN DORMANT ALFALFA

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product. Call Collect, day or night. 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES. BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product will control or suppress many weeds, including quackgrass, downy brome and cheatgrass in dormant alfalfa.

Apply 8 to 12 ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliolate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliolate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with Roundup Ultra.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product is limited to persons who have attended a Monsanto-approved training program. Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA RT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra RT and this supplement.

ROUNDUP® ULTRA RT
Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED
AUG 26 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

ROUNDUP® ULTRA RT HERBICIDE FOR USE DORMANT ALFALFA

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS. DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra RT herbicide for essential product performance information.

RECOMMENDATIONS

This product will control or suppress many weeds, including quackgrass, downy brome and cheatgrass in dormant alfalfa.

Apply 8 to 12 ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliolate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliolate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with Roundup Ultra RT.

Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product is limited to persons who have attended a Monsanto-approved training program. Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra RT herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1996

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

PM

25

524-475

8/26/97

page 172

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED
AUG 26 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

FOR PISTACHIOS UP TO THREE DAYS PRIOR TO HARVEST

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product is recommended for weed control in established groves of pistachios. Applications may be made with boom equipment, shielded sprayers, hand-held and high volume wands, lances and orchard guns, CDA equipment or with wiper application equipment.

REC: P/DPD1
97 MAR 7 P 3:41

2. 9 2

For specific rates of application and instructions, see the Weed Control Tables and Tree Nuts sections of the label booklet for Roundup Ultra herbicide.

Allow a minimum of 3 days between last application and harvest.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

ROUNDUP® W

HERBICIDE BY

Monsanto

ACCEPTED

AUG 26 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

Read the entire label before using this product. Use only according to label instructions. The contents of this container may not be REFORMULATED, nor, except in accordance with the Monsanto Repackaging Agreement, REPACKAGED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

FIRST AID: IF IN EYES, flush with plenty of water. Get medical attention if irritation persists.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if

ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

For more product information, call toll-free 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

FOR REPACKAGING INTO HERBICIDES ONLY.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its

isopropylamine salt:	41.0%
INERT INGREDIENTS:	59.0%
	100.0%

*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

©MONSANTO COMPANY 1997

®Registered trademark of Monsanto Company.

EPA Reg. No. 524-475

EPA Est. 524-LA-1

EPA Est. 524-IA-1

EPA Est. 524-NC-1



LOT NO.

PACKER

NET

GAL.

MONSANTO COMPANY, ST. LOUIS, MISSOURI 63167, U.S.A.

21145V1-1X3

PM 25 524-475 8/26/97 page 1 of 1

BACKPACK IDENTIFICATION STICKER FOR



Keep out of reach of children.

CAUTION!

ACTIVE INGREDIENT:
Glyphosate, N-(phosphonomethyl)glycine

EPA Reg. No. 524-475

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetables since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist), which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

DO NOT USE FOR STORAGE.

This container must be used only for Roundup Pro or other non-selective herbicides, or must be thoroughly cleaned and rinsed. Rinsate must be used in accordance with total directions or disposed of in accordance with applicable law. Any product residue in the container may result in injury to desirable plants and crops.

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

For more product information, call toll-free 1-800-332-3111.

*Roundup is a registered trademark of Monsanto Company.

User must always read, follow and have in possession the current "COMPLETE DIRECTIONS FOR USE" label booklet for Roundup Pro herbicide. It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Refer to these DIRECTIONS for PRECAUTIONARY STATEMENTS, FIRST AID INSTRUCTIONS, ENVIRONMENTAL HAZARDS and PHYSICAL OR CHEMICAL HAZARDS.

Below is an abbreviated listing of weeds controlled and proper use rates from the full Roundup Pro herbicide label. Use this as a guide to mixing appropriate amounts of Roundup Pro herbicide and water to control the common labeled weeds and brush listed. Remember, Roundup Pro herbicide is a non-selective post-emergent herbicide for the control of annual and perennial grasses, broadleaves and brush species.

Annual Weeds*		Perennial Weeds*		Brush*
1% Solution		1 1/2-2% Solution		2% Solution
Annual bluegrass	Marestail	Artichoke thistle	Ice plant	French broom
Annual ryegrass	Mustard	Bermudagrass	Johnsongrass	Mullein
Barnyardgrass	Pigweed	Clover	Kikuyugrass	Oak
Chickweed	Ragweed	Dallisgrass	Pampasgrass	Poison oak
Crabgrass	Russian thistle	Dandelion	Perennial ryegrass	Tree tobacco
Fixes	Sowthistle	Field bindweed	Poison hemlock	Willows
Foxgall	Spotted spurge	Fescue		
London rocket	Wild oats	Giant seed		

*For complete list of species controlled, check the label.

SPRAY SOLUTION

Desired Volume	Amount of Roundup*Pro		
	1%	1 1/2 %	2%
1 gallon	1 1/3 oz.	2 oz.	2 2/3 oz.
25 gallons	1 qt.	1 1/2 qts.	2 qts.
100 gallons	1 gal.	1 1/2 gals.	2 gals.

General Directions:

Roundup Pro herbicide, a water soluble liquid, mixes readily with water to be applied as a foliage spray for the control of undesirable plants. Apply on a spray-to-wet basis.

21136V1-1/53

RESPONSIBLE PARTY: _____ DILUTION: _____

ADDRESS: _____

ACCEPTED

AUG 26 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, we are notified for the pesticide registered under EPA Reg. No. 524-475

pm 25

524-475

8/26/97

page 181

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP® ULTRA

Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED

AUG 26 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 524-475

USE ONLY FOR SEED PRODUCTION OF SUGARBEETS WITH THE ROUNDUP READY®
GENE

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup and Roundup Ready are registered trademarks of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN SUGARBEETS WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT SUGARBEETS IN PRODUCTION FIELDS OF SUGARBEETS CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF SUGARBEETS VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE RECOMMENDATIONS

This product will control non-glyphosate tolerant sugarbeets in seed production fields of sugarbeets containing the Roundup Ready gene. This product may be applied using ground spray equipment only. Apply 24 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application of 24 ounces per acre may be applied, if needed to control non-glyphosate tolerant sugarbeets plants.

DO NOT EXCEED A MAXIMUM RATE OF 1 1/2 QUARTS OF THIS PRODUCT PER ACRE PER SEASON.

Application timing -- This product can be applied to Roundup Ready sugarbeets from emergence to canopy closure.

Treated sugarbeets may not be used for food or feed. Do not feed or graze treated sugarbeets. Do not process treated sugarbeets for food or feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

MAP

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

MON 65005 Forestry Herbicide

by Monsanto

Complete Directions for Use

EPA Reg. No. 524-475

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NONWOODY ROOTS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

1997

Read the entire label before using this product.

Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.


REFORMULATION OR REPACKAGING IS PROHIBITED.

Ingredients

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt	41.0%
INERT INGREDIENTS (including 14.5% surfactant):	59.0%
	<u>100.0%</u>

*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine



salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

PRECAUTIONARY STATEMENTS

Hazards to
Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

FIRST AID: IF IN EYES, flush with plenty of water. Get medical attention if irritation persists.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)

(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, waterproof gloves, shoes plus socks.

For more product information, call toll free 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

DISPOSAL: Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

REUSE: Do not reuse this container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous and woody plants.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most herbaceous weeds occur within 7 days but on most woody plants may not occur for 30 days or more.

After any site disturbance, such as logging, mechanical brush removal or mowing, allow stump sprouts, resprouts and foliar regrowth from woody brush and perennial herbaceous weeds sufficient time to regrow before treatment.

Always use the higher recommended rates of this product when treating dense, multicanopied sites of woody vegetation or difficult-to-control woody and herbaceous plants.

Reduced control may result when woody brush, trees and herbaceous weeds are treated under poor growing conditions caused by drought, disease or insect damage. Reduced control may result if the foliage of undesirable vegetation is covered with dust at the time of treatment.

Heavy rainfall soon after application may wash this product off the foliage and a repeat application may be required for adequate control.

This product has no herbicidal or residual activity in the soil. Weed must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide.

When this product comes in contact with the soil it is bound to soil particles. When used in accordance with label directions, once this product is bound it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treatment area or if the soil is transported off-site. The strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water. The affinity between this product and soil particles remains until this product is degraded, which is primarily a biological degradation process carried out under both aerobic and anaerobic conditions by soil microflora.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended in this label may result in reduced performance.

FORESTRY SITE PREPARATION

This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

7 7 26

APPLICATION RATES AND TIMING

APPLICATION	MON 65005	SPRAY VOLUME GAL/A
<u>BROADCAST</u>		
Aerial	2 to 10 qts/a	5 to 30
Ground	2 to 10 qts/a	10 to 60
<u>SPRAY-TO-WET</u>		
Handgun, Backpack, Mistblower	3/4 % to 2 % by volume	spray-to-wet
<u>LOW VOLUME DIRECTED SPRAY</u>		
Handgun, Backpack, Mistblower	5% to 10% by volume	partial coverage*

*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

TANK MIXTURES

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any recommended rate of this product may be used in a tank mix with the following products for forestry site preparation.

PRODUCT	BROADCAST RATE
Arsenal Applicators Concentrate	2 to 16 fl oz /a
Escort™	½ to 3 ½ oz/a
Oust™	1 to 4 oz/a
Garlon 4	1 to 4 qts/a

PRODUCT	SPRAY-TO-WET RATES
Arsenal Applicators Concentrate	1/32 % to 1/2 % by volume

PRODUCT	LOW VOLUME DIRECTED SPRAY RATES
Arsenal Applicators Concentrate	1/8% to 1/2% by volume

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

™Arsenal is a trademark of American Cyanamid Company.

™Oust is a trademark of E. I. du Pont de Nemours and Company.

™Garlon is a trademark of DowElanco Products Company.

9 8 26

FORESTRY CONIFER AND HARDWOOD RELEASE

DIRECTED SPRAY AND SELECTIVE EQUIPMENT

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries.

In hardwood plantations, tank mixtures with Oust may be used. In pine plantations, tank mixtures with Garlon 4 or Arsenal AC may be used. Comply with all site restrictions, forestry species limitations and precautions on the tank mix product label.

Avoid contact of spray, drift, mist or drips with foliage, green bark or non-woody surface roots of desirable species.

See all sections in the APPLICATION EQUIPMENT AND TECHNIQUES portion of this label for specific equipment recommendations and precautions.

For spray-to-wet applications, use a 2 percent spray solution for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution.

For low volume directed spray applications, use a 5 to 10 percent spray solution. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

For equipment calibrated for broadcast applications, use 2 to 10 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Shielded application equipment may be used to avoid contact of the spray solution with desirable plants. Shields should be adjusted to prevent spray contact with the foliage or green bark of desirable vegetation.

Wiper application equipment may be used. See the SELECTIVE EQUIPMENT portion of this label for equipment and rate recommendations

10 7 26

DO NOT APPLY THIS PRODUCT AS AN OVER-THE-TOP BROADCAST SPRAY FOR FORESTRY CONIFER OR HARDWOOD RELEASE.

**MIXING AND APPLICATION
INSTRUCTIONS**

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. HANDGUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. **NOTE:** REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

**APPLICATION EQUIPMENT
AND TECHNIQUES**

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this

product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of non-target injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

■ **AERIAL EQUIPMENT** ■

This product is recommended for aerial application in forestry sites by helicopter only. Use the recommended rates of this product in 5 to 30 gallons of spray solution per acre as a broadcast spray. (THIS PRODUCT PLUS OUST TANK MIXTURES MAY NOT BE APPLIED BY AIR IN CALIFORNIA.)

AVOID DRIFT: DO NOT APPLY DURING INVERSION CONDITION, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT; DRIFT WILL CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine droplets.

Drift control additives may be used for forestry site preparation. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace

specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear are most susceptible.

■ **GROUND BROADCAST EQUIPMENT** ■

This product is recommended for broadcast applications using suitable ground equipment in forestry sites. Use the recommended rates of this product in 10 to 60 gallons of clean water per acre as a broadcast spray. Check for even spray distribution throughout the spray pattern.

■ **BACKPACK, HANDGUN OR MISTBLOWER EQUIPMENT** ■

This product is recommended for application through backpack, handgun or hand-held mistblower* equipment. For spray-to-wet applications, coverage should be uniform and complete, but not to the point of runoff.

This product can be used for low volume directed sprays for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zigzag motion. For flat fan and cone nozzles and with mistblowers, mist the application over the foliage of the targeted vegetation. Small, open branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

*This product is not registered in California or Arizona for use in mistblowers.

■ **SELECTIVE EQUIPMENT** ■

This product may be applied through shielded sprayers or wiper application equipment. When properly used this equipment may

selectively control undesirable vegetation without harming desirable vegetation.

Shielded sprayers direct the herbicide solution onto weeds while shielding desirable vegetation from the spray solution. Any recommended rate or tank mixture of this product may be used employing this equipment.

Wiper applicators physically wipe product directly onto undesirable vegetation. Care should be taken to avoid wiping desirable vegetation. Use a 33 to 100 percent solution of this product, diluted in water for wiper applications. Use a 33 percent solution for wick or gravity feed systems. Higher concentrations may be used in pressurized systems that are capable of handling thicker solutions.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product CONTROLS, PARTIALLY CONTROLS or SUPPRESSES most woody brush, trees and herbaceous weeds, some of which are listed below.

WOODY BRUSH AND TREES

Alder

Alnus spp.

Ash

Fraxinus spp.

Aspen, quaking

Populus tremuloides

Bearmat (Bearclover)

Chamaebatia foliolosa

Beech

Fagus grandifolia

Birch

Betula spp.

Blackberry
Rubus spp.

Blackgum
Nyssa spp.

Bracken
Pteridium spp.

Broom:
French
Cytisus monspessulanus
Scotch
Cytisus scoparius

Buckwheat, California
Eriogonum fasciculatum

Cascara
Rhamnus purshiana

Catsclaw
Acacia greggi

Ceanothus
Ceanothus spp.

Chamise
Adenostoma fasciculatum

Cherry:
Bitter
Prunus emarginata

Black
Prunus serotina

Pin
Prunus pensylvanica

Coyote brush
Baccharis consanguinea

Creeper, Virginia

Parthenocissus quinquefolia

Dewberry
Rubus trivialis

Dogwood
Cornus spp.

Elderberry
Sambucus spp.

Elm
Ulmus spp.

Eucalyptus, bluegum
Eucalyptus globulus

Hasardia
Haplopappus squamosus

Hawthorn
Crataegus spp.

Hazel
Corylus spp.

Hickory
Carya spp.

Holly, Florida; Brazilian Peppertree
Schinus terebinthifolius

Honeysuckle
Lonicera spp.

Hornbeam, American
Carpinus caroliniana

Kudzu
Pueraria lobata

Locust, black
Robina pseudoacacia

Madrone*Arbutus menziesii***Manzanita***Arctostaphylos spp.***Maple***Acer spp.***Monkey Flower***Mimulus guttatus***Oak***Quercus spp.***Persimmon***Diospyros spp.***Pine***Pinus spp.***Poison Ivy***Rhus radicans***Poison Oak***Rhus toxicodendron***Poplar, yellow***Liriodendron tulipifera***Prunus***Prunus spp.***Raspberry***Rubus spp.***Redbud, eastern***Cercis canadensis***Rose, multiflora***Rosa multiflora***Sage, black***Salvia mellifera*

Sagebrush, California
Artemisia californica

Salmonberry
Rubus spectabilis

Saltbush; Sea myrtle
Baccharis halimifolia

Sassafras
Sassafras albidum

Sourwood
Oxydendrum arboreum

Sumac
Rhus vernix

Sweetgum
Liquidambar styraciflua

Swordfern
Polystichum munitum

Tallowtree, Chinese
Sapium sebiferum

Tan Oak
Lithocarpus densiflorus

Thimbleberry
Rubus parviflorus

Tobacco, tree
Nicotiana glauca

Trumpetcreeper
Campsis radicans

Waxmyrtle, southern
Myrica cerifera

Willow

Salix spp.

HERBACEOUS WEEDS

Bahiagrass

Paspalum notatum

Balsamapple

Momordica charantia

Barnyardgrass

Echinochloa crus-galli

Bassia, fivehook

Bassia hyssopifolia

Bermudagrass

Cynodon dactylon

Bindweed, field

Convolvulus arvensis

Bluegrass, Kentucky

Poa pratensis

Brackenfern

Pteridium aquilinum

Brome

Brome spp.

Bromegrass, smooth

Bromus inermis

Broomsedge

Andropogon spp.

Buttercup

Ranunculus spp.

Cheat

Bromus secalinus

Chickweed, mouseear
Cerastium vulgatum

Clover, red
Trifolium pratense

Clover, white
Trifolium repens

Cocklebur
Xanthium strumarium

Crabgrass
Digitaria spp.

Dallasgrass
Paspalum dilatatum

Dock, curly
Rumex crispus

Dwarf dandelion
Krigia cespitosa

False flax, smallseed
Camelina microcarpa

Fescue
Festuca spp.

Fiddleneck
Amsinckia spp.

Flaxleaf fleabane
Conyza bonariensis

Fleabane
Erigeron spp.

Foxtail
Setaria spp.

Groundsel, common

Senecio vulgaris

Guineagrass

Panicum maximum

Horsenettle

Solanum carolinense

Horseweed/Marestail

Conyza canadensis

Johnsongrass

Sorghum halepense

Kikuyugrass

Pennisetum clandestinum

Knapweed

Centaurea repens

Kochia

Kochia scoparia

Lambsquarters, common

Chenopodium album

Lespedeza: common, sericea

Lespedeza striata

Lespedeza cuneata

Lettuce, prickly

Lactuca serriola

Morningglory

Ipomoea spp.

Muhly, wirestem

Muhlenbergia frondosa

Mullein, common

Verbascum thapsus

Mustard, blue

Chorispora tenella

Mustard, tansy
Descurainia pinnata

Mustard, tumble
Sisymbrium altissimum

Mustard, wild
Sinapis arvensis

Napiersgrass
Pennisetum purpureum

Nightshade, silverleaf
Solanum elaeagnifolium

Nutsedge: purple, yellow
Cyperus rotundus
Cyperus esculentus

Oats, wild
Avena fatua

Orchardgrass
Dactylis glomerata

Panicum
Panicum spp.

Pampasgrass
Cortaderia jubata

Pennycress, field
Thlaspi arvense

Pigweed, redroot
Amaranthus retroflexus

Pigweed, smooth
Amaranthus hybridus

Quackgrass
Agropyron repens

Ragweed, common
Ambrosia artemisiifolia

Ragweed, giant
Ambrosia trifida

Reed, giant
Arundo donax

Ryegrass, perennial
Lolium perenne

Saltcedar
Tamarix spp.

Sandbur, field
Cenchrus spp.

Shepherdspurse
Capsella bursa-pastoris

Signalgrass, broadleaf
Brachiaria platyphylla

Smartweed, Pennsylvania
Polygonum pennsylvanicum

Sowthistle, annual
Sonchus oleraceus

Spanishneedles
Bidens bipinnata

Spurry, umbrella
Holosteum umbellatum

Starthistle, yellow
Centaurea solstitialis

Stinkgrass
Eragrostis cilianensis

Thistle, Canada
Cirsium arvense

Thistle, Russian

Salsola kali

Vaseygrass

Paaspalum urvillei

Velvetgrass

Holcus spp.

Witchgrass

Panicum capillare

INJECTION AND CUT STUMP APPLICATIONS

Woody brush and trees may be controlled using injection or cut stump applications of this product.

INJECTION APPLICATIONS

Apply the equivalent of 1 ml of this product per each 2 inches of trunk diameter. This is best achieved by applying a 25 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In these species, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, avoid applications during peak sap flow in the Spring.

CUT STUMP TREATMENTS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. **Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting.** Delays in application may result in

reduced performance. For best results, avoid applications during peak sap flow in the Spring.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS most woody brush and tree species, some of which are listed below:

Alder

Alnus spp.

Coyotebrush

Baccharis consanguinea

Dogwood

Cornus spp.

Eucalyptus

Eucalyptus spp.

Hickory

Carya spp.

Madrone

Arbutus menziesii

Maple

Acer spp.

Oak

Quercus spp.

Poplar

Populus spp.

Saltcedar

Tamarisk spp.

Sweetgum

Liquidambar styraciflua

Sycamore

Platanus occidentalis

Tan Oak

Lithocarpus densiflorus

Willow

Salix spp.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions!!) when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER

SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

Product protected by
U.S. Patent No. 4,405,531.
Other patents pending.

©MONSANTO COMPANY 1997
ST. LOUIS, MISSOURI 63167 U.S.A.