

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra herbicide and this supplement.



NOT REVIEWED
in accordance with FR Notice 82-2,
on Draft Labeling Dated 6/29/99

Herbicide

EPA Reg. No. 524-475

Roundup Ultra and Roundup Ready are registered trademarks of Monsanto Company.

FOR USE AS A BANANACIDE IN BANANA PLANTINGS INFECTED WITH
THE BANANA BUNCHY TOP VIRUS

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

Follow all applicable directions, restrictions, and precautions on the product label.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS),

DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

For control of the banana bunchy top virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy

top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

USE INSTRUCTIONS:

This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1/25 fluid ounce (1 mL) of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed.

274

All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be destroyed.

PRECAUTIONS, RESTRICTIONS:


Do not apply more than ½ fluid ounce (15 mL) of this product's concentrate per mat (or unit). Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for general weed control.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Ultra herbicide before using this product. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©1999 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

21137X1-65

6/29/99



DIRECTIONS FOR USE

For control of the banana bunchy top virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

USE INSTRUCTIONS:

This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1/25 fluid ounce (1 mL) of this product's concentrate per 2 - 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed.

All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be destroyed.

PRECAUTIONS, RESTRICTIONS:

Do not apply more than 1/2 fluid ounce (15 mL) of this product's concentrate per mat (or unit). Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for general weed control.

- **Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.**

