

PLEASE NOTE


**This image contains more than one label
approved for this product on this date.**

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ULTRA®
Herbicide by Monsanto
EPA Reg. No. 524-475



USE ONLY FOR SEED PRODUCTION OF LETTUCE WITH THE ROUNDUP READY® GENE

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra and Roundup Ready are registered trademarks of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN LETTUCE WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT LETTUCE IN PRODUCTION FIELDS OF LETTUCE CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF LETTUCE WILL RESULT IF LETTUCE VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

2 7 2

USE RECOMMENDATIONS

This product will control non-glyphosate tolerant lettuce in seed production fields of lettuce containing the Roundup Ready gene. Apply up to 64 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 64 ounces per acre may be applied, if needed to control non-glyphosate tolerant lettuce plants.

DO NOT EXCEED A MAXIMUM RATE OF 4 QUARTS OF THIS PRODUCT PER ACRE PER SEASON.


Application timing -- This product can be applied to Roundup Ready lettuce from emergence to harvest.

Treated lettuce may not be used for food or feed. Do not feed or graze treated lettuce. Do not process treated lettuce for food or feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1999

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167



NEXT


LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ULTRA®
Herbicide by Monsanto
EPA Reg. No. 524-475



USE ONLY FOR SEED PRODUCTION OF ALFALFA WITH THE ROUNDUP READY® GENE

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra and Roundup Ready are registered trademarks of Monsanto Company.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, OTHER THAN ALFALFA WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT ALFALFA IN PRODUCTION FIELDS OF ALFALFA CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF ALFALFA WILL RESULT IF ALFALFA VARIETIES WHICH DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.



292

USE RECOMMENDATIONS

This product will control non-glyphosate tolerant alfalfa in seed production fields of alfalfa containing the Roundup Ready gene. Apply up to 64 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Subsequent applications of up to 64 ounces per acre each may be applied, if needed to control non-glyphosate tolerant alfalfa plants.

DO NOT EXCEED A MAXIMUM RATE OF 8 QUARTS OF THIS PRODUCT PER ACRE PER SEASON.


Application timing -- This product can be applied to Roundup Ready alfalfa from emergence to harvest.

Treated alfalfa or the resulting seed may not be used for food or feed. Do not feed or graze treated alfalfa. Do not process treated alfalfa or resulting seed for food or feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1999

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167



NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra herbicide and this supplement.

ROUNDUP ULTRA®

Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED

FEB 16 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

ROUNDUP ULTRA® Tank-Mixed with RESOURCE® for improved symptoms in pre-plant applications in Corn or Soybeans.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra is a registered trademark of Monsanto Company.

® Resource is a registered trademark of Valent USA Corporation

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION", "MIXING", "ANNUAL WEED RATE TABLE", and "PERENNIAL WEED RATE TABLE" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This tank-mixture will significantly enhance the speed of symptomology appearance on certain weeds when applied as a pre-plant treatment prior to corn or soybeans. This mixture may be applied prior to planting or emergence of labeled crops. This mixture will not provide residual control of weeds that are un-emerged at the time of treatment.



Combine Roundup Ultra (32 fluid ounces of product per acre) and Resource (2.08 fluid ounces per acre) in the spray tank in 10 - 20 gallons of water per acre. For best results, make applications to actively growing weeds. Ammonium Sulfate at 1-2% by weight or 8.5 to 17 pounds per 100 gallons of water may be added. The addition of 2,4-D or Banvel is not recommended.

SPECIFIC MIXING RECOMMENDATIONS

Option 1

1. Fill the spray tank one-half full with water and start the agitation.
2. Add Resource at 2.08 fluid oz /acre and mix thoroughly.
3. If desired add Ammonium Sulfate at 1-2% by weight or 8-17 lbs/100 gallons of spray.
4. Add Roundup Ultra at 32 fluid oz/acre and finish filling the spray tank with water.

Option 2


1. Fill the spray tank one-half full with water and start the agitation.
2. If desired add Ammonium Sulfate at 1-2% by weight or 8-17 lbs/100 gallons of spray.
3. Add Roundup Ultra at 32 fluid oz/acre.
4. Add Resource at 2.08 fluid oz/acre to the spray tank.

Read and follow the Resource label for additional application instructions and precautions.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 2000

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167



NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra herbicide and this supplement.

ROUNDUP ULTRA®
Herbicide by Monsanto

EPA Reg. No. 524-475

ACCEPTED

FEB 16 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-475

ROUNDUP ULTRA ® Tank-Mixed with MON 46598 for improved symptoms in pre-plant applications.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION", "MIXING", "ANNUAL WEED RATE TABLE", and "PERENNIAL WEED RATE TABLE" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This tank-mixture will significantly enhance the speed of symptomology appearance on certain weeds when applied as a pre-plant treatment prior to corn, soybeans or wheat. This mixture may be applied prior to planting or emergence of labeled crops. This mixture will not provide residual control of weeds that are un-emerged at the time of treatment.

Combine Roundup Ultra (32 fluid ounces of product per acre) and MON-46598 (0.144 ounces per acre) in the spray tank in 10 - 20 gallons of water per acre. For best results, make

applications to actively growing weeds. Ammonium Sulfate at 1-2% by weight or 8.5 to 17 pounds per 100 gallons of water may be added. The addition of 2,4-D or Banvel is not recommended.

SPECIFIC MIXING RECOMMENDATIONS

Option 1

1. Fill the spray tank one-half full with water and start the agitation.
2. Add MON-46598 at 0.144 oz /acre and mix thoroughly.
3. If desired add Ammonium Sulfate at 1-2% by weight or 8-17 lbs/100 gallons of spray.
4. Add Roundup Ultra at 32 oz/acre and finish filling the spray tank with water.

Option 2


1. Fill the spray tank one-half full with water and start the agitation.
2. If desired add Ammonium Sulfate at 1-2% by weight or 8-17 lbs/100 gallons of spray.
3. Add Roundup Ultra at 32 oz/acre.
4. Prepare a slurry of MON-46598 with water and add to the spray tank.

Read and follow the MON-46598 label for additional application instructions and precautions.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 2000

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167



NEXT

LABEL


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

ROUNDUP ULTRA®
Herbicide by Monsanto

EPA Reg. No. 524-475



FOR USE IN THE FOLLOWING MEDICINAL HERBS: Burdock, Dandelion, Echinacea, Feverfew, Goldseal, Stinging Nettle, St. Johnswort, Valerian, Blue Vervain

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup and Roundup Ultra are registered trademarks of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

USE RECOMMENDATIONS

For specific rates of application and instructions for control of various annual and perennial weeds, see the Annual and Perennial Weeds Rate Tables in the Roundup Ultra label booklet.

TYPES OF APPLICATIONS: Chemical fallow, preplant fallow beds, preplant,

preemergence, post directed hooded applications and postharvest.

Chemical fallow, Preplant fallow beds

USE INSTRUCTIONS: This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields.

Preplant, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting by direct seeding and prior to transplanting the crops listed above. Applications must be made prior to emergence of the crop.

PRECAUTIONS, RESTRICTIONS: Preplant applications must be made at least 3 days prior to transplanting. When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by a single 0.5 inch application of water either by natural rainfall or via a sprinkler system. EXTREME CARE MUST BE TAKEN TO ENSURE THAT THERE IS NO PART OF A SEED OR EMERGING SEEDLING CAPABLE OF BEING CONTACTED AND THAT THE SOIL IS TOTALLY COVERING THE SEED OR DEVELOPING SEEDLING AT APPLICATION. ENSURE THAT NO PART OF ANY SEED OR EMERGING SEEDLING IS CONTACTED BY HERBICIDE SOLUTION, SPRAY DRIFT OR MIST.

For preemergence applications prior to the emergence of established perennial crops do not apply within one week prior to emergence of any plant part.

Post directed hooded applications, Post-harvest

USE INSTRUCTIONS: Post directed hooded applications of this product may be made to mulched or unmulched row middles after crop establishment.

This product may be applied after final harvest to control weeds, suppress regrowth of annual crops or for renovation of biennial or perennial crop beds. Higher rates may be needed to control large weeds which were growing in the crop at the time of harvest.

PRECAUTIONS, RESTRICTIONS: Post directed applications may be made at a maximum of 1 quart per acre and a maximum of 1 application per growing season. Post directed and post-harvest applications must be made at least 14 days prior to planting the next crop. Do not harvest or feed treated vegetation.



EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF DESIRABLE PLANTS. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

Grower assumes all responsibility for crop losses from misapplication.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 2000

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

