

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

**ROUNDUP® ULTRA**  
Herbicide by Monsanto

EPA Reg. No. 524-475

---

### WIPER APPLICATIONS OF WEEDS IN PEPPERMINT AND SPEARMINT

---

Keep out of reach of children.

### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

---

® Roundup Ultra is a registered trademark of Monsanto Company.

---

### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS). DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

---

ACCEPTED

MAY - 1 2000

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 524-475

## DIRECTIONS FOR USE

### USE INSTRUCTIONS:

This product may be used in wiper applications in peppermint and spearmint. For more information on wiper applications, see the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for Roundup Ultra herbicide.

Applicators should be adjusted so that the wiper contact point is at least 2 inches above the crop. Weeds should be a minimum of 6 inches taller than the crop. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when height of weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. Better results may be obtained if 2 applications are made in opposite directions.

**PRECAUTIONS, RESTRICTIONS:** Contact of the herbicide solution with mint may result in damage or destruction. Allow at least 7 days between application and harvest. Sequential application may be made in the same area at 30-day intervals.

---

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

---

© Monsanto Company 2000

MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167

**NEXT**

**LABEL**

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

### ROUNDUP® ULTRA

Herbicide by Monsanto

EPA Reg. No. 524-475

---

### SPOT-TREATMENT OF WEEDS IN DRY DITCHES IN CRANBERRY PRODUCTION AREAS

---

Keep out of reach of children.

### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

---

® Roundup Ultra is a registered trademark of Monsanto Company.

---

### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

---

ACCEPTED

MAY - 1 2000

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 524-475

## DIRECTIONS FOR USE

### USE INSTRUCTIONS:

Directed sprays may be used to control weeds growing in the interior and perimeter ditch areas of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in the Roundup Ultra herbicide label booklet may be used. Drop water level to remove standing water in ditches prior to application. In hand-held sprayers, use 1-2% solution of Roundup Ultra herbicide on a volume-to-volume basis with water. Spray to wet vegetation, not to run-off.

### PRECAUTIONS, RESTRICTIONS:

For treatments after drawdown of water in dry ditches, allow 2 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not apply this material through the irrigation system. Do not make applications by air. Do not apply directly to water.

Contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Use nozzles that emit medium to large-sized droplets to minimize drift. The likelihood of crop injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Avoid drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.


---

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

---

© Monsanto Company 2000

MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167



**NEXT**

**LABEL**

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.

**ROUNDUP® ULTRA**  
Herbicide by Monsanto

EPA Reg. No. 524-475

---

### POST HARVEST SPOT-TREATMENT OF WEEDS IN CRANBERRY BOGS

---

Keep out of reach of children.

#### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

---

® Roundup Ultra is a registered trademark of Monsanto Company.

---

#### DIRECTIONS FOR USE

---


It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

---



### DIRECTIONS FOR USE

#### USE INSTRUCTIONS:

Application of this product may be made after the harvest of cranberries to control weeds growing within the field. Best results will be obtained if applications are made to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in the Roundup Ultra herbicide label booklet may be used. If using hand-held sprayers, use a 1/2 to 1% solution of Roundup Ultra herbicide on a volume-to-volume basis with water. Spray to wet vegetation, not to run-off. If using hand-held boom sprayers, apply 2 to 4 quarts of this product per acre.

#### PRECAUTIONS, RESTRICTIONS:

Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not apply this product through the irrigation system. Do not make applications by air. Do not apply directly to water.

Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed. Use nozzles that emit medium to large-sized droplets to minimize drift. The likelihood of crop injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Avoid drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

---

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

---

© Monsanto Company 2000  
MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167

