


PLEASE NOTE

This image contains more than one label approved for this product on this date.

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this



ROUNDUP® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

FOR PREHARVEST APPLICATIONS IN ALFALFA

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This product will control labeled annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa.

The treated crops and weeds can be harvested and fed to livestock. Allow a minimum of 36 hours between application and harvest.

Use up to 2 quarts of this product per acre. Applications may be made at any time of the year. Make only one application to an existing stand of alfalfa per year.

For control of quackgrass, apply in the spring, late summer, or fall when quackgrass is actively growing. Control in western locations may require deep tillage following application.

For control of alfalfa, deep tillage may be required in western locations if alfalfa is drought stressed.

Do not use for alfalfa grown for seed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1999

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

	» » » »
	» » » »
	» » » »
	» » » »
» » » » »	»
	» » » »
» » » » »	» » » »
» » » » »	» » » »
» » » » »	» » » »
» » » » »	»
	» » » »
	» » » »
	» » » »
	» » » »


NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP® ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra and this supplement.



ROUNDUP® ULTRA
Herbicide by Monsanto

EPA Reg. No. 524-475

FOR PREHARVEST APPLICATIONS IN BARLEY ONLY IN THE STATES OF IDAHO, MINNESOTA, MONTANA, NORTH DAKOTA, OREGON, SOUTH DAKOTA, AND WASHINGTON

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

® Roundup Ultra is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

RECOMMENDATIONS

This product may be used prior to harvest of barley to control annual and perennial weeds listed on the label booklet for Roundup Ultra herbicide.

INSTRUCTIONS

Apply up to 1 quart of this product per acre in 5 to 10 gallons of water per acre.

This product may be applied using either aerial or ground spray equipment. See the "mixing" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of the label for complete instructions.

PRECAUTIONS, RESTRICTIONS

Preharvest applications of this product are not recommended for malting barley.

Make applications when grain has 30% grain moisture or less and at least 7 days prior to harvest. Applications made at greater than 30% moisture can reduce barley quality and increase lodging.

Note: It is not recommended that barley grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup Ultra herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© Monsanto Company 1999

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167