


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ULTRA HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Ultra herbicide, container labels and this supplement.



Roundup Ultra

Herbicide

EPA Reg. No. 524-475

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
Reg. No. 524-475

Roundup Ultra is a trademark and Roundup Ready is a registered trademark of Monsanto Technology LLC.

FOR POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS) DESIRABLE PLANTS AND TREES, OTHER THAN CORN WITH THE ROUNDUP READY GENE, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Ultra herbicide for essential product performance information.

GENERAL INFORMATION

MONSANTO COMPANY RECOMMENDS USE OF THIS PRODUCT ONLY ON CORN HYBRIDS DESIGNATED AS CONTAINING THE ROUNDUP READY GENE.

- Applying this product to corn hybrids which are not designated as Roundup Ready will result in severe crop injury and yield loss.
- The Roundup Ready designation indicates that the corn contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready corn may be obtained from your seed supplier or Monsanto representative.

Application Instructions

For Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first, this product may be applied over-the-top broadcast or with drop nozzles. For corn

heights 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles adjusted to avoid spraying into the whorls of the corn plants. When corn height is 24 to 48 inches (free standing), for optimum spray coverage and weed control, drop nozzles are recommended.

Single in-crop applications of this product should not exceed 1.5 quarts per acre.

Maximum Yearly Rates Allowed

Preplant: Maximum amount of this product which can be applied prior to crop emergence is 5 quarts per acre.

In-crop: Maximum combined total of multiple in-crop applications from emergence through the 48 inch stage is 3 quarts per acre.

Cropping Season: Combined total per year for all applications may not exceed 8 quarts per acre.

When applied as directed, this product controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Applications should be made to actively growing weeds before they reach the maximum size listed in the "WEEDS CONTROLLED" section of the label booklet for Roundup Ultra herbicide. Refer to the "MIXING" section of the label booklet for proper use instructions.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet® or Micro-Tech® herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including fertilizers and micro-nutrients are not

recommended with this product since this may result in increased potential for crop injury.

Allow a minimum of 50 days between application of this product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 10 days between in-crop applications of this product. There are no rotational crop restrictions following applications of this product.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

For ground applications: Use the recommended rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the recommended rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 1.5 quarts per acre. See "WEEDS CONTROLLED" section on this label. AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Weed Control Recommendations

Apply 24 to 32 fluid ounces of this product per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for Roundup Ultra herbicide for rate recommendations for specific annual weeds. Roundup Ultra herbicide applied at up to 1.5 quarts per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem chuhly. For additional information on perennial weeds, see the "PERENNIAL WEED RATE TABLE" of the label booklet for Roundup Ultra herbicide.

Preemergence followed by Postemergence Weed Control Program

This product may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. An in-crop application of this product at the recommended rate will provide control of emerged weeds listed on the label. This product may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first. For corn heights 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants. When corn height is 24 to 48 inches (free standing), for optimum spray coverage and weed control drop nozzles are recommended.

Postemergence Only Weed Control Program

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied over-the-top broadcast or with drop nozzles

postemergence to Roundup Ready corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first. For corn height 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants. When corn height is 24 to 48 inches (free standing), for optimum spray coverage and weed control, drop nozzles are recommended.

This product may be applied in tank mixture with a labeled rate of Harness®, Harness Xtra, Harness Xtra 5.6L, Micro-Tech, Bullet, Permit® or atrazine. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines - the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

Tank Mix Partner	Maximum Height Of Corn For Application
Harness Harness Xtra Harness Xtra 5.6L	11 inches
Bullet* Micro-Tech*	5 inches
Permit	30 inches
Atrazine	12 inches

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

Bullet, Harness and Micro-Tech are registered trademarks of Monsanto Technology LLC

Permit is a trademark of, and used under license from, Nissan Chemical Industries, Ltd.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Ultra herbicide before using. For over-the-top uses on Roundup Ready crop varieties, crop safety and weed control performance are not warranted by Monsanto when this product is used in conjunction with "brown bag" or "bin run" seed saved from the previous year's productions and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2002 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167