

524-475

11/01/2007

1/2

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PRO® herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

**Roundup PRO®**

Herbicide by Monsanto

### FOR HOLLOW STEM INJECTION

**ACCEPTED**

NOV - 1 2007

EPA Reg. No. 524-475

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended for the pesticide  
registered under  
EPA Reg. No. 524-475

Keep out of reach of children.

**CAUTION!**

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup PRO is a registered trademark of Monsanto Technology LLC.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application.

**AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup PRO herbicide for essential product performance information.

This product may be applied through hand-held injection devices that deliver recommended amounts of this product into targeted hollow-stem plants growing in any non-crop site specified on this label. For control of the following hollow-stem plants, follow the use instructions below:

**Castorbean (*Ricinus communis*)**

Inject 5 mL per plant of this product into the lower portion of the main stem.

**Hemlock, Poison (*Conium maculatum*)**

Inject one leaf cane per plant 10 to 12 inches above root crown with 6 mL of a 5% v/v solution of this product.

**Hogweed, Giant (*Heracleum mantegazzianum*)**

Inject one leaf cane per plant 12 inches above root crown with 6 mL of a 5% v/v solution of this product.

**Horsetail, Field (*Equisetum arvense*)**

2/2

Inject one segment above the root crown with 0.6 mL per stem of this product. Use a small syringe that calibrates to this rate.

**Knotweed, Bohemian (*Polygonum bohemicum*)**

Inject 6 mL per stem of this product between second and third internode.

**Knotweed, Giant (*Polygonum sachalinense*)**

Inject 6 mL per stem of this product between second and third internode.

**Knotweed, Japanese (*Polygonum cuspidatum*)**

Inject 6 mL per stem of this product between second and third internode.

**Reed, Giant (*Arundo donax*)**

Inject 8 mL per stem of this product between the second and third internode.

**Thistle, Canada (*Cirsium arvense*)**

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.6 mL per stem of this product is injected into the stem.

**NOTE:** Based on the maximum annual use rate of glyphosate for these non-crop sites, the combined total for all treatments must not exceed 10.6 quarts of this product per acre. At 6 mL per stem, 10.6 quarts should treat approximately 1700 stems.

**Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup PRO herbicide before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.**

© [DATE] MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167