

PLEASE NOTE

This image contains more than one label approved for this product on this date.

524-500

2/22/2001

1/7

ACCEPTED

FEB 22 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-500



Herbicide by Monsanto

Outrider herbicide is a selective herbicide for control of annual and perennial grass and broadleaf weeds in noncrop areas.

Complete Directions For Use

Outrider is a trademark of Monsanto Company.

EPA Reg. No. 524-500

2000-1

Read the entire label before using this product.

Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

ACTIVE INGREDIENT:

| | |
|--------------------------|---------------|
| Sulfosulfuron..... | 75.0% |
| OTHER INGREDIENTS: | 25.0% |
| | <u>100.0%</u> |

This product is protected by U.S. Patent Nos. 5,017,212 and 5,534,482. No license granted under any non U.S. Patent(s).

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling

| FIRST AID | |
|---|---|
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call collect, day or night, (314) 694-4000 for emergency medical treatment information. | |

Environmental Hazards

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is highly toxic to non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal Laws.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL FREE, 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: Store under cool, dry conditions (below 120°F). Do not store under moist conditions.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

See individual container label for disposal information.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Ground Broadcast Equipment

Apply Outrider™ herbicide uniformly with properly calibrated ground equipment in 10 to 50 gallons of water per acre. Select spray volumes that ensure thorough and uniform weed coverage. Use equipment which is capable of continuous agitation. Choose nozzles which provide optimum spray distribution and uniform coverage at the appropriate spray pressure. Avoid streaking, skips, overlaps, and spray drift during applications.

Hand-Held and High-Volume Equipment

Use hand-held spray guns, backpacks, or other similar sprayers to apply this product. Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

Do not apply product through any type of irrigation system.

IMPORTANT: Do not allow this product to contact roots or foliage of desirable vegetation, areas where roots of desirable vegetation may extend, or areas where this product may be washed or moved into contact with roots of desirable vegetation.

Desirable plants may be injured if planted into treated areas.

Do not use on food or feed crops.

Equipment cleaning

Clean application equipment after using this product by thoroughly flushing with water.

Injection Systems

This product may be used in ground injection spray systems. It may be diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injections systems, unless specifically recommended.

ATTENTION

AVOID DRIFT. Care must be used when applying this product to prevent injury to desirable plants and crops. Do not allow the herbicide solution to mist, drift, or splash onto desirable vegetation or soil areas where sensitive crops will be planted since minute quantities of this product can cause severe damage or destruction to plants on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE. Drift control agent additives should be used for ground broadcast applications. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

MIXING INSTRUCTIONS

Thoroughly clean equipment prior to mixing spray solution.

Fill the spray tank to about three-fourths of the desired volume. Add the recommended amount of this product. Complete the filling process while maintaining agitation. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. Add nonionic surfactant near the end of the filling process where required.

For tank mixtures, add individual formulations to the spray tank in the following sequence: water, water soluble bags, dry flowables, emulsifiable concentrates, drift control additive, water soluble liquids followed by a nonionic surfactant. Tank mixtures of this product with Roundup Pro® herbicide do not require addition of surfactant.

GENERAL INFORMATION

Outrider herbicide is a preemergent and postemergent, systemic herbicide for control of many annual and perennial broadleaf and grass weeds on noncrop sites where unimproved bermudagrass, bahiagrass and tall fescue are present. It is formulated as a water dispersible granule (WDG).

NONCROP SITES

Outrider herbicide may be used for general weed control on noncrop sites including roadsides, utility rights-of-way, airports, fallow areas, ditch banks, dry ditches, dry canals, fencerows, industrial sites, lumberyards, manufacturing sites, petroleum tank farms and pumping installations, railroads, storage areas, utility substations, warehouse areas, and similar industrial and noncrop sites.

BERMUDAGRASS AND BAHIAGRASS

Outrider herbicide may be used to control or partially control many annual and perennial weeds for effective release of bermudagrass or bahiagrass on roadsides and other labeled noncrop sites.

Application Information

Ground Broadcast

Apply Outrider herbicide at 0.75 to 2.0 ounces per acre. Do not exceed 2.0 ounces of this product per acre per year.

Use the higher recommended rates of this product for control of large established weeds or when weed growth is heavy or dense. Best results are obtained when weeds are in the early stage of growth and are not disturbed by mowing or other factors for 12 days prior to, or 12 days after application.

Addition of a nonionic surfactant at 0.5 percent by volume (2 quarts per 100 gallons of spray solution) is required for postemergence applications unless tank mixed with Roundup Pro herbicide. Use only nonionic surfactants which contain at least 90 percent active ingredient. When this product is applied in a tank mixture with Roundup Pro herbicide, addition of a surfactant is not necessary.

Hand-Heid and High-Volume Equipment

Add 1.0 ounce of Outrider herbicide per 100 gallons of water. Add 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Use only nonionic surfactants containing at least 90 percent active ingredient.

4/7

Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label directions, precautionary statements and all information on the labels of all products used.

Any recommended rate of this product may be tank mixed with the following products. Refer to these product labels for approved noncrop sites and application rates. Always apply tank mixtures according to the most restrictive label of the products used. Products recommended as tank mixes with Outrider herbicide for bermudagrass or bahiagrass release are restricted to those specified in the "RELEASE OF BERMUDAGRASS OR BAHIAGRASS" section.

- | | |
|------------|--------------|
| BANVEL™ | PLATEAU™ |
| CAMPAIGN® | ROUNDUP PRO® |
| DIURON | TELAR™ |
| ESCORT™ | TRANSLINE™ |
| GARLON™ 3A | VANQUISH™ |
| KARMEX™ DF | 2,4-D |
| MSMA | |
| OUST™ | |

NOTE: Tank mixtures with broadleaf herbicides formulated as amines (including 2,4-D and others) may decrease the effectiveness of Outrider herbicide.

RELEASE OF BERMUDAGRASS OR BAHIAGRASS

Dormant Applications

This product may be used to control or partially control many winter annual weeds for effective release of dormant bermudagrass or bahiagrass when treated after grass is dormant and prior to spring green-up.

Apply any labeled rate of this product alone or in a tank mixture with 8 to 64 fluid ounces of Roundup Pro herbicide or with 16 to 64 fluid ounces of Campaign herbicide per acre. In dormant bermudagrass only, up to 1 ounce per acre of Escort may be added to Outrider herbicide or tank mixtures of Outrider herbicide and Roundup Pro herbicide to increase the spectrum of broadleaf weeds controlled. Addition of Escort may delay green-up of bermudagrass. Tank mixtures of this product with Escort in highly maintained turfgrass areas will result in unacceptable turf injury.

In the state of Texas, applications of Outrider herbicide applied before September 30 will not delay green-up of bermudagrass the following spring, however some temporary discoloration of desirable spring germinating wildflowers may occur.

Actively Growing Bermudagrass Release

This product may be used to control or partially control johnsongrass and other weeds for effective release of actively growing bermudagrass. Use only on well-established bermudagrass. Apply any labeled rate of this product alone or in a tank mixture with 8 to 32 fluid ounces of Roundup Pro herbicide per acre. Use the higher rates of each product to control perennial weeds or annual weeds greater than 6 inches in height.

Up to 0.5 ounce per acre of Oust, up to 1 ounce per acre of Escort, or up to 0.5 ounce per acre of Telar may be added to Outrider herbicide or tank mixtures of Outrider herbicide and Roundup Pro herbicide. Tank mixtures of this product with Oust, Escort or Telar in highly maintained turfgrass areas will result in unacceptable turf injury.

Actively Growing Bahiagrass Release

This product may be used to control or partially control johnsongrass and other weeds for effective release of actively growing bahiagrass. Use only on well established bahiagrass. Apply any labeled rate of this product alone or in a tank mixture with 4 to 12 ounces of Roundup Pro herbicide per acre. Use the higher rates of each product to control perennial weeds or annual weeds greater than 6 inches in height.

IMPORTANT: Established bermudagrass and bahiagrass are tolerant to Outrider herbicide applications. Tank mixtures of other products with Outrider herbicide may increase grass injury.

Where bermudagrass or bahiagrass are desirable ground cover use tank mixtures only when some temporary injury or discoloration can be tolerated.

TALL FESCUE

Outrider herbicide may be used to control or partially control johnsongrass and other weeds in tall fescue on roadsides and other labeled noncrop sites.

Application Information

Ground Broadcast

Apply Outrider herbicide at 0.75 to 1.0 ounce per acre. Do not exceed 1.0 ounce of this product per acre per year.

Use the higher recommended rate for control of large established weeds or when weed growth is heavy or dense. Best results are obtained when weeds are actively growing and are not disturbed by mowing or other factors for 14 days prior to, or 14 days after application.

Addition of a nonionic surfactant at 0.5 percent by volume (2 quarts per 100 gallons of spray solution) is required. Use only nonionic surfactants containing at least 90 percent active ingredient.

Hand-Held and High-Volume Equipment

Add 1.0 ounce of Outrider herbicide per 100 gallons of water. Add 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Use only nonionic surfactants containing at least 90 percent active ingredient.

Timing

Applications should be made to actively growing johnsongrass in early boot to full seedhead stage. Symptom development will be more rapid in less mature johnsongrass.

Use this product only in well-established tall fescue. Use of labeled rates can result in transient growth reduction and discoloration of tall fescue but these cause no long-term detrimental effects.

WEEDS CONTROLLED

Barley, volunteer

Hordeum vulgare

Bedstraw, catchweed

Galium aparine

Bluegrass, roughstalk

Poa trivialis

Bluegrass, bulbous

Poa bulbosa

Brome, downy

Bromus tectorum

Brome, rigput

Bromus rigidus

Buttercup

Ranunculus arvensis

Chamomile, mayweed

Anthemus cotula

Cheat

Bromus secalinus

Chess, hairy


Bromus commutatus

Chickweed, common

Stellaria media

Horseweed

Conyza canadiensis



6/7

- Fiddleneck, tarweed**
Amsinckia lycopsoides
- Flixweed**
Descurainia sophia
- Johnsongrass**
Sorghum halepense
- Ladysthumb**
Polygonum persicaria
- Mustard, tumble**
Sisymbrium altissimum
- Mustard, wild**
Sinapis arvensis
- Pennycress, field**
Thlaspi arvense
- Purple Nutsedge**
Cyperus rotundus
- Quackgrass**
Elytrigia repens
- Ragweed, common**
Ambrosia artemisiifolia
- Shepherd's-purse**
Capsella bursa-pastoris
- Sunflower, common**
Helianthus annuus
- Tansymustard, pinnate**
Descurainia pinnata
- Vetch, sand**
Vicia acutifolia
- Yellow Nutsedge**
Cyperus esculentus

LIMIT OF WARRANTY AND LIABILITY

Monsanto Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label pamphlet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE

7/7

THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.


Roundup Pro and Campaign are registered trademarks of Monsanto Company.
Escort, Karmex, Oust, and Telar are trademarks of E.I. duPont de Nemours and Company.
Garlon and Transline are trademarks of Dow AgroSciences.
Banvel and Plateau are trademarks of BASF Corporation.
Vanquish is a trademark of Novartis Crop Protection.

EPA Reg. No. 524-500

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

©2000 MONSANTO COMPANY
ST. LOUIS, MISSOURI, 63167 U.S.A.

Formulated in the United States Using Active Ingredient Made in Japan.



NEXT

LABEL

524-500

2/22/2001
SUPPLEMENTAL LABELING

1/3

READ THE ENTIRE LABEL FOR OUTRIDER™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Outrider herbicide and this supplement.

ACCEPTED
FEB 22 2001
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-500



1000000000
1000000000
1000000000
1000000000
1000000000
1000000000
1000000000
1000000000
1000000000
1000000000

EPA Reg. No. 524-500

Outrider is a trademark of Monsanto Company.

FOR USE FOR SELECTIVE HERBACEOUS WEED CONTROL IN FORESTRY CONIFER RELEASE.

Keep out of reach of children

CAUTION!

In case of emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

Note: Applications of this product should not contact leaves of desirable plants since foliar injury, discoloration, or death may result

Environmental Hazards

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This pesticide is highly toxic to non-target plants. Do not apply to non-target areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to plants in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal Laws.

Application Equipment and Instructions

AVOID DRIFT—DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE >5 MPH OR GUSTY, OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Coarse sprays are less likely to drift; therefore, avoid combinations of pressure and nozzle types that will result in splatter or fine particles (mist) which are likely to drift.

Aerial Application

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Aerial application of this product is by helicopter only. Apply the recommended rate of this product in 5 to 30 gallons of water per acre. Use the higher recommended spray volumes where weeds are dense or form multiple canopy layers. When used according to label directions this product will give control or partial control of herbaceous weeds such as

ragweed and sedges and other weeds listed in the "WEEDS CONTROLLED" section of the Outrider label.

Aerial Spray Drift Management

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The importance of spray droplet size:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but may not prevent drift if applications are made improperly or under unfavorable environmental conditions (see the following Wind, Temperature and Humidity, and Temperature Inversion sections of this advisory).

Controlling initial droplet size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher flow rates produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation** - Orienting nozzles so the spray stream is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Controlling placement of spray droplets:

- **Boom length** - For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height** - Applications should not be greater than 10 feet above the top of the tallest plants unless a greater height is required for aircraft safety. Greater application heights result in greater droplet size reduction through evaporation and greater movement in air currents. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
- **Application speed** - Slower aircraft speeds within a safe range will produce less air turbulence and fewer small droplets.
- **Swath adjustment** - When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (wind speed, droplet size, etc.).

Key environmental factors:

- **Wind** - Drift potential is lowest between wind speeds of 2-10 mph. However, many factors including droplet size and equipment type determine drift potential at any given speed. Application should be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Applicators should be familiar with local wind patterns and how they affect spray drift.
- **Temperature and humidity** - When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.
- **Temperature inversions** - Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable air currents that are common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke detector. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive areas:

Pesticides should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water.

Drift control additives may be used. When a drift control additive is used, read and carefully observe all the cautionary statements and all other information appearing on the additive label.

Ensure uniform application—To avoid streaked, uneven, or overlapped application, use appropriate marking devices.

FORESTRY HERBACEOUS CONIFER RELEASE

This product is recommended for the control or partial control of herbaceous weeds in forestry conifer release as a Spring or early Summer application after planting. Best results with this product are obtained where Accord[®] SP or labeled tank mixes with Accord SP have been used for site preparation prior to planting.

This product is recommended for use in conifer release after planting loblolly, slash or longleaf pine and in fallow silvicultural nursery sites for these species.

APPLICATION RATES AND TIMING

Ground Broadcast

Apply Outrider herbicide at 0.75 to 2 ounces per acre. Do not exceed 2 ounces of this product per acre per year.

Use the higher recommended rates of this product or recommended tank mixes for control of large established weeds or when weed growth is heavy or dense. Best results are obtained when weeds are in the early stage of growth.

Addition of a nonionic surfactant at 0.5 percent by volume (2 quarts per 100 gallons of spray solution) is required for postemergence applications unless tank mixed with another surfactant containing product. Use only nonionic surfactants which contain at least 90 percent active ingredient.

Hand-Held and High-Volume Equipment

Add 1 to 2 ounces of Outrider herbicide per 100 gallons of water. Add 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Use only nonionic surfactants containing at least 90 percent active ingredient.

Aerial Equipment

Apply Outrider herbicide at 0.75 to 2 ounces per acre. Do not exceed 2 ounces of this product per acre per year.

Use the higher recommended rates of this product or recommended tank mixes for control of large established weeds or when weed growth is heavy or dense. Best results are obtained when weeds are in the early stage of growth.

Addition of a nonionic surfactant at 0.5 percent by volume (2 quarts per 100 gallons of spray solution) is required for postemergence applications unless tank mixed with another surfactant containing product. Use only nonionic surfactants which contain at least 90 percent active ingredient.

TANK MIXTURES FOR CONIFER RELEASE

Tank mixtures of this product may be used to increase the spectrum of herbaceous vegetation controlled. When tank

3/3
mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry conifer release, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any recommended rate of this product may be used in a tank mix with the following products for forestry use.

| PRODUCT | BROADCAST RATE |
|---|----------------------------|
| Arsenal™ Applicators Concentrate | 4 fl oz /a* |
| Arsenal Applicators Concentrate + Oust™ | 4 fl oz /a + 1 oz/a* |
| Oust | 1-2 oz/a |
| Oust + Velpar | 1-2 oz /a + 0.375-0.5 lb/a |
| Oustar™ | 8-12 oz/a |
| Velpar | 0.375-0.75 lb/a |

*Use of surfactant not recommended for slash and longleaf pine.

Any of these mixtures can be used as a broadcast spray or a banded application around trees to reduce potential for soil erosion.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement in the label booklet for Outrider herbicide before using. These terms apply to this SUPPLEMENTAL LABELING, and if these terms are not acceptable return the product unopened at once.

©2001 Monsanto Company
St. Louis, Missouri 63167

1/1/2001