

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR OUTRIDER® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING


When using Outrider herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

OUTRIDER®

Herbicide by Monsanto

EPA Reg. No. 524-500

Outrider is a registered trademark of Monsanto Technology LLC.



Weed Control in Established Stands of Non-Crop Trees In Conservation and Wildlife Areas

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See the "GENERAL INFORMATION" and "MIXING" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of the Complete Directions For Use for Outrider herbicide for essential product information.

WEED CONTROL IN ESTABLISHED STANDS OF NON-CROP TREES

Outrider can be applied as a broadcast application around or over the top of select hardwood and conifer tree species in conservation and wildlife areas to control johnsongrass, tall fescue, purple and yellow nutsedge, and other weed species listed on the Outrider label. This product generally has been shown to be safe for use on the following tree species:

- American Plum
- Cottonwood
- Burr Oak
- Swamp Oak
- Sycamore
- Bald Cypress
- Green Ash
- Pin Oak
- Pecan
- Walnut

Treated trees must be growing in areas where commercial fruit or nut harvest will not occur. Make over-the-top applications to non-bearing trees only. Treat over the top of transplanted trees after they are well established. Temporary yellowing and growth reduction may occur in some species. Do not apply by air.

Apply up to 1.33 ounces of this product per acre with a non-ionic surfactant at 0.25 to 0.5 percent by volume (1 to 2 quarts per 100 gallons of spray solution). Sequential applications of this product can be made at a minimum of 21 days between applications, up to a maximum use rate of 2.66 ounces per acre per year.

Use only nonionic surfactants that contain at least 90 percent active ingredient. Do not use nonionic surfactants or other additives that alter the pH of the spray solution below pH 5.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label pamphlet for Outrider herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.