

524-500

1-24-2007

1/2

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR MAVERICK® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Maverick herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on this supplemental labeling and on the label booklet provided with the pesticide container. This supplemental labeling must be in the possession of the user at the time of pesticide application.


MAVERICK® HERBICIDE

EPA Reg. No. 524-500

Maverick is a registered trademark of Monsanto Technology LLC.

FOR USE IN WINTER WHEAT

Keep out of reach of children.
CAUTION!



In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "DIRECTIONS FOR USE" and "MIXING INSTRUCTIONS" sections of the label booklet for Maverick herbicide for essential product performance information.

WINTER WHEAT

Maverick herbicide can be applied in winter wheat either as a single preemergence application, a single post-emergence application or as a split postemergence application to control the weeds listed in the "WEEDS CONTROLLED" section of the product label. Best weed control is obtained when soil moisture is adequate to support vigorous wheat and weed growth.

Choose one of the following application scenarios.

Preemergence in Winter Wheat

Apply Maverick herbicide preemergence to winter wheat at 2/3 ounce of product per acre in a single application. Preemergence application of Maverick herbicide should be applied after drilling wheat but before wheat or weed emergence. Do not apply preemergence if dry soil conditions will cause delayed wheat and/or weed emergence. Preemergence applications under dry conditions make the product vulnerable to wind erosion until fall moisture is received. Under these conditions, wait until crop and weeds have emerged and are showing good vigor and follow the postemergence application instructions on this label.

Preemergence applications are not recommended for no-till systems or when high crop residue levels (plant material) are present on the soil surface.

Postemergence in Winter Wheat—Single Application

Apply Maverick herbicide at 2/3 ounce of product per acre in a single application when the target weeds listed in the "WEEDS CONTROLLED" section of the product label are actively growing. Use 0.5 percent by volume nonionic surfactant concentration (2 quarts per 100 gallons of spray solution) for all postemergence applications.

In the states of KS, OK, TX and MT, the single post-emergence application should be made after the wheat is at the 2-leaf stage but prior to the jointing stage (Feekes' Scale 6). In all other states, the single postemergence application can be made after wheat emergence but prior to the jointing stage (Feekes' Scale 6).

Brome (Cheat, Downy Brome, Japanese Brome)

For best control of brome species, apply Maverick herbicide as a single postemergence fall application of 2/3 ounce of product per acre when brome is in the 2- to 3-leaf stage of growth. Best performance with fall applications of Maverick herbicide will occur with good soil moisture and/or rainfall after application.

For spring postemergence suppression of brome species, apply a single application of 2/3 ounce of this product per acre when the brome has recovered from cold weather (majority of foliage is green and not red or purple) and is actively growing. For best results, make applications when brome is less than the 5-tiller stage of growth.

Mustards and Other Winter Annual Broadleaf Weeds

For fall postemergence control of mustards and other winter annual broadleaf weeds, apply 2/3 ounce of this product per acre in a single application. For best control, make application when weeds are less than 2 inches in diameter.

For spring postemergence control of winter annual broadleaf weeds, apply 2/3 ounce of this product per acre in a single application. For best control, make applications when weeds are less than 6 inches in diameter.

Postemergence in Winter Wheat—Split Application

[Optional Statement: For use only in the State of Montana]

As an alternative to a single postemergence application, Maverick herbicide may be applied to winter wheat in a split application. Start with an initial fall application of 3/8 ounce of product per acre after winter wheat and target weeds have emerged (after the 2-leaf stage), followed by a second application of 3/8 ounce of Maverick herbicide per acre in the spring, no sooner than two weeks following the initial application but prior to boot stage (Feekes' scale 9). Use 0.5 percent by volume nonionic surfactant concentration (2 quarts per 100 gallons of spray solution) for all postemergence applications.

FOR SPLIT APPLICATION ONLY, DO NOT EXCEED 3/4 OUNCE OF PRODUCT PER ACRE PER CROPPING SEASON.

Tank Mixtures for Winter Wheat

Before mixing in the spray tank, it is recommended that compatibility be tested by mixing all components in a small container in proportionate quantities. Add individual formulations to the spray tank in the following sequence: water dispersible granules (this product), water soluble bags, dry flowables, emulsifiable concentrates, drift control additives, water soluble liquids, nonionic surfactants.

Refer to the specific product labels and observe all precautions, mixing and application instructions for all products used in the tank mixture.

Insecticides: Maverick herbicide may be tank-mixed or used sequentially with insecticides, except for Malathion. DO NOT use Maverick herbicide plus Malathion, as crop injury may result.

Tank mixtures of Maverick herbicide with insecticides are not recommended when the wheat crop has significant insect damage, is under drought stress, or when growth is influenced by other environmental stress, such as nutrient deficiency, soil pH, or disease.

Do not apply Maverick herbicide within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment, as crop injury may result.

Herbicides: For control of additional broadleaf weeds, Maverick herbicide may be applied as a spring postemergence application in a tank mixture with the following herbicides.

- 2,4-D amine^{1,2,3}
- 2,4-D LV ester²
- Bronate™ (bromoxynil + MCPA)
- Buctril™ (bromoxynil)
- Buctril 4EC
- MCPA amine^{1,2,3}
- MCPA LV ester²
- Puma™ (fenoxaprop)³
- Sencor™ 4 (metribuzin)^{3,4}
- Senor™ DF (metribuzin)^{3,4}

¹ Tank mixtures with this herbicide may result in reduced control of brome species.

² Tank mixtures with this product may be made provided the specific product being used is registered for postemergence application to wheat..

³ Not recommended for use with split application rate of 3/8 ounce of Maverick herbicide.

⁴ Different formulations of the active ingredient may be used, provided the specific product being used is registered for postemergence application on wheat.

Tank mixtures with herbicides formulated as amines may decrease the effectiveness of Maverick herbicide.

Refer to the tank mix herbicide label for application rate and restrictions related to soil texture, soil organic matter, and wheat growth stage.

Always add nonionic surfactant to a concentration of 0.5 percent by volume.

Tank mixtures with metribuzin may be applied only in the spring.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label pamphlet for Maverick herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©[year] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

[print plate number]

[date]