# SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP SOLUGRAN™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup SoluGran herbicide and this supplement.

# Roundup SoluGran<sup>TM</sup> Herbicide by Monsanto

EPA Reg. No. 524-502

### FOR AERIAL APPLICATION IN CALIFORNIA ONLY.

Keep out of reach of children.

# **CAUTION!**

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Roundup SoluGran is a trademark of Monsanto Co.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

See the "GENERAL INFORMA-TION" and "MIXING" sections of the label booklet for essential product information.

See the "CROPPING SYSTEMS" section of the label booklet for specific recommendations on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION, SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

When mixing this product mix 4 or more quarts of a non-ionic surfactant per 100 gallons of spray solution. Use a nonionic surfactant labeled for use herbicides. The surfactant must contain 80 percent ОГ more active ingredient.

Aerial applications of this product are allowed in the following situations:

- 1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
- 2. In cotton, prior to harvest. Refer to the Roundup SoluGran herbicide booklet for specific preharvest application instructions.

Do not plant subsequent crops other than those listed in the label booklet for 30 days following application.

When applied as recommended under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

DO NOT EXCEED A MAXIMUM RATE OF 28 OUNCES BY WEIGHT PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE SYSTEMS.

DO NOT EXCEED A MAXIMUM RATE OF 14 OUNCES BY WEIGHT PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN COTTON, PRIOR TO HARVEST.

# **Aerial Equipment**

Use the recommended rates of this product in 3 to 15 gallons of water per acre.

Do not apply directly to water, to areas where surface water is present or to intértidal areas below the mean high water mark. 'Do not contaminate water when disposing of equipment washwaters.

AVOID DRIFT-DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CON-DITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CON-TACTED TO WHICH TREAT-MENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION. APPROPRIATE **BUFFER** ZONES MUST BE MAIN-TAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

- Do not apply within 100 feet of all desirable vegetation or crop(s).
- If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
- Winds blowing from 5 to 10 miles per - hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
- 4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the air-stream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application—To avoid streaking, uneven, or over-lapped application, use appropriate marking devices.

Thoroughly wash aircraft. especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED** EXPOSURE OF THIS PROD-UCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND **POSSIBLE** OF THE FAILURE PART. LANDING GEAR IS MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 prevent corrosion.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup SoluGran herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1999

MONSANTO COMPANY ST. LOUIS, MO 63167 USA

ACCEPTED

AUG 1 1 1899

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 224 502