SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP ULTRADRY™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booking to the label bookin Roundup UltraDry herbicide and this supplement.

MAY - 1 2000

Roundup UltraDry Herbicide

EPA Reg. No. 524-504

Under the Federal Insecticide, Fungicide, and Rodenticide Act. as amended, for the pesticide registered under BPA Reg. No 314-504

ADDITION OF SURFACTANT FOR APPLICATIONS WITH ROUNDUP ULTRADRY.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Roundup® and Roundup Ready®are registered trademarks and Roundup UltraDry is a trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

See the "GENERAL INFORMA-TION" and "MIXING" sections of the label booklet for essential product information.

)

See the "CROPS" section of the label booklet for specific recommendations on the use of this product.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THE SPRAY WITH FOLIAGE. GREEN STEMS, OR FRUIT OF DESIRABLE CROPS (EXCEPT AS SPECIFIED **INDIVIDUAL FOR** ROUNDUP READY CROPS). PLANTS, TREES OR OTHER DÊŞIRABLE VEGETATION. SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Although generally not required, surfactant may be added to spray solutions at water carrier volumes above gallons per acre application rates below ounces of this product per асге.

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, a rate of 0.25 to 0.5 percent surfactant concentration (1 to 2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient is recommended. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

When applied as recommended under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

DO NOT add buffering agents or pH adjusting agents to the spray solution when Roundup UltraDry is the only pesticide used. DO NOT ADD SURFACTANT TO PRODUCT FOR APPLICATIONS OVER-THE-TOP OF ROUNDUP READY® **CROPS** AND PREHARVEST TO COTTON.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label for Roundup UltraDry herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product uncrened at once.

©MONSANTO COMPANY 2000

MONSANTO COMPANY ST. LOUIS, MISSOURI 63167