

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP PRODRY™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup ProDry herbicide and this supplement.

Roundup ProDry Herbicide

FOR USE FOR SELECTIVE WEED CONTROL ON ROUNDUP PRODRY HERBICIDE TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

EPA Reg. No. 524-505

Roundup ProDry is a trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

NOTE: Applications of this product should not contact leaves of desirable plants since foliar injury, discoloration or death may result.

ROUNDUP PRODRY HERBICIDE-TOLERANT TALL FESCUE SELECTIONS FOR SEED PRODUCTION

Use this product on Roundup ProDry herbicide tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 2.3 to 9.2 ounces per acre as a post-emergence spray on Roundup ProDry herbicide tolerant tall fescue selections. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd'spurse, sowthistle, wild oat, dandelion, quack-grass, and Canada thistle. See the Roundup ProDry herbicide label for a complete list of weeds controlled or suppressed.

NOTE: The recommended rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with Roundup ProDry herbicide tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

TIMING OF APPLICATIONS

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy and in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup ProDry herbicide before using this product. Those terms apply to this supplemental unopened labeling and, if those terms are not acceptable, return the product at once.

Pure Gold and Aurora Gold are trademarks of Turf-Seed, Inc.

©2000 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63137

03/14/00

