

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

524-512

5/20/2004

1/2

# SUPPLEMENTAL LABELING

For distribution and use only in the States of Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, and Washington

READ THE ENTIRE LABEL FOR Roundup UltraMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup UltraMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

## Roundup UltraMAX HERBICIDE LOGO

Roundup UltraMAX is a trademark of Monsanto Technology LLC

**ACCEPTED**

MAY 20 2004

EPA Reg. No. 524-512

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticides registered under EPA Reg. No. 524-512

## PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY PEAS, LENTILS, AND CHICKPEAS.

Keep out of reach of children.

### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup UltraMAX herbicide for essential product performance information.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup UltraMAX herbicide for essential product performance information.

### DIRECTIONS FOR USE

See the "APPLICATION EQUIPMENT AND TECHNIQUES" Section of the Roundup UltraMAX herbicide label booklet for more information.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

**BROADCAST SPRAY**

**USE INSTRUCTIONS:** This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 21 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup UltraMAX herbicide label.
- Preharvest application is not recommended for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

**SPOT TREATMENTS**

**USE INSTRUCTIONS:** This product may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in peas, lentils, and chickpeas. Apply up to 21 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine spot treatments with a preharvest broadcast spray on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup UltraMAX herbicide label.
- Do not feed treated vines and hay from these crops to livestock.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

**Read the "Limit of Warranty and Liability" in the label booklet for Roundup UltraMAX herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.**

©2002 MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167

**NEXT**

**LABEL**

524-512

5/20/2004

1/2


# SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR Roundup UltraMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup UltraMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

## Roundup UltraMAX HERBICIDE LOGO

Roundup UltraMAX is a trademark  
of Monsanto Technology LLC



## PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY BEANS

Keep out of reach of children.

### CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup UltraMAX herbicide for essential product performance information.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup UltraMAX herbicide for essential product performance information.

### DIRECTIONS FOR USE

See the "APPLICATION EQUIPMENT AND TECHNIQUES" Section of the Roundup UltraMAX herbicide label booklet for more information.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

### BROADCAST SPRAY

**USE INSTRUCTIONS:** This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 26 fluid ounces in 3 to 20 gallons of water

per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup UltraMAX herbicide label.
- Preharvest application is not recommended for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat cowpeas, since this crop is considered to be grown as livestock feed.

**SPOT TREATMENTS**

**USE INSTRUCTIONS:** This product may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 26 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine spot treatments with a preharvest broadcast spray on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup UltraMAX herbicide label.
- Do not feed treated vines and hay from these crops to livestock.
- Do not treat cowpeas, since this crop is considered to be grown as livestock feed.

**Read the "Limit of Warranty and Liability" in the label booklet for Roundup UltraMAX herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.**

©2002 MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167