

524-517

6-19-2008

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUN 19 2008

Ms. Dawn Fee-White
Monsanto Company
1300 I (Eye) Street, Suite 450 East
Washington, DC 20005

Dear Ms Fee-White:

Subject: Ranger Pro Herbicide (Update Use on Citrus in Fl)
EPA Registration No. 524-517
Application Dated March 20, 2008

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. A stamped copy of labeling is enclosed for your records.

Supplemental labeling must be incorporated into master labeling and copies of master labeling submitted to the Agency for our files at your next printing or within two (2) years from the date of acceptance of supplemental labeling, whichever comes first. The Agency will consider a convincing argument as to why a supplemental label should not appear on a master label on a case by case basis.

Sincerely,

James A. Tompkins for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR RANGER PRO® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Ranger PRO herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

ACCEPTED

JUN 19 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 524-517

RANGER PRO®

Herbicide

FOR USE IN CITRUS CROPS IN FLORIDA

EPA Reg. No. 524-517

Keep out of reach of children

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

Ranger PRO is a trademark of Monsanto Technology LLC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "PRECAUTIONARY STATEMENTS", "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Ranger PRO herbicide for essential product information.

MAXIMUM APPLICATION RATE: For applications in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate containing products do not exceed stated maximum use rate.

LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor

TYPES OF APPLICATION: Preplant (Site Preparation) Broadcast Spray, General Weed Control, Middles (between rows), Strips (within rows), Shielded Sprayer, Hooded Sprayer, Wiper Applicator, Directed Spray, Spot Treatment

USE INSTRUCTIONS: Applications may be made with boom equipment, CDA equipment, shielded sprayer, hand-held and high-volume wand, lance, orchard gun or with wiper applicator equipment, except as directed. For general weed control apply this product according to the rates listed in the "ANNUAL WEEDS" and "PERENNIAL WEEDS" sections of the Complete Directions for Use booklet for Ranger PRO herbicide. Use the higher rate in the recommended range when weeds are stressed, growing in

Refer to the tank mixture product label for restrictions and precautions; use according to the most restrictive precautionary statements for each product in the tank mixture.

PRECAUTIONS, RESTRICTIONS: Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used where there is the potential for crop contact, and then only where there is sufficient clearance. See additional use instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the directions for use booklet for Ranger PRO.

Allow a minimum of 1 day between last application and harvest in citrus crops. Allow a minimum of 3 days between application and transplanting trees.

For citron groves, apply as a directed spray only.

Read the "Limit of Warranty and Liability" in the label booklet for Ranger PRO herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© 2008 MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI 63167