

524-517

1/25/2010

10f44



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

1-25-10

Dawn Fee-White
Monsanto Company
1300 I (Eye) Street, NW
Suite 450 East
Washington, DC 20005

Subject: EPA Reg. No. 524-517 / Ranger Pro Herbicide Label Amendment

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

A handwritten signature in black ink that appears to read "Erik Kraft".

James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

21225H4-1/53

Ranger PRO[®] Herbicide

Ranger PRO is a complete broad-spectrum postemergence professional herbicide for non-crop, industrial, Turf and ornamental weed control.

Complete Directions for Use

~~The complete broad-spectrum postemergence professional herbicide for industrial, turf and ornamental weed control.~~

EPA Reg. No. 524-517

2009-1

**AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS,
EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS,
DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY
OR DESTRUCTION IS LIKELY TO RESULT.**

| |
|--|
| ACCEPTED |
| 1-25-10 |
| Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-517 |

30 of 44

MASTER LABEL FOR EPA REG. NO. 524-517
Registered Brand Names


Base Brand Name:
RangerPRO Herbicide

Table of Contents for Master Label

| | | |
|----|---|---------|
| I | Main Label for Industrial, Turf, & Ornamental Uses | Page 2 |
| II | Supplemental Labeling for Industrial, Turf, & Ornamental Uses | Page 34 |

****See each label part for more detailed table of contents****

All AG Crop uses and crop use supplemental labels previously on this registration have been removed. Ag non-crop uses are incorporated.



I. MAIN LABEL FOR INDUSTRIAL, TURF, ORNAMENTAL USES

[INSERT PRODUCT LOGO]



Ranger PRO herbicide is a complete broad-spectrum postemergence professional herbicide for industrial, turf and ornamental weed control.

Optional alternate statement: Ranger PRO herbicide is a complete broad-spectrum postemergence professional herbicide for forestry site preparation and utility rights-of-way weed control.

Complete Directions for Use

EPA Reg. No. 524-517

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product.

Use only according to label instructions.

Not all products listed in this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO (THIS COMPANY) DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

[Container label text for Refillable Container Label Statement]

THIS IS AN END-USE PRODUCT. MONSANTO (THIS COMPANY) DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A MONSANTO (VALID) REPACKAGING OR TOLL REPACKAGING AGREEMENT.

[Container label text for Non-Refillable Container Label Statement]

THIS IS AN END-USE PRODUCT. MONSANTO (THIS COMPANY) DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

- 1 1.0 INGREDIENTS
2 2.0 IMPORTANT PHONE NUMBERS
3 3.0 PRECAUTIONARY STATEMENTS
3.1 Hazards to Humans and Domestic Animals

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- 3.2 Environmental Hazards
- 3.3 Physical or Chemical Hazards
- 4 4.0 **STORAGE AND DISPOSAL**
- 5 5.0 **PRODUCT INFORMATION**
 - 5.1 Weed Resistance Management
 - 5.2 Management for Glyphosate Resistant Weed Biotypes
- 6 6.0 **MIXING**
 - 6.1 Mixing with Water
 - 6.2 Tank Mixtures
 - 6.3 Tank Mixing Procedure
 - 6.4 Mixing Percent Solutions
 - [Optional Section: Surfactants]
 - 6.5 Colorants or Dyes
 - 6.6 Drift Control Additives
- 7 7.0 **APPLICATION EQUIPMENT AND TECHNIQUES**
 - 7.1 Aerial Equipment
 - 7.2 Ground Broadcast Equipment
 - 7.3 Backpack or High-Volume Equipment
 - 7.4 Selective Equipment
 - 7.5 Injection Systems
 - 7.6 CDA Equipment
- 8 8.0 **SITE AND USE INSTRUCTIONS**
 - 8.1 Cut Stump
 - 8.2 Forestry Site Preparation
 - 8.3 Non-Crop Areas and Industrial Sites
 - 8.4 Habitat Management
 - 8.5 Hollow Stem Injection
 - 8.6 Injection and Frill (Woody Brush and Trees)
 - 8.7 Ornamentals, Plant Nurseries, and Christmas Trees
 - 8.8 Pasture Grasses, Forage Legumes
 - 8.9 Rangelands
 - 8.9 Parks, Recreational and Residential Areas
 - 8.10 Railroads
 - 8.11 Roadsides
 - 8.12 Utility Sites
- 9 9.0 **WEEDS CONTROLLED**
 - 9.1 Annual Weeds
 - 9.2 Perennial Weeds
 - 9.3 Woody Brush and Trees
- 10 10.0 **LIMIT OF WARRANTY AND LIABILITY**

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....41.0%

OTHER INGREDIENTS (including surfactant): 59.0%

100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent Nos. 5,683,958; 5,703,015; 6,063,733; 6,121,199; 6,121,200. No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE,
1-800-332-3111.
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE,
CALL COLLECT, DAY OR NIGHT,
(314)-694-4000.

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

| | |
|---|--|
| FIRST AID: | Call a poison control center or doctor for treatment advice. |
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses if present after the first 5 minutes then continue rinsing eye. |
| <ul style="list-style-type: none"> • Have the product container or label with you when calling a poison control center or doctor, or going for treatment. • You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information. • This product is identified as [INSERT BRAND NAME], EPA Registration No. 524-517. | |

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

70944

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.3 Physical or Chemical Hazards

Mix, store and apply spray solutions of this product using only stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto [insert company name] Supplemental Labeling or Fact Sheets. Supplemental labeling can be found on the Internet at www.cdms.net or www.greenbook.net websites or obtained by contacting your Authorized Monsanto [insert company name] Retailer or Monsanto Company [insert company name] Representative.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (40 CFR Part 170) for agricultural pesticides. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

4.0 STORAGE AND DISPOSAL

[FOR RIGID PLASTIC 2.5 GAL CONTAINERS OR OTHERS <= 5 GAL]

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. *[Alternate language]* Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container, including rinsate, by application in accordance with label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable Federal, State and local procedures

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state. *[Alternate language]* CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container.

CONTAINER HANDLING: Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once cleaned, some (optional: agricultural) plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-768-6387. If recycling is not available, puncture and dispose of in a sanitary landfill.

[Alternate disposal statement: Offer container for recycling, if available, or puncture and dispose of in a sanitary landfill.]

[FOR RIGID PLASTIC 30 GAL CONTAINERS OR OTHERS > 5 GAL]

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. *[Alternate language]* Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container, including rinsate, by application in accordance with label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable Federal, State and local procedures

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container.

[Alternate language] Nonrefillable container. Do not reuse this container to hold materials other

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.

CONTAINER HANDLING: Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once cleaned, some (optional: agricultural) plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-768-6387. If recycling is not available, puncture and dispose of in a sanitary landfill. [Alternate disposal statement: Offer container for recycling, if available, or puncture and dispose of in a sanitary landfill.]

[Optional Container Label Text] Return container to Monsanto for recycling. Contact 1-800-768-6387.

[Container label language for transport vehicles as defined in 40 CFR §156.3]

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, empty as much product from this transport vehicle as possible for repackaging or use in accordance with label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all precautions stated on this label until the container is cleaned, reconditioned or destroyed.

CONTAINER HANDLING: Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, and worn-out threads and closures. Clean thoroughly before reuse for transportation of a material of different composition or before retiring this transport vehicle from service.

5.0 PRODUCT INFORMATION

Product Description: This product is a postemergence, systemic herbicide with no residual soil activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant and no additional surfactant is needed or recommended. It may be applied through standard equipment after dilution and mixing with water or other carriers according to label instructions.

Optional alternate statement: It is formulated as a water-soluble liquid containing [X.X percent] surfactant and no additional surfactant is needed or recommended.

10044

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. See the "WEEDS CONTROLLED" sections of this label for specific weed rates.

Always use the higher product application rate in the range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area. Reduced weed control may result from treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, ensure spray coverage is uniform and complete. Do not spray foliage to the point of run-off.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Maximum Application Rates: The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as tank mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalents) does not exceed the maximum allowed. The combined total of all treatments must not exceed 10.6 quarts of this product (10.6 pounds of glyphosate acid) per acre per year. See the "INGREDIENTS" section of this label for necessary product information.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

110F49

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

5.1 Weed Resistance Management

GROUP 9 HERBICIDE

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or using other cultural or mechanical practices.

To minimize the occurrence of glyphosate-resistant biotypes observe the following good weed management practices:

- Scout your application site before and after herbicides applications.
- Control weeds early when they are relatively small.
- Incorporate other herbicides and cultural or mechanical practices as part of your weed control system where appropriate.
- Utilize the label rate for the most difficult weed in the site. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism) or tank mixtures which encourage rates of this product below the labeled amounts.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from site to site to minimize spread of weed seed.
- Use new commercial seed as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Monsanto *[insert company name]* representative, local retailer, or county extension agent.

5.2 Management for Glyphosate Resistant Weed Biotypes

NOTE: Appropriate testing is critical in order to confirm weed resistance to glyphosate. Contact your Monsanto *[inert company name]* representative to determine if resistance has been confirmed to any particular weed biotype in your area. Control instructions for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling or Fact Sheets for this product and may be obtained from your local retailer or Monsanto *[insert company name]* representative.

Since the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, Monsanto Company *[insert company name]* is not responsible for any losses that may result from the failure of this product to control glyphosate-resistant weed biotypes.

The following good weed management practices are encouraged to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present at your site, this product may be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices may also be used as appropriate.
- Scout treated sites after herbicide applications and control escapes of resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving sites known to contain resistant biotypes.

6.0 MIXING

Mix, store and apply spray solutions of this product using only clean stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

6.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of clean water. Add the labeled amount of this product near the end of the filling process and mix gently (well). During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

6.2 Tank Mixtures

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Always read and follow label directions for all products in the tank mixture. Use according to the most restrictive precautionary statements for each product in the mixture. Any labeled rate of this product may be used in a tank mix.

When this label describes a tank mixture with a generic active ingredient such as diuron, 2,4-D or dicamba, the user is responsible for ensuring the mixture product label allows the specific application.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly listed in this label. Mixing this product with herbicides or other materials not identified on this label may result in reduced performance.

6.3 Tank Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the labeled amount of this product.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50-mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance. Ensure that the specific tank mixture product is registered for application at the desired site.

Refer to the "Tank Mixtures" section for additional precautions.

6.4 Mixing Percent Solutions

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution

Amount of [INSERT BRAND NAME]

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| Desired Volume | 1/2% | 1% | 1-1/2% | 2% | 5% | 10% |
|----------------|--------|----------|-----------|----------|----------|--------|
| 1 gal | 2/3 oz | 1-1/3 oz | 2 oz | 2-2/3 oz | 6-1/2 oz | 13 oz |
| 25 gal | 1 pt | 1 qt | 1-1/2 qt | 2 qt | 5 qt | 10 qt |
| 100 gal | 2 qt | 1 gal | 1-1/2 gal | 2 gal | 5 gal | 10 gal |

2 tablespoons = 1 fluid ounce

For use in backpack, knapsack or pump-up sprayers, it is suggested that the labeled amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

Optional Section:

Surfactants

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

6.5 Colorants or Dyes

Approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's instructions.

6.6 Drift Control Additives

Drift reduction additives may be used with all equipment types, except wiper applicators, and sponge bars. When a drift reduction additive is used, read and carefully observe precautionary statements and all other information appearing on the additive label. The use of drift reduction additives can affect spray coverage which may result in reduced performance.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE VEGETATION.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and/or the grower is responsible for considering all these factors when making decisions.

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Use the labeled rates of this herbicide in 3 to 25 gallons of water per acre.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the air stream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application — To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aircraft Maintenance

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear is most susceptible.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of droplet size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the **Wind, Temperature and Humidity**, and **Temperature Inversion** sections of this label).

Controlling droplet size

- **Volume:** Use high flow-rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures labeled for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow-rates are needed, use higher flow-rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom Length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

- **Application Height:** Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance increases with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This product must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

7.2 Ground Broadcast Equipment

Apply the labeled rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified in this label or in separate supplemental labeling or fact sheets published by Monsanto [this company]. *(Optional statement: Use the labeled rates of this product in 10 to 60 gallons of water per acre as a broadcast spray unless otherwise specified in this label or in separate supplemental labeling or Fact Sheets published by Monsanto [this company])* As density of weeds increases, increase spray volume within the labeled range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

7.3 Backpack or High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, ensure spray coverage is uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

Refer to the "Annual Weeds" instructions of "WEEDS CONTROLLED" section for specific rates and restrictions

7.4 Selective Equipment

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

This product may be diluted with water and applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, to listed weeds growing in any site specified on this label.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION. Contact of this product with desirable vegetation may result in unwanted plant damage or destruction. **Recirculating Spray**

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

Adjust application equipment used above desired vegetation to the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation is likely to result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators and Sponge Bars

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced product performance may result from the use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators: Solutions ranging from 33 to 75 percent of this product in water may be used.

For Panel Applicators and pressure-feed systems: Solutions ranging from 33 to 100 percent of this product in water may be used.

When applied as directed, this product **CONTROLS** the following weeds:

| | |
|-----------------|------------------|
| Corn, volunteer | Sicklepod |
| Panicum, Texas | Spanishneedles |
| Rye, common | Starbur, bristly |
| Shattercane | |

When applied as directed, this product **SUPPRESSES** the following weeds:

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| | |
|------------------------|-----------------|
| Beggarweed, Florida | Ragweed, common |
| Bermudagrass | Ragweed, giant |
| Dogbane, hemp | Smutgrass |
| Dogfennel | Sunflower |
| Guineagrass | Thistle, Canada |
| Johnsongrass | Thistle, musk |
| Milkweed | Vaseygrass |
| Nightshade, silverleaf | Velvetleaf |
| Pigweed, redroot | |

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically directed.

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount directed in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 SITE AND USE INSTRUCTIONS

Detailed instructions follow alphabetically, by site.

Unless otherwise specified on this label or in separate supplemental labeling or fact sheets published by Monsanto [this company], applications may be made to control any weeds listed in the annual, perennial and woody brush tables. Also refer to the "SELECTIVE EQUIPMENT" section.

8.1 Cut Stump

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100-percent solution of this product to the freshly-cut surface **immediately after** cutting. Delays in application may result in reduced performance. For best results, make applications during periods of active growth and full leaf expansion.

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2 Forestry Site Preparation

This product is labeled for the control or partial control of woody brush, trees and herbaceous weeds in forestry sites. This product is also labeled for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is labeled for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

180444

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Use higher rates of this product within the labeled range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the labeled range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the labeled range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any labeled rate of this product may be used in a tank mix with the following products for forestry site preparation.

| | |
|---------------------------------|-----------|
| Arsenal Applicators Concentrate | Escort |
| Chopper | Garlon 3A |
| Chopper GEN2 | Garlon 4 |
| | Oust XP |

For control of herbaceous weeds, use the lower labeled tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher labeled tank mixture rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release unless otherwise directed on this label or in separately published Monsanto [insert company name] supplemental labeling or fact sheets,

8.3 Non-Crop Areas and Industrial Sites

This product may be used in non-crop areas, airports, apartment complexes, commercial sites, ditch banks, driveways, dry ditches, dry canals, fencerows, forestry sites, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, petroleum tank farms and pumping installations, railroads, recreational areas, residential areas, rights-of-way, roadsides, schools, shadehouses, sports complexes, storage areas, substations, turfgrass areas, utility sites, warehouse areas, other public areas, and wildlife management sites.

This product may also be used in non-food crop sites, Christmas tree farms, plant nurseries, sod or turf seed farms.

Unless otherwise specified, applications may be made to control any weeds listed in the WEEDS CONTROLLED section of this label.

Weed Control, Trim-and-Edge, Bare Ground

This product may be used in non-crop and non-food crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

TANK MIXTURES: This product may be tank mixed with the following products provided that the specific product is labeled for application at the use site. Refer to the individual product labels for approved sites and application rates.

| | |
|-----------------|----------------|
| Arsenal | Plateau |
| Clarity | Princep DF |
| Barricade 65WG | Princep Liquid |
| Diuron | Ronstar 50WSP |
| Endurance | Sahara |
| Escort | simazine |
| Garlon 3A | Surflan |
| Garlon 4 | Telar |
| Karmex | Vanquish |
| Krovar I DF | 2,4-D |
| | |
| Oust XP | |
| Pendulum 3.3 EC | |
| Pendulum WDG | |

This product plus dicamba tank mixtures may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 1 to 2 quarts of this product plus 2 to 8 ounces of Oust XP per acre.

| | |
|--------------|---------------|
| Bahiagrass | Fescue, tall |
| Bermudagrass | Johnsongrass |
| Broomsedge | Poorjoe |
| Dallisgrass | Quackgrass |
| Dock, curly | Vaseygrass |
| Dogfennel | Vervain, blue |

Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing - Annuals

For growth suppression of some annual grasses, annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Make applications when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

Bromus Species and Medusahead in Pastures and Rangelands

Bromus species. This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 8 to 16 fluid ounces of this product per acre on a broadcast basis.

For best results, coincide treatment with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Make

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

applications to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead. To treat medusahead, apply 16 fluid ounces of this product per acre as soon as plants are actively growing, and prior to the 4-leaf stage. Applications may be made in the fall or spring.

Applications to brome and medusahead may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre. When applied as directed in this label, there are no grazing restrictions.

Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring green-up.

Apply 8 to 64 fluid ounces of this product per acre. Apply the labeled rates in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces per acre may result in injury or delayed green-up in highly maintained areas, golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust XP in highly maintained turfgrass areas. For further uses, refer to the "RAILROADS" section of this label, which gives rates for dormant bermudagrass treatment and the "ROADSIDES" section of this label, which gives rates for dormant bermudagrass and bahiagrass treatments.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. DO NOT apply more than 16 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust XP in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses like bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Delay tillage or renovation techniques such as vertical mowing, coring or slicing for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

8.4 Habitat Management

Habitat Restoration and Management

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.5 Hollow Stem Injection

Castorbean (*Ricinus communis*)

Inject 5 ml per plant of this product into the lower portion of the main stem.

Hemlock, Poison (*Conium maculatum*)

Inject one leaf cane per plant 10 to 12 inches above root crown with 6 ml of a 5% v/v solution of this product.

Hogweed, Giant (*Heracleum mantegazzianum*)

Inject one leaf cane per plant 12 inches above root crown with 6 ml of a 5% v/v solution of this product.

Horsetail, Field (*Equisetum arvense*)

Inject one segment above the root crown with 0.6 ml per stem of this product. Use a small syringe that calibrates to this rate.

Knotweed, Bohemian (*Polygonum bohemicum*), Giant (*Polygonum sachalinense*), Japanese (*Polygonum cuspidatum*)

Inject 6 ml per stem of this product between second and third internode.

Reed, Giant (*Arundo donax*)

Inject 8 ml per stem of this product between the second and third internode.

Thistle, Canada (*Cirsium arvense*)

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.6 ml per stem of this product is injected into the stem.

NOTE: Based on the maximum annual use rate of glyphosate for these non-crop sites, the combined total for all treatments must not exceed 10.6 quarts of this product per acre. At 6 ml per stem, 10.6 quarts treats approximately 1700 stems.

8.6 Injection and Frill (Woody Brush and Trees)

This product may be used to control or partially control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 1/25 fluid ounce (1 ml) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In these species, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of this product. For best results, make application during periods of active growth and after full leaf expansion.

8.7 Ornamentals, Plant Nurseries, and Christmas Trees

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Post-Directed, Trim-and-Edge

This product may be used prior to the planting of and as a post-directed spray around established woody ornamental species, arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jobo, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew, growing in plant nurseries, on Christmas tree farms, or on other non-food tree production sites. This product may also be used to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a nursery setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. THIS PRODUCT IS NOT LABELED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be exercised to avoid contact of spray, drift or mist with foliage or bark of established ornamental species.

Site Preparation

This product may be used prior to planting any ornamental, nursery or Christmas tree species.

Wiper Applications

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established eucalyptus or poplar trees. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

Greenhouse/Shadehouse

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

8.8 PASTURE GRASSES, FORAGE LEGUMES

Alfalfa, Clover, and Other Forage Legumes

This product may be used on Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, and Vetch (all types). This product can be applied preplant, preemergence, at-planting, or for renovation.

This product may also be used as a spot treatment (Alfalfa and Clover Only), over-the-top wiper applications (Alfalfa and Clover Only), or preharvest (Alfalfa Only).

Preplant, Preemergence, At-Planting

This product may be applied before, during or after planting crops listed in this section. Applications must be made prior to emergence of the crop.

If a single application is made at rates of 2 quarts per acre or less, no waiting period between treatment and feeding or grazing is required. If application rates greater than 2 quarts per acre are made, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Renovation

This product may be applied as a broadcast spray to renovate existing stands of alfalfa, clover, and other labeled forage legumes.

Remove domestic livestock before application. If application rates of 2 quarts per acre or less are used wait 36 hours after application before grazing or harvesting. If application rates greater than 2 quarts per acre are used, wait 8 weeks between applications and grazing or harvesting.

Spot Treatment, Over-the-Top Wiper Applications (Alfalfa and Clover Only)

This product may be applied as a spot treatment in alfalfa or clover. This product may be applied with wiper applicators to control or suppress the weeds listed below. Applications may be made in the same area at 30-day intervals.

230f44

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. Treat no more than 10 percent of the total field area at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

When applied as directed, this product CONTROLS the following weeds:

- Corn, volunteer
- Panicum, Texas
- Rye, common
- Shattercane
- Sicklepod
- Spanishneedles
- Starbur, bristly

When applied as directed, this product SUPPRESSES the following weeds:

- Beggarweed, Florida
- Bermudagrass
- Dogbane, hemp
- Dogfennel
- Guineagrass
- Johnsongrass
- Milkweed
- Nightshade, silverleaf
- Pigweed, redroot
- Ragweed, common
- Ragweed, giant
- Smutgrass
- Sunflower
- Thistle, Canada
- Thistle, musk
- Vaseygrass
- Velvetleaf

Preharvest (Alfalfa Only)

This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This application will severely injure or destroy the stand of alfalfa. This product will control annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa. The treated crop and weeds can be harvested and fed to livestock after 36 hours. Allow a minimum of 36 hours between application and harvest. Applications may be made at any time of the year. Make only one application to an existing stand of alfalfa per year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

Do not apply more than 2 quarts of this product per acre as a preharvest treatment. Preharvest application to alfalfa grown for seed may result in a reduction in germination or crop vigor.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

8.9 Rangelands

This product can be applied postemergence and will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands.

Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds.

Delay grazing of treated areas to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Do not use ammonium sulfate when spraying rangeland grasses with this product. Do not apply more than 3 quarts per acre per year.

Apply 12 to 16 fluid ounces of this product per acre to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Make Fall applications where spring moisture is limited and fall germination allows for good weed growth.

For medusahead, apply 16 fluid ounces of this product per acre at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

8.10 Parks, Recreational and Residential Areas

This product may be used in parks, recreational and residential areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around trees, fences, and paths, around buildings, sidewalks, and other objects in these areas. This product may be used for spot treatment of unwanted vegetation. This product may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the "Non-Crop Areas and Industrial Sites" section apply to park and recreational areas.

8.11 Railroads

All of the instructions in the "Non-Crop Areas and Industrial Sites" section apply to railroads.

Bare Ground, Ballast and Shoulders, Crossings, and Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

This product may be tank mixed with the following products provided that the specific product is registered for ballast, shoulder, spot, bare ground and crossing treatments:

| | |
|-----------|-------------|
| Arsenal | Krovar I DF |
| Clarity | Oust XP |
| Diuron | Sahara |
| Escort | Spike |
| Garlon 3A | Telar |
| Garlon 4 | Vanquish |

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| | |
|---------|-------|
| Hyvar X | 2,4-D |
|---------|-------|

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 3/4 to 2-percent solution of this product when using high-volume spray-to-wet applications. Apply a 5- to 10-percent solution of this product when using low volume directed sprays for spot treatment. This product may be mixed with the following products for enhanced control of woody brush and trees:

| | |
|-----------|----------|
| Arsenal | Garlon 4 |
| Escort | Tordon K |
| Garlon 3A | |

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

| | |
|------------------|----------------|
| Bahiagrass | Johnsongrass |
| Bluestem, silver | Trumpetcreeper |
| Fescue, tall | Vaseygrass |

This product may be tank-mixed with Oust XP. If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

| | |
|------------------|----------------|
| Bahiagrass | Fescue, tall |
| Blackberry | Johnsongrass |
| Bluestem, silver | Poorjoe |
| Broomsedge | Raspberry |
| Dallisgrass | Trumpetcreeper |
| Dewberry | Vaseygrass |
| Dock, curly | Vervain, blue |
| Dogfennel | |

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Do not makerepeat applications in the same season since severe injury may occur.

8.12 Roadsides

All of the instructions in the "Non-Crop Areas and Industrial Sites" section apply to roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails and Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Tank Mixtures

This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments provided that the specific product is registered for use on such sites. Read and carefully observe label directions, cautionary statements and all information on the labels of all products used.:

| | |
|-----------------|----------------|
| Clarity | Princep DF |
| Diuron | Princep Liquid |
| Endurance | Ronstar 50WSP |
| Escort | Sahara |
| Krovar I DF | simazine |
| Oust XP | Surflan |
| Outrider | Telar |
| Pendulum 3.3 EC | Vanquish |
| Pendulum WDG | 2,4-D |

See the "Tank Mixture" section of this label for tank mixing instructions.

Release of Bermudagrass or Bahiagrass

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring green-up. This product may also be tank-mixed with Outrider or Oust XP for residual control. Tank mixtures of this product with Oust XP may delay green-up.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 8 to 64 fluid ounces of this product in a tank mixture with 3/4 to 1-1/3 ounces Outrider herbicide per acre. Read and follow all label directions for Outrider herbicide.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 1/4 to 1 ounce per acre of Oust XP. Apply the labeled rates in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in green-up and minimize injury, add no more than 1 ounce of Oust XP per acre on bermudagrass and no more than 1/2 ounce of Oust XP per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 1 to 3 pints of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

| | |
|------------------|----------------|
| Bahiagrass | Johnsongrass |
| Bluestem, silver | Trumpetcreeper |
| Fescue, tall | Vaseygrass |

This product may be tank mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 8 to 32 fluid ounces of this product with 3/4 to 1 1/3 ounces of Outrider. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank mixed with Oust XP. If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust XP label. Use the higher

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

| | |
|------------------|----------------|
| Bahiagrass | Fescue, tall |
| Bluestem, silver | Johnsongrass |
| Broomsedge | Poorjoe |
| Dallisgrass | Trumpetcreeper |
| Dock, curly | Vaseygrass |
| Dogfennel | Vervain, blue |

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season causes severe injury.

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

This product may be used for control or partial control of Johnsongrass and other weeds listed on the Outrider label in actively growing bahiagrass. Apply 1-1/2 to 4-3/4 ounces of this product with 0.75 to 1.33 ounces of Outrider per acre. Use the higher rates for control of perennial weeds or annual weeds greater than 6 inches in height. Use only on well established bahiagrass.

A tank mixture of this product plus Oust XP may be used. Apply 6 fluid ounces of this product plus 1/4 ounce of Oust XP per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.13 Utility Sites

This product is labeled for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

This product may be used in utility sites and substations for bare ground, trim-and-edge, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product is also labeled for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees provided that the specific product is registered for application to the desired site. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any labeled rate of this product may be used in a tank mix.

For control of herbaceous weeds, use the lower labeled tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher labeled rates.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

TANK MIXTURES: This product may be tank mixed with the following products for use in utility sites, provided that the specific product used is labeled for use on these sites. Refer to the individual product labels for approved sites and application rates.

| | |
|------------------------|----------------------------|
| Arsenal | atrazine ¹ |
| Barricade 65WG | dicamba ¹ |
| diuron ¹ | Endurance |
| Escort | Escort XP |
| Garlon 3A ² | Garlon 4 ³ |
| Krenite | Krovar I DF |
| Oust | Oust XP |
| Outrider | pendimethalin ¹ |
| Plateau | |
| Ronstar 50WP | Sahara |
| simazine ¹ | Surflan AS |
| Surflan WDG | Telar DF |
| Transline | Vanquish |
| Velpar DF | Velpar L |
| 2,4-D ¹ | |

- ¹ Tank mixtures with product containing this generic active ingredient may be made provided the specific product is labeled for application at the use site.
- ² Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray incompatibility problems.
- ³ For side trimming treatments, use this product alone or in a tank mixture with Garlon 4.

Bare Ground and Trim-and-Edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product may be tank mixed with the following products. Refer to these products' labels for approved non-crop sites and application rates.

| | |
|-----------------|-----------------|
| Arsenal | Plateau™ |
| Banvel | Princep™DF |
| Barricade™ 65WG | Princep™ Liquid |
| Diuron | Ronstar™ 50WP |
| Endurance™ | Sahara™ |
| Escort | simazine |
| Garlon 3A | Surflan™ |

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the labeled range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

290944

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

For low-volume directed spray applications, use a 5- to 10-percent solution of this product for control or partial control of annual weeds, perennial weeds, or woody brush and trees. Spray coverage must be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple resprouts.

Refer to the following label sections for labeled rates for the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results.

9.1 Annual Weeds

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions.

For spray-to-wet applications, apply a 1/2-percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall.

WEED SPECIES

| |
|--|
| Anoda, spurred |
| Barley* |
| Barley, little * |
| Barnyardgrass* |
| Bassia, fivehook |
| Bittercress* |
| Bluegrass, annual* |
| Bluegrass, bulbous* |
| Brome, downy* |
| Brome, Japanese* |
| Buttercup* |
| Castorbean |
| Cheatgrass* |
| Cheeseweed (<i>Malva parviflora</i>) |
| Chervil* |
| Chickweed* |
| Cocklebur* |
| Copperleaf, hophornbeam |
| Copperleaf, Virginia |
| Coreopsis, plains/tickseed * |
| Corn* |
| Crabgrass* |
| Cupgrass, woolly * |
| Dwarfdandelion* |
| Eclipta* |
| Falsedandelion* |
| Falseflax, smallseed* |
| Fiddleneck |
| Filaree |
| Fleabane, annual* |
| Fleabane, hairy (<i>Conyza bonariensis</i>)* |

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| |
|--|
| Fleabane, rough* |
| Foxtail* |
| Foxtail, Carolina * |
| Geranium, Carolina |
| Goatgrass, jointed* |
| Goosegrass |
| Groundsel, common* |
| Henbit |
| Horseweed/Marestail (<i>Conyza canadensis</i>) |
| Itchgrass* |
| Johnsongrass, seedling |
| Junglerice |
| Knotweed |
| Kochia |
| Lamb's-quarters* |
| Lettuce, prickly * |
| Mannagrass, eastern * |
| Mayweed |
| Medusahead* |
| Morningglory (<i>Ipomoea spp</i>) |
| Mustard, blue* |
| Mustard, tansy* |
| Mustard, tumble* |
| Mustard, wild* |
| Nightshade, black * |
| Oats |
| Panicum browntop * |
| Panicum, fall * |
| Panicum, Texas * |
| Pennycress, field * |
| Pepperweed, Virginia * |
| Pigweed* |
| Puncturevine |
| Purslane, common |
| Pusley, Florida |
| Ragweed, common* |
| Ragweed, giant |
| Rice, red |
| Rocket, London * |
| Rocket, yellow |
| Rye* |
| Ryegrass* |
| Sandbur, field* |
| Sesbania, hemp |
| Shattercane* |
| Shepherd's-purse* |
| Sicklepod |
| Signalgrass, broadleaf* |
| Smartweed, ladysthumb* |
| Smartweed, Pennsylvania* |
| Sorghum, grain (milo)* |
| Sowthistle, annual |
| Spanishneedles |

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| |
|----------------------|
| Speedwell, corn * |
| Speedwell, purslane* |
| Sprangletop* |
| Spurge, annual |
| Spurge, prostrate* |
| Spurge, spotted* |
| Spurry, umbrella* |
| Starthistle, yellow |
| Stinkgrass* |
| Sunflower* |
| Teaweed/Prickly sida |
| Thistle, Russian |
| Velvetleaf |
| Wheat* |
| Wild oats* |
| Witchgrass* |

*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 1 pint of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the labeled range.

Use a 2-percent solution on tough-to-control perennials like bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low-volume directed spot treatments, apply a 5- to 10-percent solution of this product.

Allow 7 or more days after application before tillage.

| Weed Species | Rate (QT/A) | Hand-Held % Solution |
|--------------|-------------|----------------------|
|--------------|-------------|----------------------|

| | | |
|--|-----|-------|
| Alfalfa* | 1 | 2 |
| Alligatorweed* | 4 | 1.5 |
| Anise (fennel) | 2-4 | 1-2 |
| Artichoke, Jerusalem | 3-5 | 2 |
| Bahiagrass | 3-5 | 2 |
| Beachgrass, European (<i>Ammophila arenaria</i>) | -- | 5 |
| Bentgrass* | 1.5 | 2 |
| Bermudagrass | 5 | 2 |
| Bermudagrass, water (knotgrass) | 1.5 | 2 |
| Bindweed, field | 4-5 | 2 |
| Bluegrass, Kentucky | 2 | 2 |
| Blueweed, Texas | 4-5 | 2 |
| Brackenfern | 3-4 | 1-1.5 |
| Bromegrass, smooth | 2 | 2 |
| Bursage, woolly-leaf | -- | 2 |
| Canarygrass, reed | 2-3 | 2 |
| Cattail | 3-5 | 2 |

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| | | |
|--------------------------|-----|--------|
| Clover; red, white | 3-5 | 2 |
| Cogongrass | 3-5 | 2 |
| Dallisgrass | 3-5 | 2 |
| Dandelion | 3-5 | 2 |
| Dock, curly | 3-5 | 2 |
| Dogbane, hemp | 4 | 2 |
| Fescue (except tall) | 3-5 | 2 |
| Fescue, tall | 1-3 | 2 |
| Guineagrass | 3 | 1 |
| Hemlock, poison | 2-4 | 1-2 |
| Horsenettle | 3-5 | 2 |
| Horseradish | 4 | 2 |
| Iceplant | 2 | 1.5-2 |
| Ivy, German | 2-4 | 1-2 |
| Johnsongrass | 2-3 | 1 |
| Kikuyugrass | 2-3 | 2 |
| Knapweed | 4 | 2 |
| Lantana | -- | 1-1.25 |
| Lespedeza | 3-5 | 2 |
| Milkweed, common | 3 | 2 |
| Muhly, wirestem | 2 | 2 |
| Mullein, common | 3-5 | 2 |
| Napierrgrass | 3-5 | 2 |
| Nightshade, silverleaf | 2 | 2 |
| Nutsedge; purple, yellow | 3 | 1-2 |
| Orchardgrass | 2 | 2 |
| Pampasgrass | 3-5 | 1.5-2 |
| Paragrass | 3-5 | 2 |
| Pepperweed, perennial | 4 | 2 |
| Phragmites* | 3-5 | 1-2 |
| Quackgrass | 2-3 | 2 |
| Redvine* | 2 | 2 |
| Reed, giant | 4-5 | 2 |
| Ryegrass, perennial | 2-3 | 1 |
| Smartweed, swamp | 3-5 | 2 |
| Spurge, leafy* | -- | 2 |
| Sweet potato, wild* | -- | 2 |
| Thistle, artichoke | 2-3 | 1-2 |
| Thistle, Canada | 2-3 | 2 |
| Timothy | 2-3 | 2 |
| Torpedograss* | 4-5 | 2 |
| Trumpetcreeper* | 2-3 | 2 |
| Vaseygrass | 3-5 | 2 |
| Velvetgrass | 3-5 | 2 |
| Wheatgrass, western | 2-3 | 2 |

*Partial control

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed in this label or separate supplemental label or Fact Sheet published by Monsanto [*this company*]. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low-volume directed-spray spot treatments, apply a 5- to 10-percent solution of this product.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

| Weed Species | Broadcast Rate (QT/A) | Hand-Held Spray-to-Wet % Solution |
|----------------------------|-----------------------|-----------------------------------|
| Alder | 3-4 | 1-1.5 |
| Ash* | 2-5 | 1-2 |
| Aspen, quaking | 2-3 | 1-1.5 |
| Bearclover (Bearmat)* | 2-5 | 1-2 |
| Beech* | 2-5 | 1-2 |
| Birch | 2 | 1 |
| Blackberry | 3-4 | 1-1.5 |
| Blackgum* | 2-5 | 1-2 |
| Bracken | 2-5 | 1-2 |
| Broom; French, Scotch | 2-5 | 1.5-2 |
| Buckwheat, California* | 2-4 | 1-2 |
| Cascara* | 2-5 | 1-2 |
| Catsclaw* | -- | 1-1.5 |
| Ceanothus* | 2-5 | 1-2 |
| Chamise* | 2-5 | 1 |
| Cherry; bitter, black, pin | 2-3 | 1-1.5 |
| Coyotebrush | 3-4 | 1.5-2 |
| Creepers, Virginia | 2-5 | 1-2 |
| Deerweed | 2-5 | 1 |
| Dogwood* | 2-5 | 1-2 |
| Elderberry | 2 | 1 |
| Elm* | 2-5 | 1-2 |
| Eucalyptus | -- | 2 |
| Flower, monkey flower* | 2-4 | 1-2 |
| Gorse* | 2-5 | 1-2 |
| Hasardia* | 2-4 | 1-2 |
| Hawthorn | 2-3 | 1-1.5 |
| Hazel | 2 | 1 |
| Hickory* | 2-5 | 1-2 |
| Honeysuckle | 3-4 | 1-1.5 |
| Hornbeam, American* | 2-5 | 1-2 |
| Ivy, poison | 4-5 | 2 |
| Kudzu | 4 | 2 |
| Locust, black* | 2-4 | 1-2 |
| Madrone resprouts* | -- | 2 |
| Manzanita* | 2-5 | 1-2 |
| Maple, red* | 2-4 | 1-1.5 |
| Maple, sugar | -- | 1-1.5 |
| Maple, vine * | 2-5 | 1-2 |

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

| | | |
|--|-----|-------|
| Oak; black, white* | 2-4 | 1-2 |
| Oak, post | 3-4 | 1-1.5 |
| Oak; northern pin | 2-4 | 1-1.5 |
| Oak, poison | 4-5 | 2 |
| Oak, scrub* | 2-4 | 1-1.5 |
| Oak, southern red | 2-3 | 1-1.5 |
| Olive, Russian * | 2-5 | 1-2 |
| Peppertree, Brazilian (Florida holly)* | 2-5 | 1-2 |
| Persimmon* | 2-5 | 1-2 |
| Pine | 2-5 | 1-2 |
| Poplar, yellow* | 2-5 | 1-2 |
| Redbud, eastern | 2-5 | 1-2 |
| Rose, multiflora | 2 | 1 |
| Sage, black | 2-4 | 1 |
| Sage, white* | 2-4 | 1-2 |
| Sage brush, California | 2-4 | 1 |
| Salmonberry | 2 | 1 |
| Saltcedar* | 2-5 | 1-2 |
| Sassafras* | 2-5 | 1-2 |
| Sourwood* | 2-5 | 1-2 |
| Sumac; laurel, poison, smooth, sugarbush, winged * | 2-4 | 1-2 |
| Sweetgum | 2-3 | 1-1.5 |
| Swordfern* | 2-5 | 1-2 |
| Tallowtree, Chinese | -- | 1 |
| Tanoak resprouts* | -- | 2 |
| Thimbleberry | 2 | 1 |
| Tobacco, tree* | 2-4 | 1-2 |
| Toyon* | - | 2 |
| Trumpetcreeper | 2-3 | 1-1.5 |
| Waxmyrtle, southern* | 2-5 | 1-2 |
| Willow | 3 | 1 |
| Yerba Santa, California* | - | 2 |

*Partial control

10.0 LIMIT OF WARRANTY AND LIABILITY

Monsanto Company [This company] warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

35 of 44

I. MAIN LABEL FOR INDUSTRIAL, TURF, AND ORNAMENTAL USES

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

[INSERT BRAND NAME], Outrider, Monsanto and Vine symbol are trademarks of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

This product is protected by U.S. Patent Nos. 5,683,958; 5,703,015; 6,063,733; 6,121,199; 6,121,200. No license granted under any non-U.S. patent(s).

EPA Reg. No. 524-517

In case of an emergency involving this product,
or for medical assistance,
Call Collect, day or night, (314) 694-4000.

©[DATE] MONSANTO COMPANY [INSERT COMPANY NAME]

Packed For: *[Produced (Manufactured) for:]*

[Insert company name & mailing address]

Monsanto Company

800 N. Lindbergh Blvd.

ST. LOUIS, MISSOURI, 63167 U.S.A.

II. SUPPLEMENTAL LABELING FOR INDUSTRIAL, TURF, & ORNAMENTAL USES

Table of Contents: Industrial, Turf, and Ornamental Supplemental labeling

| | Name | EPA APPROVAL |
|---|---|-------------------------|
| A | FOR USE FOR SELECTIVE WEED CONTROL ON [INSERT BRAND NAME] TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS. | Nov 20, 2002 |
| B | [INSERT BRAND NAME] HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS, INCLUDING RAILROAD RIGHTS-OF-WAY, SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES | Nov 20, 2002 |
| C | AERIAL APPLICATIONS IN CALIFORNIA | Amended this submission |
| D | FOR USE IN CITRUS CROPS IN FLORIDA | June 19, 2008 |

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **[INSERT BRAND NAME]** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **[INSERT BRAND NAME]** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

[INSERT BRAND NAME HERE]

Herbicide

[INSERT SUPPLEMENTAL TITLE HERE]

EPA Reg. No. 524-517

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

[INSERT BRAND NAME] is a trademark of Monsanto Technology LLC *[insert Company name]*.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "PRODUCT INFORMATION" and "MIXING" sections of the label booklet for **[INSERT BRAND NAME]** for essential product performance information.

[INSERT SPECIFIC DIRECTIONS FOR USE HERE]

Read the "Limit of Warranty and Liability" in the label booklet for **[INSERT BRAND NAME]** before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [DATE] MONSANTO COMPANY

800 N. Lindbergh Blvd.

ST. LOUIS, MISSOURI 63167

[Insert company name and mailing address]

380944

A. FOR USE FOR SELECTIVE WEED CONTROL ON [INSERT BRAND NAME] TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

[INSERT BRAND NAME] Tolerant Tall Fescue Selections For Seed Production

Use this product on [INSERT BRAND NAME] tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 4 to 16 fluid ounces per acre as a postemergence spray on [INSERT BRAND NAME] tolerant tall fescue selections. See the label booklet for application instructions, rates, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the [INSERT BRAND NAME] label booklet for a complete list of weeds controlled or suppressed.

NOTE: The labeled rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with **INSERT BRAND NAME** tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

Timing Of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy and in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

B. [INSERT BRAND NAME] HERBICIDE AND TANK MIXTURES FOR NON-CROP AREAS, INCLUDING RAILROAD RIGHTS-OF-WAY, SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the [INSERT BRAND NAME] label booklet for application rates. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quarts per acre of this product may be used for improved results.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product provided that the specific product is registered for application to the desired site. Any labeled rate of this product may be used in a tank mixture with these products.

Tank-mix Product

| | |
|-------------------------|--|
| Arsenal™ | |
| Banvel | |
| 2,4-D | |
| Garlon™ 3A | |
| Garlon 4 | |
| Diuron | |
| Diuron + 2,4-D | |
| Diuron + Garlon 3A | |
| Diuron + Garlon 4 | |
| Hyvar™ X | |
| Hyvar X + 2,4-D | |
| Hyvar X + Garlon 3A | |
| Hyvar X + Garlon 4 | |
| Krovar™ I DF | |
| Krovar I DF + 2,4-D | |
| Krovar I DF + Garlon 3A | |
| Krovar I DF + Garlon 4 | |
| Oust XP™ | |
| Oust XP + 2,4-D | |
| Oust XP + Garlon 3A | |
| Oust XP + Garlon 4 | |
| Spike™ 80W | |
| Spike 80W + 2,4-D | |
| Spike 80W + Garlon 3A | |
| Spike 80W + Garlon 4 | |

Arsenal is not approved for use in the state of California.

Refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, labeled use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mix products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

When used in combination as recommended by Monsanto Company [this company], the liability of Monsanto [this company] shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Monsanto [this] product in such combination use.

| |
|--|
| <p>C. LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRESNO COUNTY, CALIFORNIA</p> |
|--|

DIRECTIONS FOR USE

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this supplemental labeling and in the product label booklet. Refer to Aerial Equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the product label for additional information. Refer to the individual use site section of the product label, or to other supplemental labeling or technical fact sheets published separately for this product by Monsanto [*insert company name*], for specific use instructions.

AVOID DRIFT—DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be used for aerial application in California. Tank mixtures with 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and for alfalfa and pasture renovation applications only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

ADDITIONAL INFORMATION FOR FRESNO COUNTY, CALIFORNIA

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

North: Fresno County line
 South: Fresno County line
 East: State Highway 99
 West: Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Recommendations

A written recommendation MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation MUST state the proximity of surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Applications at Night—Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

To report known or suspected misuse of this product, call 1-800-332-3111.

For additional information on the proper aerial application of this product, call 916-784-1718.

420944

D. FOR USE IN CITRUS CROPS

DIRECTIONS FOR USE

LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor

TYPES OF APPLICATION: Preplant (Site Preparation) Broadcast Spray, Middles (between rows), Strips (within rows), Shielded Sprayer, Hooded Sprayer, Wiper Applicator, Directed Spray, Spot Treatment

USE INSTRUCTIONS: Applications may be made with boom equipment, CDA equipment, shielded sprayer, hand-held and high-volume wand, lance, orchard gun or with wiper applicator equipment, except as directed. Apply this product according to the rates listed in the "ANNUAL WEEDS" and "PERENNIAL WEEDS" sections of the Complete Directions for Use booklet for [Insert Brand Name] herbicide. Use the higher rate in the labeled range when weeds are stressed, growing in dense populations or greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts of product per acre per year.

For burndown or control of the weeds listed below, apply the labeled rate of this product in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 2 to 3 quarts of this product per acre in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Krovar I or Karmex may improve weed control.

Perennial Weeds:

S = Suppression B = Burndown
PC = Partial Control C = Control

| WEED SPECIES | [Insert Brand Name] Herbicide RATE PER ACRE | | | |
|-------------------|---|------|------|------|
| | 1 QT | 2 QT | 3 QT | 5 QT |
| Bermudagrass | B | — | PC | C |
| Guineagrass | | | | |
| Florida Ridge | B | C | C | C |
| Florida Flatwoods | — | B | C | C |
| Paragrass | B | C | C | C |
| Torpedograss | S | — | PC | C |

Refer to the individual product labels for listing of specific crops, rates, geographic restrictions and precautionary statements.

Middles (between rows)

USE INSTRUCTIONS: This product will control or suppress annual and perennial weeds, and ground covers growing in the middles (between rows of trees). If weeds are under drought stress, irrigate prior to application. Reduced weed control may result if weeds have been recently mowed at the time of application.

TANK MIXTURES: A tank mixture of this product plus Goal 2XL may be applied for annual weed control in-between rows (middles) of citrus trees, especially when weeds are stressed or growing in dense populations. Application of 16 to 32 fluid ounces of this product, plus 3 to 12 fluid ounces of Goal 2XL, per acre will control annual weeds with a maximum height of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). This tank mixture will also control common cheeseweed (malva) or hairy fleabane (*Conyza bonariensis*) with a maximum height or diameter of 3 inches. Read and follow label directions for all products in the tank mixture.

Strips (in rows)

USE INSTRUCTIONS: For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used, and then only where there is sufficient clearance, in order to minimize the potential for overspray or drift of this product onto the crop.

TANK MIXTURES: This product may be applied within rows of trees in tank mixtures with the following products:

Devrinol 50-DF, Direx 4L, Goal 2XL, Karmex DF, pendimethalin, Princep Caliber 90, Simazine 4L, Sim-Trol 4L, Solicam DF, Surflan AS,

Refer to the tank mixture product label for restrictions and precautions; use according to the most restrictive precautionary statements for each product in the tank mixture.

PRECAUTIONS, RESTRICTIONS: Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or bark of trunk, branches, suckers, fruit or other parts of trees. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used where there is the potential for crop contact, and then only where there is sufficient clearance. See additional use instructions and precautions in the "**Application Equipment And Techniques**" section of the directions for use booklet for [Insert Brand Name].

Allow a minimum of 1 day between last application and harvest in citrus crops. Allow a minimum of 3 days between application and transplanting trees.

For citron groves, apply as a directed spray only.

44
/ 44

Revisions this submission for 000524-00517.20091214.amend.pdf

Removal of alternate brand names

Support of Ranger PRO as base brand name

Changes requested by EPA October 29, 2009 approval to remove general.

Change/delete uses of recommend and other advisory statements as requested on other recent EPA approvals resulting in slight re-wording of some sentences.