

## Supplemental Labeling

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container for each product and on this supplemental labeling.

### Roundup WeatherMAX Herbicide Logo

EPA Reg. No. 524-537

Roundup WeatherMAX and Roundup Ready are registered trademarks of Monsanto Company

**FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT HORSEWEED  
(MARESTAIL, *Conyza canadensis*) IN COTTON, CORN, AND SOYBEANS.**

Keep out of reach of children.  
**CAUTION!**

In case of an emergency involving this product, Call Collect, day or night (314) 694-4000.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of herbicide application. **AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS).**

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

For ground applications, use 10 to 20 gallons of water per acre. For aerial applications, use 3 to 15 gallons of water per acre. **For tank-mix recommendations, read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental labeling or fact sheets published separately for all herbicides used.**

#### Cotton:

##### Preplant:

For control of horseweed, apply this product (22 fluid ounces per acre) in a tank-mix with Clarity® (8 fluid ounces per acre). This application must be made 21 to 35 days before planting and before horseweed reaches 6 inches in height. In order to avoid crop injury, a minimum interval of 21 days during which there is at least 1 inch of cumulative rainfall must be observed between Clarity application and planting of cotton.

ACCEPTED

FEB - 6 2003

In the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
registered for the pesticide  
with label  
524-537

**Post-directed (Roundup Ready® Cotton varieties only):**

Management of early season weed competition and the development of a crop height differential between cotton and the horseweed is often achieved by a combination of preplant burndown and post emergent over-the-top and/or directed applications of Roundup WeatherMAX herbicide. These measures enhance the development of a height differential that is necessary to successfully make post-directed treatments. In-crop post-directed applications of MSMA (2 pounds active ingredient per acre) tank-mixed with diuron (0.5 to 0.75 pounds active ingredient per acre) should be made when the temperature is 80°F or higher.

**Soybeans:**

**Preplant:**

Apply a tank mixture of this product (22 fluid ounces per acre) with 2,4-D (0.5 pounds a.i. per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting. For areas where 2,4-D cannot be applied due to application restrictions or proximity to a sensitive crop, contact your local retailer and Monsanto representative.

**In-crop (Roundup Ready Soybean varieties only):**

It is strongly encouraged that horseweed be controlled prior to planting using recommended preplant burndown treatments. In-crop Roundup Ready soybeans, apply a tank mixture of this product (22 fluid ounces per acre) with Amplify™ (0.3 ounces per acre). This treatment should be used as a salvage treatment only for a horseweed infestation that was not controlled preplant. Application should be made between full emergence of the first trifoliate leaf and 50% flowering stage of soybeans. At the time of treatment, horseweed should not exceed 6 inches in height.

**Corn:**

**Preplant, At-Planting, Preemergence:**

Apply a tank mixture of this product (22 fluid ounces per acre) plus 2,4-D (0.5 pounds a.i. of per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

Atrazine (1 to 2 pounds active ingredient per acre) may be included in the tank-mixture to provide residual control. Refer to the atrazine product label for specific use instructions.

**In-crop (Roundup Ready Corn hybrids only):**

In-crop Roundup Ready corn, apply a tank-mixture of this product (22 fluid ounces per acre) plus Clarity (8 to 16 fluid ounces per acre) or 2,4-D (0.5 to 1.0 pounds a.i. per acre). Apply between corn emergence and the 5-leaf stage of growth (approximately 8 inches tall).

**ATTENTION**

**AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.**

Amplify, Roundup WeatherMAX, Roundup Ready, and Transorb II are registered trademarks of Monsanto Company. Clarity is a trademark of BASF Corp.