

PLEASE NOTE

This image contains more than one label approved for this product on this date.

SUPPLEMENTAL LABELING

For distribution and use only in the States of Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, and Washington


READ THE ENTIRE LABEL FOR Roundup WeatherMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX HERBICIDE LOGO

EPA Reg. No. 524-537

Roundup WeatherMAX is a
trademark
of Monsanto Technology LLC



PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY PEAS, LENTILS, AND CHICKPEAS.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

DIRECTIONS FOR USE

See the "APPLICATION EQUIPMENT AND TECHNIQUES" Section of the Roundup WeatherMAX herbicide label booklet for more information.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray

drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

2/2

BROADCAST SPRAY

USE INSTRUCTIONS: This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 17 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTIONS, RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup WeatherMAX herbicide label.
- Preharvest application is not recommended for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

SPOT TREATMENTS

USE INSTRUCTIONS: This product may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in peas, lentils, and chickpeas. Apply up to 17 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

PRECAUTIONS, RESTRICTIONS:

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine spot treatments with a preharvest broadcast spray on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup WeatherMAX herbicide label.
- Do not feed treated vines and hay from these crops to livestock.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2002 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

NEXT

LABEL

524-537

5/20/2004

1/2

SUPPLEMENTAL LABELING


READ THE ENTIRE LABEL FOR Roundup WeatherMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX HERBICIDE LOGO

EPA Reg. No. 524-537

Roundup WeatherMAX is a
trademark
of Monsanto Technology LLC



PREHARVEST AND SPOT-TREATMENTS OF WEEDS IN DRY BEANS

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

DIRECTIONS FOR USE

See the "APPLICATION EQUIPMENT AND TECHNIQUES" Section of the Roundup WeatherMAX herbicide label booklet for more information.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

2/2

BROADCAST SPRAY

USE INSTRUCTIONS: This product may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 22 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

PRECAUTIONS, RESTRICTIONS:

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup WeatherMAX herbicide label.
- Preharvest application is not recommended for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system.
- Do not treat cowpeas, since this crop is considered to be grown as livestock feed.

SPOT TREATMENTS

USE INSTRUCTIONS: This product may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 22 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

PRECAUTIONS, RESTRICTIONS:

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine spot treatments with a preharvest broadcast spray on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the Roundup WeatherMAX herbicide label.
- Do not feed treated vines and hay from these crops to livestock.
- Do not treat cowpeas, since this crop is considered to be grown as livestock feed.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2002 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167


NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



Roundup WeatherMAX
Herbicide

EPA Reg. No. 524-537

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup WeatherMAX is a registered trademark of Monsanto Technology LLC.

FOR AID IN SUGARCANE RIPENING

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX for essential product performance information.

This product is a foliar-applied plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane. It is effective in both low and high-tonnage sugarcane.

When applied as directed under the conditions described, this product will hasten ripening and extend the period of high sucrose level in sugarcane.

As a result of leaf desiccation, improved trash burn can be expected.

Most of the sucrose increase is concentrated in the top nodes of the treated cane stalk. In order to recover the maximum sugar where topping is practiced during harvest, top at the base of the fourth leaf.

Prior to application, consult your state sugarcane authority or local Monsanto representative regarding the degree of sucrose response anticipated from the variety of sugarcane to be treated. Do not plant subsequent crops in treated fields other than the following for 30 days after application: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybeans, squash (all types) or wheat.

PRECAUTIONS, RESTRICTIONS: Application of this product may initiate development of shooting eyes. This product may not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2 to 3 weeks after application, this product may produce a slight yellowing to pronounced browning and drying of leaves, and a shortening of upper internodes. Spindle death may occur.

Rainfall within 6 hours after application may reduce effectiveness.

Application is not recommended for sugarcane grown for seed, as a reduction in germination or vigor may occur.

Do not feed or graze treated sugarcane forage following application.

Do not apply for enhanced ripening to any crops other than sugarcane. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

APPLICATION RATES:

Use the following application rates and timing instructions according to the State in which the sugarcane is grown.

NOTE: Use the higher rate within the recommended range when treating sugarcane under adverse ripening conditions or when less responsive varieties are to be treated.

FLORIDA-Apply 5 to 12 fluid ounces of this product per acre 3 to 6 weeks before harvest of LAST RATOON CANE ONLY.

HAWAII-Apply 9 to 21 fluid ounces of this product per acre 4 to 10 weeks before harvest.

LOUISIANA-Apply 3.5 to 12 fluid ounces of this product per acre 3 to 7 weeks before harvest of RATOON CANE ONLY.

PUERTO RICO-Apply 5 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

TEXAS-Apply 5 to 12 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX
Herbicide

ROUNDUP WEATHERMAX FOR WEED CONTROL IN VEGETABLE CROPS AND MISCELLANEOUS CROPS

EPA Reg. No. 524-537

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

Roundup WeatherMAX is a registered trademark of Monsanto Technology LLC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.


AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX for essential product performance information.

1 Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL LISTED VEGETABLE CROPS WITHIN THE SECTIONS GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, Prior to Transplanting Vegetables, At-Planting, Hooded Sprayers in Row Middles, Shielded Sprayers in Row Middles, Wiper Applications in Row Middles, Post-Harvest, Directed Applications (Non-Bearing Ginseng), Over-the-Top Wiper Applications (Rutabagas Only).



PRECAUTIONS, RESTRICTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to insure that the wash water flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for additional information.

1.1 Brassica Vegetables

LABELED CROPS: Broccoli, Chinese broccoli (gai lan), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

1.2 Bulb Vegetables

LABELED CROPS: Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

1.3 Cucurbit Vegetables and Fruits

LABELED CROPS: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Melons (all), *Momordica spp* (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon.

PRECAUTIONS, RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

1.4 Leafy Vegetables

LABELED CROPS: Amaranth (Chinese spinach), Arugula (roquette), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

PRECAUTIONS, RESTRICTIONS: For Watercress, avoid applications within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of injury.

1.5 Fruiting Vegetables

LABELED CROPS: Eggplant, Groundcherry (*Physalis spp*), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.

PRECAUTIONS, RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For tomato and tomatillo, do not make hooded or shielded sprayer applications in row middles because of the potential for crop injury.

1.6 Legume Vegetables (Succulent or Dried)

LABELED CROPS: Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

1.7 Root and Tuber Vegetables

LABELED CROPS: Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

Directed Applications (Non-Bearing Ginseng Only)

USE INSTRUCTIONS: This product may be used for general weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, and orchard guns or with wiper application equipment.

PRECAUTIONS, RESTRICTIONS: Direct applications so that there is no contact of this product with the ginseng plant. Applications must be made at least one year prior to harvest.

Over-the-Top Wiper Applications (Rutabagas Only)

USE INSTRUCTIONS: Wiper applicators may be used over-the-top of rutabagas.

PRECAUTIONS, RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas.

2 Miscellaneous Crops

LABELED CROPS: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following: General Weed Control, Site Preparation, Spot Treatment (Asparagus), Post-Harvest (Asparagus).

For Roundup Ready sugar beets, see the "ROUNDUP READY CROPS" section of the label booklet.

PRECAUTIONS, RESTRICTIONS: Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of the label booklet for additional information.

General Weed Control, Site Preparation

USE INSTRUCTIONS: This product may be applied for general weed control or for site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS, RESTRICTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to insure that the wash water flushes off the plastic mulch and does not enter transplant holes. Allow at least 21 days between residue removal and transplanting. Applications made at emergence will result in injury or death to emerged seedlings.

Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.

Spot Treatment (Asparagus)

USE INSTRUCTIONS: This product may be applied immediately after cutting, but prior to the emergence of new spears.

PRECAUTIONS, RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Post-Harvest (Asparagus)

USE INSTRUCTIONS: This product may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears.


PRECAUTIONS, RESTRICTIONS: Direct contact of the spray with the asparagus may result in serious crop injury. Select and use recommended types of spray equipment for postemergence post-harvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© 2004 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

NEXT

LABEL



SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX Herbicide

EPA Reg. No. 524-537

Roundup WeatherMAX and Roundup Ready are trademarks of Monsanto Technology LLC.

FOR COTTON WITH THE ROUNDUP READY GENE FOR APPLICATIONS IN THE STATE OF ARIZONA ONLY.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

General Information

ATTENTION: MONSANTO RECOMMENDS THIS PRODUCT FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN CROPS WITH THE ROUNDUP READY GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

ROUNDUP READY COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "ROUNDUP READY", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup WeatherMAX herbicide for essential product performance information.

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Over-the-top, Selective Equipment, Preharvest

Maximum Allowable Combined Application Quantities Per Season

Combined total per year for all applications	5.3 quarts per acre
Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total in-crop applications from ground cracking to layby	2.5 quarts per acre
<ul style="list-style-type: none"> Total in-crop over-the-top from ground cracking to 4-leaf stage 	2.0 quarts per acre
<ul style="list-style-type: none"> Total in-crop applications using selective equipment through layby 	1.3 quarts per acre
Maximum preharvest application rate	44 fluid ounces per acre

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of the Roundup WeatherMAX herbicide label booklet for general precautionary instructions for use in Roundup Ready crops. Do NOT combine these instructions with other recommendations made for crop varieties that do not contain the Roundup Ready gene, in the "ANNUAL AND PERENNIAL

CROPS (Alphabetical)" section of the Roundup WeatherMAX herbicide label booklet. The combined total application of this product from cotton emergence until harvest must not exceed 4 quarts per acre.

NO MORE THAN TWO OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT.

NO MORE THAN TWO POST-DIRECTED APPLICATIONS SHOULD BE MADE FROM THE 5-LEAF STAGE THROUGH LAYBY.

SEQUENTIAL IN-CROP OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT MUST BE AT LEAST 10 DAYS APART AND COTTON MUST HAVE AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS.

ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST. APPLICATIONS MADE IN EXCESS OF MAXIMUM LABEL RATES ARE EXPECTED TO RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS AND ARE THE SOLE RESPONSIBILITY OF THE GROWER.

THE MAXIMUM USE RATES STATED THROUGHOUT THIS PRODUCT'S LABELING APPLY TO THIS PRODUCT COMBINED WITH THE USE OF ALL OTHER HERBICIDES CONTAINING GLYPHOSATE OR SULFOSATE AS THE ACTIVE INGREDIENT, WHETHER APPLIED AS MIXTURES OR SEPARATELY. CALCULATE THE APPLICATION RATES AND ENSURE THAT THE TOTAL USE OF THIS AND OTHER GLYPHOSATE OR SULFOSATE CONTAINING PRODUCTS DOES NOT EXCEED STATED MAXIMUM USE RATE.

Preplant, At-Planting, and Preemergence:

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton.

Over-the-Top:

USE INSTRUCTIONS: This product may be applied by aerial or ground application equipment at rates up to 22 fluid ounces per acre per application postemergence to Roundup Ready cotton from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Unless otherwise directed in this supplemental label, any single over-the-top broadcast application should not exceed 22 fluid ounces per acre. Combined over-the-top applications between ground cracking until the 4-leaf (node) stage should not exceed 2.0 quarts per acre.

NOTE: For specific rates of application and instructions, refer to the "ANNUAL" and "PERENNIAL WEEDS RATE TABLES" in the label booklet for Roundup WeatherMAX herbicide.

Selective Equipment:

USE INSTRUCTIONS: This product may be applied in-crop using precision post-directed or hooded sprayers to Roundup Ready cotton through layby. Unless otherwise directed in this supplemental label, any single application using selective equipment should not exceed 22 fluid ounces per acre.

Sequential in-crop applications using selective equipment may be made up to a maximum of 1.3 quarts per acre.

At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches).

PRECAUTIONS, RESTRICTIONS: See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the Roundup WeatherMAX herbicide label booklet for information on proper use and calibration of this equipment. The combined total of in-crop over-the-top plus selective equipment applications must not exceed 2.6 quarts per acre.

Salvage Treatment:

USE INSTRUCTIONS: From the ground cracking stage through layby, where weeds threaten to cause the loss of the crop, applications of up to 32 fluid ounces per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: CROP TOLERANCE OF ROUNDUP READY COTTON HAS NOT BEEN FULLY TESTED AT THIS APPLICATION RATE. SALVAGE TREATMENTS ARE EXPECTED TO RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS AND ARE THE SOLE RESPONSIBILITY OF THE GROWER. NO MORE THAN TWO SALVAGE TREATMENTS SHOULD BE USED PER GROWING SEASON.**

PRECAUTIONS, RESTRICTIONS: The combined total of in-crop over-the-top plus selective equipment applications must not exceed 2.6 quarts per acre.

Preharvest:

USE INSTRUCTIONS: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20 percent boll crack. Up to 1.3 quarts per acre of this product may be applied using either aerial or ground spray equipment. **NOTE:** This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Do not apply Roundup WeatherMAX herbicide preharvest to Roundup Ready cotton grown for seed.

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT

IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX herbicide before using. For over-the-top uses on Roundup Ready crop varieties crop safety and weed control performance are not warranted by Monsanto when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2004 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167