

PLEASE NOTE

This image contains more than one label approved for this product on this date.

SUPPLEMENTAL LABELING


READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX®
Herbicide

EPA Reg. No. 524-537

WeatherMAX and Roundup Ready are trademarks of Monsanto Company



Tank Mixtures for In-Crop Use with Roundup Ready® Alfalfa

Keep out of reach of children

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

ROUNDUP READY CROPS CONTAIN A PATENTED GENE THAT PROVIDES TOLERANCE TO GLYPHOSATE, THE ACTIVE INGREDIENT IN THIS PRODUCT. THIS PRODUCT WILL CAUSE SEVERE CROP INJURY OR DESTRUCTION AND YIELD LOSS IF APPLIED TO CROPS THAT ARE NOT GLYPHOSATE TOLERANT. AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS THAT DO NOT CONTAIN A GLYPHOSATE TOLERANCE GENE, SINCE SEVERE PLANT INJURY OR DESTRUCTION WILL RESULT.

See the "GENERAL INFORMATION", "MIXING" and "ROUNDUP READY CROPS" sections of the label booklet for Roundup WeatherMAX herbicide for essential product information.

DIRECTIONS FOR USE

New Stand Establishment (Seeding Year)

TANK MIXTURES: This product may be applied postemergence (in-crop) over the top of Roundup Ready alfalfa in the seeding year at up to 44 fluid ounces per acre in a tank mixture with the following products. Application should be made after weeds have emerged but before alfalfa growth or regrowth interferes with spray coverage of the target weeds. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to alfalfa. Read and follow label directions of all products in the tank mixture.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa:

clethodim, imazamox, imazethapyr, sethoxydim, quizalofop p-ethyl

Assure II, Poast, Prism, Pursuit, Raptor, SelectMAX]

[Optional statement: Pursuit or Raptor applied to seedling alfalfa may result in a temporary reduction in growth. Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury may result.]

Established Stands (Non-seeding Year)

TANK MIXTURES: This product may be applied postemergence (in-crop) over the top of established stands of Roundup Ready alfalfa in tank mixtures described below according to the growing conditions of the crop. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to alfalfa. Read and follow label directions of all products in the tank mixture

Actively growing alfalfa: For control of emerged annual grasses and broadleaf weeds when alfalfa is actively growing, this product may be applied at up to 44 fluid ounces per acre in a tank mixture with the following herbicides.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa:

clethodim, imazamox, imazethapyr, sethoxydim
Assure II, Poast, Prism, Pursuit, Raptor, SelectMAX]

[Optional statement: Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury may result.]

Dormant alfalfa: For control of emerged annual grasses and broadleaf weeds when alfalfa is dormant, this product may be applied at up to 44 fluid ounces per acre in a tank mixture with the following herbicides. Apply when the daily temperature remains above freezing.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use on alfalfa:

imazamox, metribuzin, propyzamide
Kerb 50-W, Lexone, Pursuit, Raptor, Sencor]

[Optional statement: Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury may result.]


PRECAUTIONS, RESTRICTIONS: Tank mixtures of this product with herbicides, insecticides, or fungicides may result in crop injury and/or reduced weed control. Applications should not be made to frozen or snow covered ground. Refer to individual tank mixture product label for restrictions and precautions; use according to the most restrictive precautionary statements for each product in the tank mixture.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX Herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [YEAR] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

[INSERT DATE]

[INSERT PRINT PLATE NUMBER]



NEXT

LABEL


SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP WEATHERMAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup WeatherMAX Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

Roundup WeatherMAX®
Herbicide

EPA Reg. No. 524-537
WeatherMAX and Roundup Ready are trademarks of Monsanto Company



Tank Mixtures for In-Crop Use with Roundup Ready® Sugar Beet

Keep out of reach of children

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

ROUNDUP READY CROPS CONTAIN A PATENTED GENE THAT PROVIDES TOLERANCE TO GLYPHOSATE, THE ACTIVE INGREDIENT IN THIS PRODUCT. THIS PRODUCT WILL CAUSE SEVERE CROP INJURY OR DESTRUCTION AND YIELD LOSS IF APPLIED TO CROPS THAT ARE NOT GLYPHOSATE TOLERANT. AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS THAT DO NOT CONTAIN A GLYPHOSATE TOLERANCE GENE, SINCE SEVERE PLANT INJURY OR DESTRUCTION WILL RESULT.

See the "GENERAL INFORMATION", "MIXING" and "ROUNDUP READY CROPS" sections of the label booklet for Roundup WeatherMAX herbicide for essential product information.

DIRECTIONS FOR USE

TANK MIXTURES: This product may be tank-mixed with the following products and applied postemergence (in-crop) over the top of Roundup Ready sugar beet. Ensure that the specific product being used in the tank mixture is registered for application postemergence (in-crop) to sugar beet. Read and follow label directions of all products in the tank mixture

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use postemergence (in-crop) to sugar beet:

clethodim, clopyralid, desmedipham, dimethenamid, ethofumesate, s-metolachlor, phenmedipham, quizalofop-p-ethyl, trisulfuron-methyl

Assure II, Betamix, Betanex, Dual MAGNUM, Nortron SC, Outlook, Progress, Select, Stinger, Upbeet]

[Optional statement: Betamix, Betanex, Nortron SC and Progress can cause significant sugar beet injury. Refer to these product labels for crop injury precautions.]

PRECAUTIONS, RESTRICTIONS: The combined total application of this product from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 32 fluid ounces per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 22 fluid ounces per acre. Allow a minimum of 30 days between last application and sugar beet harvest. Tank mixtures of this product with herbicides,

282

insecticides or fungicides may result in crop injury and/or reduced weed control. Refer to individual tank mixture product label for restrictions and precautions; use according to the most restrictive precautionary statements for each product in the tank mixture.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup WeatherMAX Herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [YEAR] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

[INSERT DATE]

[PRINT PLATE NUMBER]