

PLEASE NOTE

This image contains more than one label approved for this product on this date.

all products used in the tank mixture by mixing equal proportional quantities of this product with other herbicides may cause incompatibility, antagonism, or a reduction in product efficacy. Monsanto has not tested all tank mix product formulations for compatibility, performance, and crop safety.

Management of glyphosate resistant ryegrass:

524-539

05/18/2006

1/2

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ORIGINAL MAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup Original MAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container for each product and on this supplemental labeling.

Roundup Original MAX Herbicide

EPA Reg. No. 524-539


Roundup Original MAX and Roundup Ready are trademarks of Monsanto Technology LLC.

FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT RYEGRASS

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night (314) 694-4000.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS).

GENERAL INFORMATION

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Original MAX herbicide for essential product performance information.

This product may be tank-mixed with other herbicides for application in accordance with label directions. Follow all precautions and use instructions contained within the product's labeling in accordance with the most restrictive label limitations. Some products have the potential to cause crop injury under certain conditions, at certain crop growth stages and/or other circumstances. Read all labels for products used in tank mixtures to determine the potential for injury prior to use. Always predetermine the compatibility of all products used in the tank mixture by mixing small proportional quantities in advance. A tank mixture of this product with other herbicides may cause incompatibility, antagonism, or a reduction in product efficacy. Monsanto has not tested all tank mix product formulations for compatibility, performance, and crop safety.

Management of glyphosate resistant ryegrass:

If a naturally occurring glyphosate resistant biotype of ryegrass is present, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control. Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.

Not all herbicides are registered in each state or for all use-sites for the management of ryegrass. When Roundup Original MAX herbicide is used in combination with other herbicides, refer to each product's label and observe all precautions and limitations on the label.

Tree and Vine Crops:

Preemergence to glyphosate resistant ryegrass:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with a preemergence herbicide product such as oryzalin for preemergence ryegrass control.

Postemergence with a residual to emerged glyphosate resistant ryegrass:

Apply Roundup Original MAX herbicide, to control other emerged weeds, in a tank-mix with a preemergence herbicide product such as oryzalin and a graminicide product such as Poast. Make applications before ryegrass exceeds 4" in height.

Postemergence to glyphosate resistant ryegrass:

Apply Roundup Original MAX herbicide to control emerged weeds in a tank-mix with a graminicide product such as Poast. Make applications before ryegrass exceeds 4" in height.


NEXT

LABEL

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP ORIGINAL MAX HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup Original MAX herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container for each product and on this supplemental labeling.



Roundup Original MAX Herbicide

EPA Reg. No. 524-539

Roundup Original MAX and Roundup Ready are trademarks of Monsanto Technology LLC.

FOR CONTROL AND MANAGEMENT OF GLYPHOSATE RESISTANT PALMER AMARANTH (*Amaranthus palmeri*) IN COTTON, CORN, AND SOYBEANS

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS).

GENERAL INFORMATION

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Original MAX herbicide for essential product performance information.

This product may be tank-mixed with other herbicides for application in accordance with label directions. Follow all precautions and use instructions contained within the product's labeling in accordance with the most restrictive label limitations. Some products have the potential to cause crop injury under certain conditions, at certain crop growth stages, and/or other circumstances. Read all labels for products used in tank mixtures to determine the potential for injury prior to use. Always predetermine the compatibility of all products used in the tank mixture by mixing small proportional quantities in advance. A tank mixture of this product with other herbicides may cause incompatibility, antagonism, or a reduction in product efficacy. Monsanto has not tested all tank mix product formulations for compatibility, performance, and crop safety.

Management of glyphosate resistant Palmer amaranth:

If a naturally occurring glyphosate resistant biotype of Palmer amaranth is present, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of

action to achieve control. Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.

Not all herbicides are registered in each state or for all sites or crops for the management of Palmer amaranth. When Roundup Original MAX is used in combination with other herbicides, refer to each product's label and observe all precautions and limitations on the label.

Tillage or a burndown herbicide application is encouraged prior to planting.

Cotton:

Preplant, At-Planting, Preemergence:

Apply Roundup Original MAX herbicide to control emerged weeds, in a tank-mix with a preemergence soil residual herbicide product labeled for control of Palmer amaranth such as pendimethalin (Prowl H20) and/or fluometuron (Cotoran).

Postemergence in Roundup Ready Flex Cotton and Roundup Ready Cotton (In-crop):

Preemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with metolachlor (prior to the 4-leaf stage in Roundup Ready cotton) for the control of Palmer amaranth.

Preemergence to glyphosate resistant Palmer amaranth at layby:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with a residual herbicide product such as diuron (Direx) or flumioxazin (Valor) post-directed for the control of Palmer amaranth.

Postemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX, to control emerged weeds, in a tank-mix of MSMA and diuron (Direx) or flumioxazin (Valor), to control emerged Palmer amaranth.

Soybeans:

Preplant, At-Planting, Preemergence:

Preemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with a preemergence residual herbicide product such as alachlor (INTRRO) or another residual herbicide for control of Palmer amaranth. For preplant applications 2,4-D may be added to the tank-mix to help control emerged broadleaf weeds. Follow label instructions regarding application timing relative to soybean planting.

Postemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with lactofen (Cobra) or fomesafen (Flexstar) to control emerged Palmer amaranth.

Corn:

Preplant, At-Planting, Preemergence:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with a preemergence residual herbicide product such as Harness Xtra, Harness Xtra 5.6L, Degree Xtra or another residual herbicide for the control of Palmer amaranth.

Postemergence in Roundup Ready Corn 2 (in-crop):

Preemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX herbicide, to control emerged weeds, in a tank-mix with a preemergence residual herbicide product such as Harness Xtra, Harness Xtra 5.6L, Degree Xtra or another residual herbicide for the control of Palmer amaranth.

Postemergence to glyphosate resistant Palmer amaranth:

Apply Roundup Original MAX herbicide in a tank-mix with other herbicides such as 2,4-D or dicamba (Clarity, Banvel, or Distinct) to control emerged weeds and a residual herbicide product such as Harness Xtra, Harness Xtra 5.6L, Degree Xtra for continue control of Palmer amaranth. Observe the labeled maximum annual application rate and timing restrictions.