



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 6, 2023

Gaganpreet Kaur  
Senior Regulatory Affairs Manager  
Bayer CropScience LP  
700 Chesterfield Parkway West  
Chesterfield, Missouri 63017 USA

Subject: Label Amendment –Incorporating supplemental labels, and extension of expiration date on supplemental labels  
Product Name: ROUNDUP ORIGINAL 2K HERBICIDE  
EPA Registration Number: 524-539  
Application Dates: 3/17/2023 and 3/27/2023  
Case Numbers: 480415, 483998, 475547, 475866

Dear Gaganpreet Kaur:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find

Page 2 of 2

EPA Reg. No. 524-539

Case No. 480415, 483998, 475547, 475866

or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Sarah Meadows at 202-566-2828 or at [meadows.sarah@epa.gov](mailto:meadows.sarah@epa.gov).

Sincerely,



Emily Schmid, Product Manager 25  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

# MASTER LABEL FOR EPA REG. NO. 524-539

Primary Brand Name:

**Roundup Original 2K Herbicide**

Alternate Brand Names:

**Roundup Original MAX Herbicide  
RT Master II Herbicide  
Honcho K6 Herbicide**

**A C C E P T E D**

**11/6/2023**

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 524-539

## Master Label Table of Contents

I.	Directions for Use with Food and Feed Crops	2 - 113
II.	Directions for Use on Industrial, Turf and Ornamental Sites	114 -167

**\*\*See each label part for more detailed table of contents\*\***

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

GLYPHOSATE

GROUP

9

HERBICIDE

[INSERT BRAND NAME] [Logo]



Complete Directions for Use

EPA Reg. No. 524-539

A [Optional text: complete] broad-spectrum postemergence herbicide for weed control in many agricultural systems

[Alternative product statement: Selective broad-spectrum weed control in Roundup Ready® crops]

[Alternative product statement: Non-selective, broad-spectrum weed control for many agricultural systems and farmsteads]

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC AT (800) 424-9300]

[Optional label text: For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 hours a day 1-800-334-7577]

[Optional label text: For **PRODUCT USE** information Call 1-866-99BAYER (1-866-992-2937)]

[Optional statement for limited-geography product distribution: This product is not registered in all states.]

[Alternative or additional statements for limited-geography product distribution: For control of annual and perennial weeds in [list states and or county-level information, as appropriate, where product is registered and distributed]. [Optional text: \*County Distribution:] [Optional text: see inside for details.] [Optional text: In [list states] this product is distributed in the counties listed below:] [list counties by states]

[Optional statement for limited-geography product distribution: This product is not registered for use in [list states, if any, where this product is not registered for use]

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, [Optional text, if applicable: EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® AND OTHER LISTED GLYPHOSATE-TOLERANT CROPS,] AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

[Optional label text: Transorb® (Logo)]

[Optional label text: Transorb® II Technology (Logo)]

[Optional label text: Transorb® PLUS (Logo)]

[Optional label text: Transorb® PLUS Technology (Logo)]

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

[Optional label text: Trisorb® (Logo)]

[Optional label text: CROPSHIELD® Formulation]

[Optional label text: CROPSHIELD® Formula – Specially formulated for Roundup Ready® crops (CROPSHIELD® Logo)]

[Optional label text: Herbicide for Roundup Ready® [Optional text: and other listed Glyphosate-Tolerant] Crops]

[Optional label text: Roundup® – Powerful Performance at a Practical Price]

[Optional label text: Roundup Ready PLUS™ – Weed Management Solutions]

[Optional label text: A member of the Roundup® Family of [Optional text: Agricultural] Herbicides by Bayer CropScience]

[Optional label statement, if applicable: See attached labeling [Alternative text: See inside] for Complete Directions for Use [Optional text: in English and Spanish].

[Optional label statement for container labels with attached labeling: See Complete Directions for Use attached to this label [Alternative text: See attached label booklet] for complete [Optional text, as applicable: Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use [and] Limit of Warranty and Liability] [Optional text: in English and Spanish]

Not all products listed on this label are registered for use in California. Check the registration status of each product in California before using.

Read the entire label before using this product.

Use only according to label directions.

Read the LIMIT OF WARRANTY AND LIABILITY statement [Optional text, if applicable: at the end of this labeling] [Alternative optional text, if applicable: on side panel] [Alternative optional text, if statement not included on the container label: at the end of the attached labeling] before buying or using. If terms are not acceptable, return at once unopened.

[Insert net contents, as appropriate: NET [insert appropriate value or leave blank line for refillable container] GAL [or other appropriate unit of measure] [Alternative text: NET CONTENTS] [Alternative label statement for transport vehicles only: NET [Optional: CONTENTS]: See Bill of Lading]

LOT [Insert Lot number or blank space] [Alternative label statement for transport vehicles only: LOT: See Bill of Lading]

[Optional text for container labels: ATTACH COMPLETE DIRECTIONS FOR USE HERE]

[Optional text for container labels with attached Directions for Use labeling that will be hidden behind the attached labeling and only become visible if the labeling is removed: COMPLETE DIRECTIONS FOR USE HAVE BEEN REMOVED]

### CONTENTS

<b>1</b>	<b>1.0</b>	<b>INGREDIENTS</b>
<b>2</b>	<b>2.0</b>	<b>IMPORTANT PHONE NUMBERS</b>
<b>3</b>	<b>3.0</b>	<b>PRECAUTIONARY STATEMENTS</b>
	3.1	Hazards to Humans and Domestic Animals
	3.2	Environmental Hazards
	3.3	Physical or Chemical Hazards

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

- 4 4.0 STORAGE AND DISPOSAL**
- 5 5.0 PRODUCT INFORMATION**
- 6 6.0 WEED RESISTANCE MANAGEMENT**
  - 6.1 Weed Management Practices
  - 6.2 Management of Glyphosate-Resistant Biotypes
- 7 7.0 MIXING**
  - 7.1 Mixing with Water
  - 7.2 Tank Mixtures
  - 7.3 Tank-Mixing Procedure
  - 7.4 Mixing Spray Solution Concentrations
  - 7.5 Surfactants [*this section is optional in the final printed label*]
  - 7.6 Ammonium Sulfate
  - 7.7 Colorants and Dyes
  - 7.8 Drift Reduction Additives
- 8 8.0 APPLICATION EQUIPMENT AND TECHNIQUES**
  - 8.1 Spray Drift Management
  - 8.2 Aerial Application Equipment
  - 8.3 Ground Application Equipment
  - 8.4 Handheld Sprayers
  - 8.5 Selective Application Equipment
  - 8.6 Injection Systems
  - 8.7 Controlled Droplet Applicator (CDA)
- 9 9.0 ANNUAL AND PERENNIAL CROPS**
  - 9.1 Cereal and Grain Crops
  - 9.2 Corn
  - 9.3 Cotton
  - 9.4 Fallow Systems
  - 9.5 Grain Sorghum (Milo)
  - 9.6 Herbs and Spices
  - 9.7 Oilseed Crops
  - 9.8 Soybean
  - 9.9 Sugarcane
  - 9.10 Vegetable Crops
  - 9.11 Miscellaneous Crops
- 10 10.0 TREE, VINE, AND SHRUB CROPS**
  - 10.1 Berry and Small Fruit Crops
  - 10.2 Citrus Fruit Crops
  - 10.3 Pome Fruit Crops
  - 10.4 Stone Fruit Crops
  - 10.5 Tree Nut Crops
  - 10.6 Tropical and Subtropical Trees and Fruit Crops
  - 10.7 Vine Crops
  - 10.8 Miscellaneous Tree Food Crops
  - 10.9 Non-Food Tree Crops
- 11 11.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELAND**
  - 11.1 Alfalfa, Clover, and Other Forage Legumes
  - 11.2 Conservation Reserve Program (CRP)
  - 11.3 Grass Seed and Sod Production

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

- 11.4 Pastures
- 11.5 Rangeland
- 12 12.0 ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS**
  - 12.1 Alfalfa with Roundup Ready Technology
  - 12.2 Canola with Roundup Ready Technology (Spring Varieties)
  - 12.3 Canola with Roundup Ready Technology (Winter Varieties)
  - 12.4 Canola with TruFlex Technology (Spring Varieties)
  - 12.5 Field Corn Hybrids with Roundup Ready 2 Technology
  - 12.6 Sweet Corn Hybrids with Roundup Ready 2 Technology
  - 12.7 Field Corn Hybrids with Agrisure GT Technology
  - 12.8 Cotton with Roundup Ready Technology
  - 12.9 Cotton with Roundup Ready Flex Technology and XtendFlex Technology
  - 12.10 Cotton with GlyTol Technology
  - 12.11 Soybean with Roundup Ready Technology
  - 12.12 Soybean with Roundup Ready 2 Yield Technology, Roundup Ready 2 Xtend Technology, and XtendFlex Technology
  - 12.13 Soybean with GT27 Technology and Enlist E3 Technology
  - 12.14 Sugarbeet with Roundup Ready Technology
- 13 13.0 FARMSTEAD USE**
  - 13.1 Farmstead Weed Control, Trim-and-Edge
  - 13.2 Greenhouse/Shadehouse
  - 13.3 Chemical Mowing
  - 13.4 Cut Stump Application
  - 13.5 Habitat Management
- 14 14.0 ANNUAL WEEDS RATE SECTION**
  - 14.1 Annual Weeds—Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K
  - 14.2 Annual Weeds—Handheld Sprayers
  - 14.3 Annual Weeds—Tank Mixtures for Fallow and Reduced Tillage Systems
- 15 15.0 PERENNIAL WEEDS RATE SECTION**
- 16 16.0 WOODY BRUSH, TREES AND VINES RATE SECTION**
- 17 17.0 TANK-MIX PRODUCT INFORMATION** [*this section printed only if required*]
- 18 18.0 LIMIT OF WARRANTY AND LIABILITY**

**1.0 INGREDIENTS**

ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its potassium salt..... 48.7%

OTHER INGREDIENTS: ..... 51.3%

100.0%

\*Contains 660 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 5.5 pounds per U.S. gallon, which is equivalent to 540 grams of the acid, glyphosate, per liter or 4.5 pounds per U.S. gallon (39.8% by weight).

[Optional label text that will be updated at the time of printing, if necessary: This product is protected by [Optional text: one or more of the following] U.S. Patent No's. 5,668,085, RE 37,866 and 6,365,551.]

[Optional label text, if applicable: Other Patents Pending.]

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

[Optional label text, if applicable: No license granted under any non-U.S. patent(s.)

[Option to insert reference to a Patent Website: For a list of patents, if any, covering this product or its use, go to [www.monsantotechnology.com](http://www.monsantotechnology.com).]

EPA Est. [Insert appropriate EPA establishment number: 524-IA-1; 524-LA-1; or Other; or blank space]  
[Optional text: (Establishment Number when entered here supersedes all others)]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites, as appropriate.]

**2.0 IMPORTANT PHONE NUMBERS**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 hours a day **1-800-334-577**

For **PRODUCT USE** information Call **1-866-99BAYER (1-866-992-2937)**

**3.0 PRECAUTIONARY STATEMENTS**

**3.1 Hazards to Humans and Domestic Animals**

Keep out of reach of children

**CAUTION**

Causes moderate eye irritation

Avoid contact with eyes, skin, or clothing

[Optional label statement, if applicable: See inside [Alternative text: See back panel] for additional] [Optional text, as applicable: Precautions [and] First Aid]

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<ul style="list-style-type: none"> <li>• Have the product container or labeling with you when calling a poison control center or doctor, or going for treatment.</li> <li>• For emergency medical treatment information, call toll-free 24 hours a day <b>1-800-334-7577</b>.</li> <li>• This product is identified as <b>[INSERT BRAND NAME], EPA Registration No. 524-539</b>.</li> </ul>	

[Optional text, if applicable: LABEL CONTINUED ON BACK]

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.



---

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes, and waterproof gloves.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

#### User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### 3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.

### 3.3 Physical or Chemical Hazards

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

---

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or on separately published supplemental labeling. Supplemental labeling for this product can be obtained from your Authorized Bayer CropScience retailer or company representative.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, socks and shoes, and waterproof gloves.

### **Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

[*Optional label text for container labels with attached labeling:* See complete Directions for Use attached to this label for complete [*Optional text, as applicable:* Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use [and] Limit of Warranty and Liability] [*Optional text, if applicable:* in English and Spanish]

## **4.0 STORAGE AND DISPOSAL**

### **STORAGE AND DISPOSAL**

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. [*Optional label text, if applicable for separate Directions for Use labeling:* See individual container label for additional storage conditions, if any.]

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** [*Optional label text, if applicable for separate Directions for Use labeling:* See label attached to the container for container handling and disposal instructions and refilling limitation.]

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID CONTAINERS OF LESS THAN 1-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill the [this] container.

[*Alternative container statement:* Nonrefillable container. Do not reuse the [this] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [this] container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse the [this] container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once properly rinsed, some plastic [*Optional text:* agricultural] pesticide containers can be taken to a collection site or picked up for recycling. [*Alternative container disposal statement:* Then offer the container for recycling, if available.]

[*Optional container disposal statement:* To find the nearest collection site, contact your chemical dealer or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 2.5-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 1 GALLON, BUT EQUAL TO OR LESS THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse the [this] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [this] container. Contact your state regulatory agency to determine allowable practices in your state.

[*Alternative container statement:* Nonrefillable container. Do not reuse or refill the [this] container.]

Triple rinse or pressure rinse (or equivalent) the [this] container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [*Optional text:* agricultural] pesticide containers can be taken to a collection site or picked up for recycling. [*Alternative container disposal statement:* Then offer the container

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

for recycling, if available.]

[*Optional container disposal statement:* To find the nearest collection site, contact your chemical dealer] [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[Insert UN packaging certification, if applicable]

[Optional packaging text: Pull at scored cutouts for easier opening]

[Optional packaging text: THIS SIDE UP] [Optional graphic of two arrows pointing upward]

[Optional packaging text: Liquid] [Optional graphic of a liquid droplet]

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 30-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill the [this] container.

[Alternative container statement: Nonrefillable container. Do not reuse the [this] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [this] container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse or pressure rinse (or equivalent) the [this] container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank. [Optional label text: For containers not equipped with pumping systems,] Fill the container ¼ full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth for 30 seconds, ensuring at least one complete revolution. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

[Alternative or additional triple rinsing instructions for large containers equipped with pumping systems: [Optional label text: For containers equipped with pumping systems,] Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Repeat this procedure two more times.]

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a collection site or picked up for recycling. [Alternative container disposal statement: Then offer the container for recycling, if available.]

[Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the [this] container, if available. If no recycling information is available on the [this] container, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)] to find the nearest recycling

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

location].]

[*Optional container disposal statement:* To find the nearest collection site, contact your chemical dealer  
[*Optional text:* or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[*Optional container disposal statement:* Return Properly Rinsed Container for Recycling – Call 1-866-99BAYER (1-866-992-2937)]

[*Optional container disposal statement:* IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389)  
www.schuetz.net/ticket; Schuetz ticket service]

[*Optional container disposal statement:* FREE IBC PICKUP] [For continental USA and Canada only]

[*Optional container disposal statement:* RETURNnet SYSTEM – To return empty IBC's Email or Call –  
www.returnnetsystem.com – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA –  
Clarkston, Michigan – USA)]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR ALL REFILLABLE CONTAINERS, EXCEPT TRANSPORT VEHICLES]

Refillable container. Refill the [this] container with pesticide only. Do not reuse the [this] container for any other purpose.

Cleaning the [this] container before refilling is the responsibility of the refiller. Cleaning the [this] container before final disposal is the responsibility of the person disposing of the container.

To clean the [this] container before final disposal, empty the remaining contents from the container into application equipment or mix-tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Then offer the container for recycling, if available.

[*Optional container disposal statement:* Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the [this] container, if available. If no recycling information is available on the [this] container, contact your chemical dealer  
[*Optional text:* or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[*Optional container disposal statement:* IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389)  
www.schuetz.net/ticket; Schuetz ticket service]

[*Optional container disposal statement:* FREE IBC PICKUP] [For continental USA and Canada only]

[*Optional container disposal statement:* RETURNnet SYSTEM – To return empty IBC's Email or Call –  
www.returnnetsystem.com – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA –  
Clarkston, Michigan – USA)]

[*Optional container disposal statement:* To obtain information about recycling refillable containers, contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937).]

[*Optional container disposal statement:* Return Properly Rinsed Container for Recycling – Call 1-866-99BAYER (1-866-992-2937)]

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

[Optional container label statements for the CUBE refillable packaging system only:

**CUBE Refillable Delivery System**

FEATURES INCLUDE:

- Automatic Venting
  - Heavy duty one-way 2-inch camloc ball valve with protective shield door
  - Complete coated steel protective enclosure
  - Durable 4-way plastic pallet
- Lift door to access one-way valve]

*[CONTAINER HANDLING AND DISPOSAL STATEMENT FOR ALL TRANSPORT VEHICLES, AS DEFINED IN 40 CFR 156.3]*

Emptied container retains vapor and product residue. Observe all precautions stated on this label until the container is cleaned, reconditioned or destroyed. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, and worn-out threads and closures. Clean thoroughly before reuse for transportation of a material of different composition or before retiring this transport vehicle from service.

*[Additional label statement for transport vehicles only: FOR BULK PESTICIDE TRANSPORT ONLY]*

*[Additional label statement for transport vehicles only: THIS LABEL FOR USE WITH TRANSPORT VEHICLES ONLY]*

*[STORAGE AND DISPOSAL FOR THE NONREFILLABLE SMART-PAK CONTAINER – A COMPOSITE PACKAGING CONSISTING OF A LIGHTWEIGHT PLASTIC CONTAINER INSIDE RIGID CARDBOARD PACKAGING. ONCE IT IS FILLED, THE LIGHTWEIGHT PLASTIC CONTAINER IS NOT STORED, TRANSPORTED OR SOLD WITHOUT THE OUTER CARDBOARD PACKAGING.]*

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. Do not store the *[this]* container unprotected from the weather, especially precipitation, for extended periods of time. **DO NOT REMOVE THE PLASTIC CONTAINER FROM THE PACKAGE UNTIL IT HAS BEEN EMPTIED AND PROPERLY RINSED.**

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the *[this]* container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not reuse or refill the *[this]* container.

Triple rinse or pressure rinse (or equivalent) the inner plastic container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the inner plastic container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into the application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing or collect rinsate for later use or disposal. Insert pressure rinsing nozzle through the bottom of the container, ensuring that you puncture the inner plastic container, and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

After rinsing, open the outer cardboard packaging and remove the inner plastic container.

Once properly rinsed, some plastic [*Optional text: agricultural*] pesticide containers can be taken to a collection site or picked up for recycling. Recycle the cardboard separately. [*Alternative container disposal statement: Then offer the outer cardboard packaging and inner plastic container for recycling, if available.*]

[*Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or call 1-866-99BAYER (1-866-992-2937)].*]

If recycling is not available, dispose of each component in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed inner plastic container and disposing both the outer cardboard packaging and inner plastic container in a sanitary landfill.

[*The inner plastic container of the Smart-Pak must be labeled with the following information*]

**[INSERT BRAND NAME]** [*Logo optional*]

ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its potassium salt..... 48.7%

OTHER INGREDIENTS: ..... 51.3%  
100.0%

\*Contains 660 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 5.5 pounds per U.S. gallon, which is equivalent to 540 grams of the acid, glyphosate, per liter or 4.5 pounds per U.S. gallon (39.8% by weight).

Keep out of reach of children

## CAUTION

EPA Reg. No. 524-539

**UNLESS THIS PLASTIC CONTAINER IS EMPTY AND HAS BEEN PROPERLY RINSED FOR DISPOSAL, IT IS NOT TO BE REMOVED FROM ITS OUTER CARDBOARD PACKAGING. FOR MORE INFORMATION, CALL 1-899-99BAYER (1-866-992-2937).**

[*Optional Smart-Pak container label statements:*

- Protect this package from precipitation until emptied and properly rinsed.
- Use the cap to cut the foil seal.
- Invert cap over foil seal and rotate clockwise to cut the foil seal. Remove foil seal.
- East to Handle
- Uses less plastic than ordinary [*Alternative text: typical*] jugs
- Produces less plastic waster than ordinary [*Alternative text: typical*] jugs
- Compact Design
- Space Efficient
- Stackable
- Easy to Use
- Easy to Pour
- Easy to Recycle]

## 5.0 PRODUCT INFORMATION

**Product Description:** This product is a postemergence, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual and perennial weeds, woody brush, trees and vines. It is formulated as a water-soluble liquid containing surfactant and may be applied using standard and specialized pesticide application equipment after dilution and thorough mixing with water or other carrier according to label directions.

[*Optional label text:* Do not add [*Optional label text:* surfactants, additives containing surfactants,] buffering agents or pH adjusting agents to the spray solution when [INSERT BRAND NAME] is the only pesticide being applied unless otherwise directed. See the "MIXING" section of this label for instructions regarding other additives.]

[*Optional label text:* No additional surfactant is needed in the spray solution. This includes additives containing surfactants, buffering agents or pH adjusting agents when [INSERT BRAND NAME] is the only pesticide used unless otherwise directed.]

**Mechanism of Action:** Glyphosate works by targeting an enzyme that is essential for plant growth.

**No Soil Activity:** This product binds tightly to soil particles and does not provide residual weed control. Weeds must be emerged at the time of application to be controlled by foliar application of this product. Weed seeds in the soil will not be affected by this product and will continue to germinate. Unattached plant rhizomes and rootstocks beneath the soil surface will also not be affected by this product.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.

**Stage of Weeds:** Annual weeds are easiest to control when they are small. Performance of this product on most perennial weeds is best when applied at late growth stages approaching maturity. Refer to the "ANNUAL WEEDS RATE SECTION," "PERENNIAL WEEDS RATE SECTION" and "WOODY BRUSH, TREES AND VINES RATE SECTION" for more information on the control of specific weeds.

**Cultural Considerations:** Reduced weed control could result when this product is applied to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to re-grow prior to application. Always use a higher product application rate within the given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area. Reduced weed control could also result when this product is applied to weeds that show signs of disease or insect damage, are covered with dust, or are surviving under poor growing conditions.

**Spray Coverage:** For best results with this product, spray coverage must be uniform and complete. Do not spray foliage to the point of runoff.

**Rainfastness:** Rainfall within 4 hours of application could wash this product off of the foliage and a second application might then be needed to achieve acceptable weed control. Refer to specific use sections of this label for additional information on the minimum intervals required before re-application of this product.

**Time to Symptoms:** This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of aboveground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds, effects might not be visible for 7 or more days after application. Extremely cool or cloudy weather following application could slow activity of this product and delay development of visual symptoms.

**Maximum Application Rates:** The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowable application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or in a tank mixture, on a basis of total pounds of



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalents) does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Unless otherwise specified on this label, the combined total application of this product on a site must not exceed 5.3 quarts (6 pounds of glyphosate acid) per acre per year. For applications on non-crop sites, or on tree, vine or shrub crop production sites, the combined total application of this product must not exceed 7 quarts (8 pounds of glyphosate acid) per acre per year.

Unless otherwise specified on this label in the sections that follow, the maximum single application rate of this product is 3.3 quarts (3.7 pounds of glyphosate acid) per acre when applied using ground application equipment and 44 fluid ounces (1.55 pounds of glyphosate acid) per acre when applied using aerial application equipment. Unless otherwise specified on this label, multiple applications of this product made be made onto a site up to the maximum annual application rates defined above. Allow a minimum of 14 days between applications.

The following table provides the glyphosate application rate (pounds of glyphosate acid equivalents per acre) when this product is applied at the application rates indicated (fluid ounces or quarts of this product per acre).

<b>Application Rate of [INSERT BRAND NAME] (amount of product per acre)</b>	<b>Application Rate of Glyphosate Acid Equivalents (ae) (pounds of ae per acre)</b>
11 fluid ounces	0.39
16 fluid ounces	0.56
22 fluid ounces	0.77
32 fluid ounces	1.125
44 fluid ounces	1.55
64 fluid ounces	2.25
3.3 quarts	3.7
4.1 quarts	4.6
5.3 quarts	6
7 quarts	8

To determine the Glyphosate acid equivalents (pounds of Glyphosate ae per acre) for application rates of [INSERT BRAND NAME] not listed here, multiply the application rate (fluid ounces per acre) by 0.0352.

$$\text{Application Rate} \times 0.0352 = \text{Glyphosate (ae) per acre}$$

### 6.0 WEED RESISTANCE MANAGEMENT

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mechanism of action classification system of the Weed Science Society of America. Any weed population can contain plants that are naturally resistant to Group 9 herbicides. Weeds resistant to Group 9 herbicides can be effectively managed by using another herbicide from a different Group (either alone or in a mixture according to label directions), by using other cultural or mechanical methods of weed control, or a combination of the two. Consult your local company representative, state cooperative extension agent, professional consultant or other qualified authority to determine appropriate actions for controlling specific resistant weeds.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### 6.1 Weed Management Practices

Resistant populations arise when rare individual plants are uncontrolled by a normal dose of a given herbicide under normal environmental conditions. In the absence of other control measures these individuals survive, produce seed, and eventually become the dominant biotype in the field through continuous selection. The best means of reducing this selection is to use diverse weed control practices such as multiple herbicides with different mechanisms of action, and often in combination with various mechanical and cultural practices.

Suspected herbicide resistance can be identified by these factors:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially when control is achieved on adjacent weeds
- A spreading patch of non-controlled plants of a particular species
- Surviving plants mixed with controlled individuals of the same species

To minimize the occurrence of herbicide-resistant biotypes, including those resistant to glyphosate, implement the following weed management practice options that are practical to your situation. These management practices are applicable to reduce the spread of confirmed resistant biotypes (managing existing resistant biotypes) and to reduce the potential for selecting for resistance in new species (proactive resistance management).

- Use a diversified approach toward weed management focused on preventing weed seed production and reducing the number of weed seeds in the soil.
- Plant crops into fields that are as weed-free as possible and then keep them as weed-free as possible.
- Plant crop seed that is as weed-free as possible.
- Scout fields routinely, before and after herbicide application.
- Use multiple herbicide mechanisms of action that are effective against the most troublesome weeds in your field and against those with known resistance.
- Apply herbicides at application rates listed on the label when weeds are within the size range indicated on the label.
- Emphasize cultural practices that suppress weeds by using crop competitiveness.
- Use mechanical and biological weed management practices, where appropriate.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules.
- Manage weed seed at harvest and after harvest to prevent a buildup of the weed seedbank.

### 6.2 Management of Glyphosate-Resistant Biotypes

Appropriate testing is needed to determine if a weed is resistant to glyphosate. Call 1-866-99BAYER (1-866-992-2937) or contact your Bayer CropScience representative to report any incidence of non-performance of this product against a particular weed species. To determine if resistance in any particular weed biotype has been confirmed in your area, or for additional information on glyphosate-resistant biotypes, go to [www.weedscience.org](http://www.weedscience.org).

Glyphosate-resistant weeds can be controlled or managed by applying this product in combination with residual preemergence herbicides and/or other postemergence herbicides labeled for control of the targeted weed in the crop being grown. For more information, see the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Since the occurrence of resistant weeds is difficult to detect prior to use, to the extent consistent with applicable law, Bayer CropScience accepts no liability for any losses that result from the failure of this product to control resistant weeds.

### 7.0 MIXING

Spray solutions of this product may be mixed, stored and applied using clean stainless steel, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.**

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

A 50-mesh nozzle screen or line strainer on the spray equipment is adequate.

Clean sprayer parts promptly after using this product by thoroughly flushing with water.

#### 7.1 Mixing with Water

PERFORMANCE OF THIS PRODUCT CAN BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS OR DITCHES THAT IS VISIBLY MUDDY OR MURKY.

This product mixes readily with water. Mix spray solutions of this product as follows. Begin filling the mixing tank or spray tank with clean water. Add the required amount of this product near the end of the filling process and mix gently. Foaming of the spray solution can occur during mixing. To prevent or minimize foaming, mix gently, terminate bypass and return lines at the bottom of the tank, and, if necessary, add an appropriate anti-foam or defoaming agent to the spray solution.

#### 7.2 Tank Mixtures

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control in the soil, a broader weed control spectrum, or an alternate mechanism of action.

Some tank-mix products have the potential to cause crop injury under certain conditions, at certain growth stages, and/or under other circumstances. Read the label of all products to be used in the tank mixture prior to use to determine the potential for crop injury.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers could result in reduced weed control or crop injury. Bayer CropScience has not tested all tank-mix product formulations for compatibility, antagonism or reduction in product performance. To the extent consistent with applicable law, buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label, or on separate supplemental labeling or Fact Sheets published for this product.

When a tank-mix with a generic active ingredient, such as 2,4-D, atrazine, dicamba, diuron, pendimethalin, or any other product or material, is listed on this label, it is the responsibility of the pesticide user to ensure that the intended use is included on the label of each product added to the mix.

Refer to all individual product labels, supplemental labeling and Fact Sheets for all products in the tank mixture, and observe all precautions and limitations on the label, including any application timing restrictions, soil restrictions, minimum re-cropping intervals and/or crop rotation restrictions. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

For optimal overall weed control, apply tank mixtures with this product at a minimum spray volume rate of 10 gallons per acre.

### 7.3 Tank-Mixing Procedure

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Mix only the quantity of spray solution that will be applied that day. Application of tank-mix solutions that stand overnight could result in reduced weed control.

Prepare tank mixtures of this product as follows:

1. Place a 20 to 35-mesh screen or wetting basket over the filling port of the tank.
2. Through the screen, fill the tank one-half full with water and start gentle agitation.
3. If ammonium sulfate is to be used, add it slowly through the screen into the tank and continue adding water into the tank through the screen. If dry ammonium sulfate is being used, ensure that it is completely dissolved in the tank before adding other products.
4. If a wettable powder is used, first prepare a slurry of it with water and add it SLOWLY through the screen into the tank while continuing gentle agitation.
5. If a flowable formulation is to be added, premix one part flowable with one part water and add the diluted mixture SLOWLY through the screen into the tank while continuing gentle agitation.
6. If an emulsifiable concentrate formulation is to be added, premix one part emulsifiable concentrate with two parts water and add the diluted mixture SLOWLY through the screen into the tank while continuing gentle agitation.
7. Continue filling the tank with water through the screen and add the required amount of this product near the end of the filling process.
8. [*Optional label statement:* If a nonionic surfactant is to be added, add it to the tank before completing the filling process.]
9. Add individual tank-mix components to the tank in the following order: wettable powders, flowables, emulsifiable concentrates, drift reduction additives, water soluble liquids (this product) [*Optional text:* , surfactant].

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, agitate thoroughly to re-suspend the mixture before resuming application.

Keep by-pass and return lines on or near the bottom of the tank to minimize foaming.

A 50-mesh nozzle screen or line strainer on the spray equipment is adequate.

### 7.4 Mixing Spray Solution Concentrations

All reference throughout this label to concentration of this product in a spray solution is on a percentage-of-volume basis.

Prepare the desired volume of spray solution at a given concentration by mixing the amount of this product indicated in the following table with water.

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Desired Volume of Spray Solution	Amount of [INSERT BRAND NAME] to Achieve Indicated Concentration in Spray Volume (percent by volume)					
	0.4%	0.7%	1%	1.5%	4%	7%
1 gallon	0.5 fl oz	1 fl oz	1.3 fl oz	2 fl oz	5 fl oz	9 fl oz
25 gallons	13 fl oz	22 fl oz	1 qt	1.5 qts	4 qts	7 qts
100 gallons	1.6 qts	2.8 qts	1 gal	1.5 gals	4 gals	7 gals

2 tablespoons = 1 fluid ounce (fl oz)

For filling backpack and pump-up sprayers, consider mixing the appropriate amount of this product with water in a larger container and then filling the sprayer from the larger container.

**7.5 Surfactants** [*this section is optional in the final printed label*]

Although not always required, nonionic surfactants that are labeled for use with herbicides may be added to spray solutions of this product. Do not reduce application rates or concentration of this product when applying in spray solutions containing additional surfactant.

[*Optional text: Additional surfactant can increase the performance of this product at water carrier volumes above 30 gallons per acre or at application rates below 16 fluid ounces of product per acre.*]

[*Optional text: Use a surfactant concentration of 0.25 to 0.5 percent (1 to 2 quarts per 100 gallons of spray solution) when adding surfactant that contains at least 70 percent active ingredient, or a 1-percent surfactant concentration (4 quarts per 100 gallons of spray solution) when adding surfactant that contains less than 70 percent active ingredient.*]

Read and follow the directions for use and observe all precautionary statements and other information on the surfactant label.

[*Optional text: DO NOT add buffering agents or pH adjusting agents to the spray solution when [INSERT BRAND NAME] is the only pesticide product being applied.*]

[*Optional text only if use in glyphosate-tolerant crops is included on the final printed labeling: DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR PREHARVEST APPLICATION TO COTTON OR ANY POSTEMERGENCE (IN-CROP) APPLICATION TO COTTON WITH ROUNDUP READY, ROUNDUP READY FLEX, XTENDFLEX, OR ANY OTHER GLYPHOSATE-TOLERANT COTTON TECHNOLOGY.*]

**7.6 Ammonium Sulfate**

Unless otherwise directed, the addition of 1 to 2 percent dry ammonium sulfate by weight (8.5 to 17 pounds per 100 gallons of water), could increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank-mixed with certain residual herbicides. An equivalent amount of a liquid formulation of ammonium sulfate may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water promptly after use to reduce corrosion.

When using ammonium sulfate, apply this product at rates or spray solution concentrations as directed on this label; lower application rate or concentration could result in reduced weed control.

**7.7 Colorants and Dyes**

Colorants and marking dyes may be added to spray solutions of this product; however, they can reduce the performance of this product. Use colorants and dyes according to the manufacturer's directions.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### 7.8 Drift Reduction Additives

Drift reduction additives may be used with all application equipment types, except wiper applicators, sponge bars and controlled droplet applicators (CDA). However, use of drift reduction additives can affect spray coverage, which could reduce the performance of this product. When a drift reduction additive is used, read and follow all directions for use, precautions, limitations and other information on the product label.

### 8.0 APPLICATION EQUIPMENT AND TECHNIQUES

This product may be applied using the following equipment:

**Aerial Application Equipment**—fixed-wing and helicopter

**Ground Application Equipment**—boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast application equipment

**Handheld Sprayers**—backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other handheld and motorized spray equipment used to direct the spray onto undesirable foliage

\* This product is not registered in California or Arizona for use in mistblowers.

**Selective Application Equipment**—shielded and hooded sprayers, wiper applicator, sponge bar

**Injection Systems**—aerial or ground injection sprayers

**Controlled Droplet Applicator (CDA)**—handheld or boom-mounted applicators that produce a spray pattern consisting of a narrow range of droplet sizes

APPLY THIS PRODUCT USING PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF ACCURATELY DELIVERING DESIRED VOLUMES.

Do not apply this product through any type of irrigation system.

### 8.1 Spray Drift Management

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [*Optional text:* GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, [*Optional text, if applicable:* EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® AND OTHER LISTED GLYPHOSATE-TOLERANT CROPS,] AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation, as small quantities of this product can cause severe damage or destruction to the crop, plants or other vegetation on which application was not intended.

AVOID DRIFT. USE EXTREME CARE TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHEN APPLYING THIS PRODUCT.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and grower are responsible for considering all these factors when making decisions regarding the application of this product.

The likelihood of injury occurring as the result of spray drift while applying this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or generation of fine particles (mist) that are likely to drift.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFERS MUST BE MAINTAINED.

AVOID APPLYING THIS PRODUCT AT EXCESSIVE SPEED OR SPRAYER PRESSURE.

### **8.2 Aerial Application Equipment**

Unless otherwise prohibited, all applications of this product described on this label may be made using aerial application equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label or on separate supplemental labeling published for this product.

DO NOT APPLY THIS PRODUCT USING AERIAL APPLICATION EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED ON THIS LABEL OR ON SEPARATELY PUBLISHED SUPPLEMENTAL LABELING FOR THIS PRODUCT.

FOR SPECIFIC USE INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS RELATED TO THE AERIAL APPLICATION OF THIS PRODUCT IN ARKANSAS AND CALIFORNIA, OR SPECIFIC COUNTIES THEREIN, REFER TO THE LIMITATIONS ON AERIAL APPLICATION IN THAT STATE OR COUNTY PRESENTED IN THIS SECTION.

Unless otherwise directed, the maximum single application rate of this product is 44 fluid ounces per acre when using aerial application equipment. Apply this product at the appropriate rate in 3 to 15 gallons of water per acre unless otherwise directed on this label or on separate supplemental labeling for this product. Refer to the individual use sections of this label for application rates, spray volumes and additional directions for use.

Drift control reduction additives may be used.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

#### **Aircraft Maintenance**

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES COULD RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) that meets aerospace specification MIL-C-38413 can help prevent corrosion.

#### **AERIAL SPRAY DRIFT MANAGEMENT**

The following drift management requirements must be followed to minimize off-target drift movement during aerial application. [*Optional label text, if applicable:* These requirements do not apply to forestry applications.]

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be followed.

#### **Importance of Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly or under unfavorable environmental conditions, such as in windy, high temperature with low humidity, and/or inversion conditions as described below.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Operate sprayer at a pressure towards the lower end of the range listed for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing the pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length could further reduce drift without reducing swath width.
- **Application height:** Application must be made at a height of 10 feet or less above the top of the largest plants unless a greater height is required for aircraft safety. Making the application at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

### Swath Adjustment

When an application is made with a crosswind present, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase the swath adjustment distance with increasing drift potential (higher wind, smaller droplets, etc.).

### Wind

Drift potential is lowest at wind speeds of between 2 and 10 miles per hour. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Avoid application when wind speeds are below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making an application in low relative humidity, set application equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversion

Do not apply this product during a temperature inversion as drift potential is high under these conditions. Temperature inversions restrict vertical air mixing, which causes small droplets to remain suspended in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from a sensitive area).

Avoid direct application to any body of water.

**State Specific Limitations on Aerial Application** [*The following section is optional to be used only if required by the States indicated.*]

---

### LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY

---

DO NOT apply this product using aerial application equipment in residential areas.

AVOID DRIFT – DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT OF THIS PRODUCT ONTO ANY VEGETATION TO WHICH APPLICATION WAS NOT INTENDED CAN CAUSE DAMAGE. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, USE PROPER AERIAL APPLICATION EQUIPMENT FITTED WITH APPROPRIATE NOZZLES AND MAINTAIN ADEQUATE BUFFERS.

Follow the directions below when making an aerial application near non-target crops, desirable annual vegetation, or desirable perennial vegetation after bud break and before total leaf drop.

1. Do not apply this product within 100 feet of all desirable vegetation or non-target crops.
2. If winds are blowing up to 5 miles per hour TOWARD desirable vegetation or non-target crops, do not apply this product within 500 feet of the desirable vegetation or crops.
3. If winds are blowing between 5 and 10 miles per hour TOWARD desirable vegetation or non-target crops, a buffer zone greater than 500 feet might be needed to protect the desirable vegetation or crops.
4. Do not apply this product using aerial application equipment when winds are blowing in excess of 10 miles per hour.
5. Do not apply this product using aerial application equipment when inversion conditions exist.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be applied in California using aerial application equipment. Tank mixtures of this product with 2,4-D amine formulations may be applied by air in California in fallow fields and in reduced tillage systems, and for alfalfa and pasture renovation applications only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

---

### ADDITIONAL LIMITATIONS ON AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

---

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

North: Fresno County line  
South: Fresno County line  
East: State Highway 99  
West: Fresno County line

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

### **Written Directions**

Written directions MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. These written directions MUST state the proximity of surrounding crops and that conditions of each manufacturer's product label and this label have been satisfied.

### **Aerial Applicator Training and Equipment**

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to ensure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

**Application at Night** – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

For additional information on the proper aerial application of this product in Fresno County, call 1-866-99BAYER (1-866-992-2937).

---

### **LIMITATIONS ON AERIAL APPLICATION IN ARKANSAS ONLY**

---

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Apply this product at the appropriate rate in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range have a lower drift potential.

Applications are typically to be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when winds are in excess of 10 miles per hour.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions can occur when wind speeds are less than 2 miles per hour.

Follow the directions below when an aerial application is made near non-target crops or other desirable vegetation:

1. Do not apply this product within 100 feet of non-target crops or any desirable vegetation.
2. If winds are blowing up to 5 miles per hour TOWARD non-target crops or desirable vegetation, do not apply this product within 500 feet upwind of the desirable vegetation or crop.
3. If winds are blowing between 5 and 10 miles per hour TOWARD non-target crops or desirable vegetation, a buffer zone greater than 500 feet might be needed to protect the crop or desirable vegetation.

---

### 8.3 Ground Application Equipment

Apply this product at an appropriate rate specified on this label in 3 to 40 gallons of water per acre when making a broadcast application using ground application equipment, unless otherwise directed on this label or on separate supplemental labeling or Fact Sheets published for this product. As the weed density increases, increase the spray volume towards the upper end of this range to ensure complete coverage. Use nozzles that will avoid generating a fine mist. For best performance of this product when using ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

### 8.4 Handheld Sprayers

When using a handheld sprayer, apply spray solutions of this product uniformly and completely to the foliage of target weeds using a coarse droplet spectrum and a spray-to-wet technique; do not spray to the point of runoff. For the appropriate concentration of this product in the spray solution and timing of application to control certain weeds, woody brush, trees and vines, refer to the "ANNUAL WEEDS RATE SECTION," "PERENNIAL WEEDS RATE SECTION" and "WOODY BRUSH, TREES AND VINES RATE SECTION" of this label.

Spot application of this product for weed control in a cropping system using a handheld sprayer may be made only when specifically directed on this label or on separate supplemental labeling for this product. The crop sprayed with this product will be killed along with the weeds. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.

### 8.5 Selective Application Equipment

Selective application equipment allows this product to be applied to weeds growing near a crop or other desirable vegetation without killing the desirable vegetation. Selective application equipment must be capable of preventing all contact of the herbicide solution with the crop or other desirable vegetation and operated without spray mist escape, leakage or dripping of the herbicide solution.

AVOID CONTACT OF THIS HERBICIDE WITH DESIRABLE VEGETATION. Contact of this product with desirable vegetation could result in unwanted plant damage or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

### Shielded and Hooded Sprayers

A shielded sprayer directs the herbicide solution to the target weeds while using an impervious material, or shield, to protect nearby desirable vegetation from coming into contact with the herbicide spray. To provide maximum protection for desirable vegetation, keep shield properly adjusted and use spray nozzles that provide uniform coverage within the application area.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

A hooded sprayer is a type of shielded sprayer where the spray pattern is fully enclosed, including the top, sides, front and back, thereby shielding the crop or other desirable vegetation from the spray solution.

This product may be diluted in water and applied using a shielded or hooded sprayer to weeds listed on this label growing on any terrestrial non-crop site described on this label and in between rows of plants (row middles) in any cropping system listed on this label.

Properly adjust the hood to protect desirable vegetation. Ensure that the hood is capable of completely enclosing the spray pattern. If necessary when applying around crops grown on raised beds, extend the front and rear flaps of the hooded sprayer downward to reach the ground in deep furrows

A hooded sprayer must be configured and operated in a manner that minimizes bouncing and avoids raising the hood up off the ground surface at any time. If the hood is raised, spray particles can escape and come into contact with the crop, causing damage to or destruction of the crop or other desirable vegetation. Avoid operating this equipment on rough or sloping terrain where the spray hood is likely to rise up off the ground surface.

Use hoods designed to minimize excessive dripping or runoff down the inside of the hood, such as a single, low pressure, low-drift, flat-fan nozzle with an 80 to 95-degree spray angle positioned at the top center of the hood, with a spray volume of 20 to 30 gallons per acre.

The following procedures will help reduce the potential for crop injury when using a hooded sprayer:

- Operate the sprayer with the hood on the ground or skimming across the ground surface.
- Leave at least an 8-inch untreated strip over the drill row. (For example, if the crop row width is 38 inches, make the maximum width of the spray hood 30 inches.)
- Operate at a ground speed of no greater than 5 miles per hour to minimize bouncing of the hooded sprayer.
- Apply when wind speed is 10 miles per hour or less.
- Use low-drift nozzles that will provide uniform coverage within the application area.

Injury to a crop or other desirable vegetation can occur when application is made to foliage of weeds that come into direct contact with the crop or desirable vegetation. Do not apply this product when leaves of desirable vegetation are growing in direct contact with weeds. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting or destruction.

### **Wiper Applicator**

A wiper applicator is a device that physically wipes this product or solutions of this product directly onto the target weed or cut stump. Any handheld device that is capable of physically wiping this product or solutions of this product directly onto the target weed or cut stump, such as a paint brush, may be used.

Wiper applicators may be used over the top of food or feed crops **ONLY** if specifically permitted for use over that crop by this label or by separately published supplemental labeling for this product.

A mechanical wiper applicator, such as a rope wick or sponge bar that can be driven through a field over the top of a crop or other desirable vegetation to control tall weeds growing above the desirable vegetation, must be designed, maintained and operated to prevent the herbicide solution from coming into contact with desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. To protect the crop or other desirable vegetation, adjust the height of the applicator to ensure that the wiper contact point is a minimum of 2 inches above the desirable vegetation.

Weeds that do not come into contact with the herbicide solution will not be affected. The more weed foliage exposed above the desirable vegetation the better mechanical wiper applicators work. Better results can

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

also occur when weeds are a minimum of 6 inches above the desirable vegetation. Poor contact can also occur when weeds are growing in dense clumps, when operating in an area of severe weed infestation, or when weed height varies dramatically. In these situations, more than one application of this product might be necessary.

For optimal results, operate wiper applicators at a ground speed of no greater than 5 miles per hour. Performance in areas of heavy weed infestation can be improved by reducing speed, which will provide more time for re-saturation of the wiper with the herbicide solution and more contact time of the wiper with the weed. Better weed control using a wiper applicator can be obtained when two applications are made travelling in opposite directions in the field.

Keep wiper surfaces clean.

Be aware that on sloping ground the herbicide solution can migrate to one side, causing dripping on the lower end and drying of the wiper on the upper end of the applicator.

Do not apply this product using a wiper applicator when weeds are wet.

Do not add surfactant to the herbicide solution when using a wiper applicator.

With **Rope Wick and Sponge Wick Applicators**, apply solutions ranging from 33 to 75 percent of this product by volume in water.

With **Panel Applicators**, apply solutions ranging from 33 to 100 percent (undiluted) of this product by volume in water.

Mix only the amount of this product that will be used during a 1-day period, as reduced product performance can result from the use of solutions held in storage.

Clean wiper parts promptly after using this product by thoroughly flushing with water.

### 8.6 Injection Systems

This product may be used in aerial and ground injection spray systems as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this concentrated product with the undiluted concentrate of other products for use in injection systems, unless otherwise directed.

### 8.7 Controlled Droplet Applicator (CDA)

The amount of this product applied per acre using a controlled droplet applicator (CDA) must be no less than the rate specified on this label for application using conventional broadcast application equipment.

A controlled droplet applicator produces a spray pattern that is not easily visible. Use extreme care to avoid spray or drift from contacting the foliage or any other green tissue of desirable vegetation, as plant damage or destruction could result.

## 9.0 ANNUAL AND PERENNIAL CROPS

THIS SECTION PROVIDES DIRECTIONS FOR USE OF THIS PRODUCT THAT APPLY TO ALL CROPS LISTED IN THE FOLLOWING SECTIONS. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC USE INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

*[Optional label text to be used only if Roundup Ready crops are included on the final printed label: See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label, or separately published supplemental labeling for this product, for directions for use in Roundup Ready and other glyphosate-tolerant crops listed on this label.]*

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**TYPES OF APPLICATION:** Chemical Fallow; Preplant Fallow Beds; Preplant; At-Planting; Preemergence; Hooded Sprayer in Row Middles; Shielded Sprayer in Row Middles; Wiper Applicator in Row Middles; Post-Harvest

**USE INSTRUCTIONS:** This product may be applied during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergence to annual and perennial crops listed on this label, except where specifically limited. For any crop not listed on this label, application must be made a minimum of 30 days prior to planting. Unless otherwise directed, apply this product according to the rates listed in the “ANNUAL WEEDS RATE SECTION,” “PERENNIAL WEEDS RATE SECTION” and “WOODY BRUSH, TREES AND VINES RATE SECTION” of this label. Application rates specified on this label for hard-to-control weeds, or those specified on separate supplemental labeling for this product, supersede the rates in the “ANNUAL WEEDS RATE SECTION,” “PERENNIAL WEEDS RATE SECTION” and “WOODY BRUSH, TREES AND VINES RATE SECTION” of this label. Additional information on hard-to-control weeds can be found on Fact Sheets published for this product.

Application of this product may be repeated as needed up to a maximum of 5.3 quarts per acre per year. Refer to specific use sections of this label for additional information on minimum intervals required before re-application of this product.

Hooded and shielded sprayers and wiper applicators capable of preventing all contact of the herbicide solution with the crop may be used in mulched or unmulched row middles after crop establishment. Wiper applicators may be used over the top of crops to control tall weeds only when specifically directed in the individual crop sections that follow. Crop injury is possible with these methods of application. Refer to the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for information regarding the potential for crop injury using selective application equipment.

Spot application of this product for weed control in a cropping system may be made only when specifically directed in the individual crop sections that follow.

Unless otherwise prohibited, all applications of this product described in the sections that follow may be made using aerial application equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label and on all supplemental labeling published for this product. Refer to the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for information on aerial application and procedures for avoiding spray drift that could cause injury to any vegetation not intended for application. Use of appropriate buffers will help prevent injury to adjacent vegetation.

**TANK MIXTURES:** This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mechanism of action. Always read and follow label directions for all products in the tank mixture. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Use all products according to rates and timing specified on the label. Some tank-mix products have the potential to cause crop injury. Read the label for all products in the tank mixture prior to use to determine the potential for crop injury. Always predetermine the compatibility of tank-mix products together in the carrier by mixing small proportional quantities in advance. Mixing other products with this herbicide in the spray tank can cause incompatibility, antagonism, or a reduction in the efficacy of this product. Bayer CropScience has not tested all product formulations for compatibility or performance in a tank-mix with this product. To the extent consistent with applicable law, buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not specifically identified on this label or on separate supplemental labeling or Fact Sheets for this product. See the “MIXING” section of this label for more information on tank mixtures.

**PRECAUTIONS:** Avoid contact of this herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops, as severe crop injury or destruction could result. Transplant seedlings coming into contact with weeds that are still wet with a spray solution of this product could result in significant crop injury. When making preemergence applications, application must

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

be made before crop emergence to avoid severe crop injury. Broadcast application of this product at emergence will result in injury or death of emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops where spot application is allowed, the crop sprayed with this product will be killed along with the weeds. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction. See the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for additional information.

Preharvest application on crops grown for seed could result in a reduction in germination or vigor. To the extent consistent with applicable law, buyer and all users are responsible for such loss or damage in connection with the preharvest use of this product on any crop grown for seed.

**RESTRICTIONS:** Before using this product, read and follow the Maximum Application Rate information in the “PRODUCT INFORMATION” section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined with the use of any and all other herbicides containing glyphosate as the active ingredient, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate.

Unless otherwise directed on this label, application using selective equipment, including wiper applicators and hooded and shielded sprayers, must be made a minimum of 14 days prior to harvest. Post-harvest and fallow applications must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

Do not harvest or feed vegetation from an area for 8 weeks following broadcast postemergence application, unless otherwise directed.

In crops where spot application is allowed, do not apply this product to more than 10 percent of the total field to be harvested, unless otherwise directed.

When applying this product as a tank mixture with one or more products, refer to each individual tank-mix product label for restrictions and apply the mixture in accordance with the most restrictive statements for each product in the tank.

### **9.1 Cereal and Grain Crops**

**LABELED CROPS:** Barley; Buckwheat; Millet (pearl, proso); Oats; Rice; Rye; Quinoa; Teff; Teosinte; Triticale; Wheat (all types); Wild Rice

**TYPES OF APPLICATION:** Those listed in Section 9.0, plus Red Rice Control Prior to Planting Rice; Spot Application (except rice); Control of Barnyardgrass in Rice Using Renovation Treatment (California only); Wiper Applicator (wheat and feed barley only); Preharvest (wheat and feed barley)

#### **Preplant, At-Planting, Preemergence**

**USE INSTRUCTIONS:** This product may be applied before, during or after the planting of cereal crops, but prior to crop emergence.

#### **Red Rice Control Prior to Planting Rice**

**USE INSTRUCTIONS:** Flush fields prior to application to obtain uniform germination and stand of red rice and then apply 32 fluid ounces of this product in 5 to 10 gallons of water per acre when the majority of the red rice plants are at the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves might only be partially controlled. Avoid spraying during conditions of low humidity, as reduced control of red rice could result.

**RESTRICTIONS:** Do not apply this product to rice fields or levees when fields contain floodwater. Do not flood fields for a minimum of 8 days following application.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### **Spot Application (Except Rice)**

USE INSTRUCTIONS: This product may be applied as a spot application in cereal crops, except rice. Apply before heading in small grains.

RESTRICTIONS: Do not apply this product to more than 10 percent of the total field area to be harvested.

### **Control of Barnyardgrass in Rice Using Renovation Treatment (California Only)**

THIS APPLICATION FOR USE IN CALIFORNIA ONLY

USE INSTRUCTIONS: This product may be applied as a renovation treatment in rice crops to control barnyardgrass (*Echinochola crus-galli*) infestations using ground broadcast application equipment or a handheld sprayer. Renovation is defined as an herbicide application that will result in crop and weed destruction in an entire field or contiguous area within a field.

RESTRICTIONS: Rice straw and stubble from the application area plus an additional 25 feet on all sides of the area may not be used for animal bedding, grazing, or any other feed purpose. DO NOT make this application using aerial application equipment.

### **Wiper Applicator (Wheat and Feed Barley Only)**

USE INSTRUCTIONS: This product may be applied over the top of wheat and feed barley using a wiper applicator to control tall weeds. To control common rye or cereal rye, apply after weeds have headed and achieved maximum growth. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow a minimum of 35 days between application and harvest. Do not use roller applicator.

### **Preharvest (Wheat and Feed Barley Only)**

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of wheat and feed barley. Apply up to 22 fluid ounces this product in 10 to 20 gallons of water per acre when using ground application equipment and in 3 to 10 gallons of water per acre when using aerial application equipment. For wheat, apply after the hard-dough stage when grain moisture is 30 percent or less. Stubble may be grazed immediately after harvest. For feed barley, make application after the hard-dough stage when grain moisture is 20 percent or less.

RESTRICTIONS: Do not apply more than 22 fluid ounces of this product per acre for preharvest application. Allow a minimum of 7 days between application and harvest or grazing.

### **Post-Harvest**

USE INSTRUCTIONS: This product may be applied up to 3.3 quarts per acre as a single ground broadcast application for weed control after harvest of cereal crops. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for weed control following harvest of cereal crops. Read and follow label directions for all products in the tank mixture.

RESTRICTIONS: Do not exceed a total combined application rate of 5.3 quarts of this product per acre per year. Allow a minimum of 7 days between application and harvest or feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

## **9.2 Corn**

TYPES OF CORN: Field corn; Popcorn; Seed corn; Silage corn; Sweet corn

TYPES OF APPLICATION: Those listed in Section 9.0, plus Spot Application; Preharvest



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

For directions for use with field corn hybrids with Roundup Ready 2 Technology (including Roundup Ready Corn 2 and field corn products displaying the Roundup Ready 2 Technology logo), or with sweet corn hybrids with Roundup Ready 2 Technology (including Roundup Ready Sweet Corn and sweet corn products displaying the Roundup Ready 2 Technology logo), and field corn hybrids with Agrisure GT Technology, see the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank-mix before, during or after planting corn, but prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*: , or one or more of the products listed]. Ensure that the product used is labeled for application prior to the planting or the emergence of the type of corn being grown. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; dimethenamid; dimethenamid-P; flufenacet; flumetsulam; flumiclorac pentyl ester; fluthiacet-methyl; isoxaflutole; linuron; mesotrione; metolachlor; s-metolachlor; metribuzin; pendimethalin; rimsulfuron; saflufenacil; simazine; thiencazone-methyl

**Axiom DF** (EPA Reg. No. 264-766; *metribuzin, flufenacet*); **Balance Flexx** (EPA Reg. No. 264-1067; *isoxaflutole*); **Capreno** (EPA Reg. No. 264-1063; *tembotrione, thiencazone-methyl*); **Corvus** (EPA Reg. No. 264-1066; *isoxaflutole, thiencazone-methyl*); **Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **DiFlexx** (EPA Reg. No. 264-1173; *dicamba*); **DiFlexx DUO** (EPA Reg. No. 264-1184; *dicamba, tembotrione*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*); **Harness MAX** (EPA Reg. No. 524-636; *acetochlor, mesotrione*); **TripleFLEX II** (EPA Reg. No. 524-614; *acetochlor, clopyralid, flumetsulam*)

AAtrex 4L; AAtrex Nine-O; Acuron; Acuron Flexi; Aim EC; Aim EW; Anthem; Anthem ATZ; Anthem Flex; Anthem Maxx; Axiom DF; Balance Flexx; Banvel; Banvel 480; Bicep II MAGNUM; Bicep II MAGNUM FC; Bicep Lite II MAGNUM; Callisto; Capreno; Cinch; Cinch ATZ; Cinch ATZ Lite; Clarity; Corvus; Degree Xtra; DiFlexx; DiFlexx DUO; Distinct; Dual MAGNUM; Dual II MAGNUM; FulTime; FulTime NXT; Guardsman MAX; Harness; Harness MAX; Harness Xtra; Harness Xtra 5.6L; Hornet WDG Broadleaf Blend; Keystone; Keystone LA; Keystone LA NXT; Keystone NXT; Leadoff; Linex 4L; Lorox DF; Marksman; Me-Too-Lachlor II; Outlook; Prowl 3.3 EC; Prowl H2O; Python WDG; Resicore; Resolve DF; Resolve Q; Resolve SG; Resource; Shark EW; Shark H2O; Sharpen Powered by Kixor; Stalwart; Stalwart C; Stalwart Xtra; Surpass EC; Surpass NXT; TopNotch; TripleFLEX II; Verdict powered by Kixor; Zidua]

---

For hard-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply 22 fluid ounces of this product per acre in these tank mixtures. For other annual weeds listed on this label, apply

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

16 to 22 fluid ounces of this product per acre when weeds are less than 6 inches tall and 22 to 32 fluid ounces per acre when weeds are over 6 inches tall. When using a nitrogen solution as the carrier, higher application rates might be needed for acceptable weed control.

**RESTRICTIONS:** Application of 2,4-D or dicamba must be made a minimum of 7 days prior to planting corn.

In Southern states, do not mix this product in nitrogen solutions for application to hard-to-control grasses, such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. This area includes Illinois and Indiana south of Route 50, Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

### **Selective Equipment Application**

**USE INSTRUCTIONS:** This product may be applied using a hooded sprayer for weed control in between rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instructions on the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**RESTRICTIONS:** Corn must be at least 12 inches tall, measured without extending leaves. Do not apply more than 22 fluid ounces of this product per acre for each hooded sprayer application and no more than 64 fluid ounces per acre per year total.

### **Spot Application**

**USE INSTRUCTIONS:** This product may be applied as a spot application prior to silking of corn. See the "ANNUAL WEEDS RATE SECTION" and the "PERENNIAL WEEDS RATE SECTION" of this label for appropriate spray solution concentrations.

**RESTRICTIONS:** Do not apply this product to more than 10 percent of the total field area to be harvested.

### **Preharvest**

**USE INSTRUCTIONS:** Up to 64 fluid ounces of this product per acre may be applied just prior to corn harvest using ground application equipment, or up to 44 fluid ounces per acre using aerial application equipment, when kernel-fill is complete and the corn is physiologically mature (black layer formed) and grain moisture is 35 percent or less.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied up to 3.3 quarts per acre as a single ground broadcast application for weed control after harvest of corn. Higher rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in corn. Read and follow label directions for all products in the tank mixture.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or feeding of vegetation within the application area. Application of this product must be made a minimum of 30 days prior to planting any crop not listed on this label. Do not exceed a total combined application rate of 5.3 quarts of this product per acre per year.

### **9.3 Cotton**

**TYPES OF APPLICATION:** Those listed in Section 9.0, plus Selective Equipment Application; Spot Application; Preharvest

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

For directions for use with cotton with Roundup Ready Technology, cotton with Roundup Ready Flex Technology, and cotton with GlyTol Technology, see the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label.

### **Preplant, At-Planting, Preemergence**

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton, but prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may also be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text: , or one or more of the products listed,*] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to planting or the emergence of cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clomazone; diuron; flumioxazin; fluometuron; fomesafen; metolachlor; s-metolachlor; norflurazon; pendimethalin; prometryn; pyriithiobac-sodium; saflufenacil

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

Caparol 4L; Command 3ME; Cotoran 4L; Cotton Pro; Direx 4L; Dual MAGNUM; Dual II MAGNUM; Karmex DF; Prowl 3.3 EC; Prowl H2O; Reflex; Sharpen Powered by Kixor; Stalwart; Staple LX; Valor SX; Warrant; Warrant Ultra]

---

### **Selective Equipment Application**

USE INSTRUCTIONS: This product may be applied using a hooded or shielded sprayer, or over the top of cotton using a wiper applicator to control tall weeds. See additional instructions on the use of this selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

### **Spot Application**

USE INSTRUCTIONS: This product may be applied as a spot application to targeted weeds in cotton prior to boll opening.

RESTRICTIONS: Do not apply this product to more than 10 percent of the total field area to be harvested.

### **Preharvest**

USE INSTRUCTIONS: This product provides weed control and cotton re-growth inhibition when applied prior to harvest. For weed control, apply at rates given in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label. For cotton re-growth inhibition, apply 16 to 44 fluid ounces of this product per acre. Make preharvest application only after sufficient bolls have developed to produce the desired yield. Application made prior to this time could affect maximum yield potential.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

TANK MIXTURES: This product may be tank-mixed with DEF 6 (EPA Reg. No. 264-730; *tribufos*), Folex 6 EC (EPA Reg. No. 5481-504; *tribufos*), or Ginstar EC (EPA Reg. No. 264-634; *thidiazuron*, *diuron*), to enhance cotton leaf-drop. Read and follow label directions for all products used in the tank mixture.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR PREHARVEST APPLICATION TO COTTON.

### 9.4 Fallow Systems

This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

TYPES OF APPLICATION: Chemical Fallow; Preplant Fallow Beds; Aid-to-Tillage

#### Chemical Fallow

USE INSTRUCTIONS: This product may be used as a substitute for tillage to control annual weeds in fallow fields. Broadcast or spot application will also control or suppress many perennial weeds in fallow fields. Tank-mix this product with 2,4-D or dicamba for a broader weed control spectrum. Application of up to 44 fluid ounces of this product per acre may be made onto fallow fields using aerial application equipment where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

PRECAUTIONS: Some crop injury could occur if dicamba is applied within 45 days of planting.

#### Preplant Fallow Beds

USE INSTRUCTIONS: This product will control weeds listed in the “ANNUAL WEEDS RATE SECTION,” “PERENNIAL WEEDS RATE SECTION” and “WOODY BRUSH, TREES AND VINES RATE SECTION” of this label prior to planting.

TANK MIXTURES: Apply 8 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Goal 2XL (EPA Reg. No. 62719-424; *oxyfluorfen*) to control the following weeds up to the maximum height or length indicated: 3 inches — common cheeseweed, chickweed, groundsel; 6 inches — London rocket, shepherd's-purse.

Apply 11 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Goal 2XL to control the following weeds up to the maximum height or length indicated: 6 inches — common cheeseweed, groundsel, mare's tail (*Coryza canadensis*); 12 inches — chickweed, London rocket, shepherd's-purse.

#### Aid-to-Tillage

USE INSTRUCTIONS: This product may be used in conjunction with tillage practices in fallow systems, or prior to the planting of crops listed on this label (preplant), to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of this product in 3 to 10 gallons of water per acre before weeds are 6 inches in height. Application must be followed by conventional tillage no later than 15 days after application and before re-growth occurs. Allow a minimum of 1 day after application before tillage.

PRECAUTIONS: Tank mixtures with residual herbicides could result in reduced performance of this product.

### 9.5 Grain Sorghum (Milo)

TYPES OF APPLICATION: Those listed in Section 9.0, plus Spot Application; Wiper Application; Preharvest

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank mixture before, during or after planting grain sorghum, but prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to planting or the emergence of grain sorghum. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products used in the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products to be added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; atrazine; metolachlor; s-metolachlor; saflufenacil

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Bicep II MAGNUM; Bicep II MAGNUM FC; Bicep Lite II MAGNUM; Degree Xtra; Dual MAGNUM; Dual II MAGNUM; Sharpen Powered by Kixor; Warrant]

---

For hard-to-control annual weeds, such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply 22 fluid ounces of this product per acre in a tank mixture with one or more of the products listed here.

For control of other annual weeds listed on this label, apply 16 to 22 fluid ounces of this product per acre when weeds are less than 6 inches tall and 22 to 32 fluid ounces per acre when weeds are over 6 inches tall. When using a nitrogen solution as the carrier, the application rate might need to be increased to achieve acceptable weed control.

### Spot Application, Wiper Application

USE INSTRUCTIONS: This product may be applied as a targeted spot application in grain sorghum before heading. This product may also be applied over the top of grain sorghum using a wiper applicator to control or suppress tall weeds growing above the crop canopy. See additional instructions on the use of wiper applicators in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label.

RESTRICTIONS: For spot application, do not apply this product to more than 10 percent of the total field area to be harvested. When applied using a wiper applicator, allow a minimum of 40 days between application and harvest. Do not use a roller applicator. Do not feed or graze grain sorghum fodder or ensile vegetation collected from within the application area.

### Selective Equipment Application

USE INSTRUCTIONS: This product may be applied using a hooded sprayer for weed control in between rows of grain sorghum. Make application before grain sorghum sends tillers between the drill rows. If tillers are sprayed with this herbicide, the main plant could be damaged or destroyed. Contact of this product in any manner with any vegetation to which application is not intended could cause damage. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instructions on the use of hooded sprayers in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** Grain sorghum must be at least 12 inches tall, measured without extending leaves. Do not graze or feed grain sorghum forage or fodder following application of this product using a hooded sprayer. Do not apply more than 22 fluid ounces of this product per acre per hooded sprayer application and no more than 64 fluid ounces per acre per year total.

### **Preharvest**

**USE INSTRUCTIONS:** Up to 44 fluid ounces of this product per acre may be applied after sorghum grain has reached 30 percent moisture or less. As with other herbicides that cause sudden plant death, avoid preharvest application of this product on grain sorghum (milo) infected with charcoal rot as lodging can occur.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest of grain sorghum. Preharvest application of this product on grain sorghum (milo) is not registered for use in California.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied up to 3.3 quarts per acre for weed control after harvest of grain sorghum. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in grain sorghum (milo). Read and follow label directions for all products in the tank mixture.

This product may be applied to grain sorghum stubble following harvest to control or suppress re-growth. Apply 22 fluid ounces of this product per acre for control or 16 fluid ounces per acre for suppression.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label. Do not exceed a total combined application rate of 5.3 quarts of this product per acre per year

## **9.6 Herbs and Spices**

**LABELED CROPS:** Allspice; Angelica; Star anise; Annatto (seed); Balm; Basil; Borage; Burnet; Camomile; Caper buds; Caraway; Black caraway; Cardamom; Cassia bark; Cassia buds; Catnip; Celery seed; Chervil (dried); Chive; Chinese chive; Cinnamon; Clary; Clove buds; Coriander leaf (cilantro or Chinese parsley); Coriander seed (cilantro); Costmary; Culantro (leaf); Culantro (seed); Cumin; Curry (leaf); Dill (dillweed); Dill (seed); Epazote; Fennel seed (common and Florence); Fenugreek; White ginger flower; Grains of paradise; Horehound; Hyssop; Juniper berry; Lavender; Lemongrass; Lovage (leaf and seed); Mace; Marigold; Marjoram (including oregano); Mexican oregano; Mioga flower; Mustard (seed); Nasturtium; Nutmeg; Parsley (dried); Pennyroyal; Pepper (black and white); Pepper leaves; Peppermint; Perilla; Poppy (seed); Rosemary; Rue; Saffron; Sage; Savory (summer and winter); Spearmint; Stevia leaves; Sweet bay; Tansy; Tarragon; Thyme; Vanilla; Wintergreen; Woodruff; Wormwood

**TYPES OF APPLICATION:** Those listed in Section 9.0, plus Spot Application (peppermint, spearmint); Wiper Application (peppermint, spearmint)

**PRECAUTIONS:** This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Remove residual product from the plastic prior to planting with a single 0.5-inch application of water, either by natural rainfall or by irrigation. Ensure that the wash water flushes off the plastic mulch and does not enter the transplant holes. Application made at crop emergence will result in injury or death of emerged seedlings.

### **Spot Application, Wiper Application (Peppermint and Spearmint Only)**

**USE INSTRUCTIONS:** This product may be applied as a targeted spot application in peppermint and spearmint or over the top of peppermint and spearmint using a wiper applicator to control tall weeds.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Application may be repeated in the same area at 30-day intervals. See additional instructions on the use of wiper applicators in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. For spot application, do not apply this product to more than 10 percent of the total field area to be harvested.

### 9.7 Oilseed Crops

LABELED CROPS: Borage; Buffalo gourd; Calendula; Canola; Castor oil plant; Chinese tallowtree; Crambe; Cuphea; Echium; Euphorbia; Evening primrose; Flax; Gold of pleasure; Hare’s ear mustard; Jojoba; Lesquerella; Meadowfoam; Milkweed; Mustard; Niger seed; Oil radish; Poppy; Rape; Rose hip; Safflower; Sesame; Stokes aster; Sunflower; Sweet rocket; Tallowwood; Tea oil plant; Vernonia

For directions for use with canola with Roundup Ready Technology and canola with TruFlex™ Technology, see the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label.

TYPES OF APPLICATION: Those listed in Section 9.0, plus Preharvest (except buffalo gourd)

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product for use in safflower, sunflower and all other oilseed crops listed in this section, if a preharvest application is to be made. If a preharvest application is NOT to be made, the maximum application rate of this product for all preemergence, selective equipment and post-harvest applications in any oilseed crop listed in this section is limited only by the maximum of 5.3 quarts per acre per year. If a preharvest application is intended to be made to any crop listed in this section, except buffalo gourd, the maximum combined total of all preemergence and selective equipment applications is limited as indicated in the following table. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

<b>Maximum Application Rates if a Preharvest Application is Made</b>	
<b>Safflower</b>	
Combined total for all Preemergence and Selective Equipment applications	64 fluid ounces per acre
Preharvest application	64 fluid ounces per acre
<b>Sunflower</b>	
Combined total for all Preemergence and Selective Equipment applications	22 fluid ounces per acre
Preharvest application	22 fluid ounces per acre
<b>All Other Oilseed Crops Listed (Except Buffalo Gourd)</b>	
Combined total for all Preemergence and Selective Equipment applications	44 fluid ounces per acre
Preharvest application	32 fluid ounces per acre

RESTRICTIONS: Do not exceed a total application rate of 5.3 quarts of this product per acre per year. Preharvest application is not permitted on buffalo gourd.

#### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting oilseed crops listed in this section, but must be applied prior to crop emergence. Observe the maximum application rates listed at the beginning of this section.

TANK MIXTURES: For sunflower, a tank mixture with pendimethalin may be applied before, during or after planting into conventionally tilled soil, a cover crop, established sod or previous crop residue.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** See the use instructions at the beginning of this section for important information on maximum application rates for preemergence and selective equipment applications of this product.

### **Selective Equipment Application**

**USE INSTRUCTIONS:** This product may be applied using a wiper applicator or shielded sprayer between crop rows once the crop is established. Observe the maximum application rates listed at the beginning of this section. See additional instructions on the use of wiper applicators and hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

### **Preharvest (Except Buffalo Gourd)**

**USE INSTRUCTIONS:** This product provides weed control and serves as a harvest aid when applied to a physiologically mature oilseed crop listed in this section. For safflower, up to 64 fluid ounces of this product may be applied per acre when seed has lost its opaque character, approximately 20 to 30 days after the end of flowering of the secondary branches. For sunflower, up to 22 fluid ounces of this product per acre may be applied when the backsides of sunflower heads are yellow and bracts are turning brown, and seed moisture content is less than 35 percent. For all other oilseed crops listed in this section (except buffalo gourd), up to 32 fluid ounces of this product per acre may be applied prior to harvest.

**RESTRICTIONS:** DO NOT MAKE A PREHARVEST APPLICATION if you have exceeded the maximum application rates for the combined total of all preemergence and selective equipment applications listed in the table at the beginning of this section. Make only 1 preharvest application of this product and allow a minimum of 7 days between application and harvest or feeding to livestock. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label. Preharvest application is not allowed on buffalo gourd or on canola with Roundup Ready Technology, canola with TruFlex™ Technology, or other listed glyphosate-tolerant canola.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied at up to 3.3 quarts per acre for weed control after harvest of oilseed crops. Higher application rates might be needed for control of large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in the crop harvested. Read and follow label directions for all products in the mix.

**RESTRICTIONS:** Do not exceed a total application rate of 5.3 quarts of this product per acre per year. Allow a minimum of 7 days between application of this product and harvest or feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

## **9.8 Soybean**

**TYPES OF APPLICATION:** Those listed in Section 9.0, plus Spot Application; Selective Equipment Application; Preharvest

For directions for use with soybean with Roundup Ready Technology, soybean with Roundup Ready 2 Yield Technology, soybean with GT27 Technology, and soybean with Enlist E3 Technology, see the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label.

### **Preplant, At-Planting, Preemergence**

**USE INSTRUCTIONS:** This product may be applied at up to 3.3 quarts per acre alone or in a tank mixture before, during or after planting soybean, but prior to crop emergence.

**TANK MIXTURES:** This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may also be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*., or one or more of the products listed,] and applied prior to planting or the



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

emergence of soybean. Ensure that the product used is labeled for application prior to planting or the emergence of soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for all products in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

[*Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fenoxaprop-p-ethyl; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; quizalofop-p-ethyl; saflufenacil; sulfentrazone; tribenuron methyl; trifluralin

**Axiom DF** (EPA Reg. No. 264-766; *metribuzin, flufenacet*); **Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

Aim EC; Aim EW; Anthem; Anthem Flex; Anthem Maxx; Assure II; Authority Assist; Authority Elite; Authority First DF; Authority MAXX; Authority MTZ DF; Authority Supremem; Authority XL; Axiom DF; Blanket 4F; Boundary 6.5 EC; Cadet; Canopy; Canopy Blend; Canopy EX; Classic; Cobra; Command 3ME; Dual MAGNUM; Dual II MAGNUM; Enlite; Envive; Fierce; Fierce MTZ; Fierce XLT; FirstRate; Flexstar; Fusion; Linex 4L; Lorox DF; Mauler; Me-Too-Lachlor; Optill Powered by Kixor; Outlook; Phoenix; Prefix; Prowl 3.3 EC; Prowl H2O; Pursuit; Python WDG; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Sharpen Powered by Kixor; Spartan 4F; Treflan 4L; Treflan 4 EC; TriCor 4F; TriCor DF; Valor SX; Valor XLT; Verdict powered by Kixor; Warrant; Warrant Ultra; Zidua; Zidua PRO Powered by Kixor; Zidua SC]

---

For hard-to-control annual weeds, such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply 22 fluid ounces of this product per acre in a tank mixture with one of the products listed. For other annual weeds listed on this label, apply 16 to 22 fluid ounces of this product per acre when weeds are less than 6 inches tall and 22 to 32 fluid ounces per acre when weeds are over 6 inches tall.

### Spot Application

USE INSTRUCTIONS: This product may be applied as a targeted spot application in soybean prior to initial pod set.

RESTRICTIONS: Do not apply this product to more than 10 percent of the total field area to be harvested.

### Selective Equipment Application

USE INSTRUCTIONS: This product may be applied in soybean using a shielded applicator, hooded sprayer, wiper applicator or sponge bar. See additional instructions on the use of selective application equipment in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

### Preharvest

USE INSTRUCTIONS: This product may be applied to soybean prior to harvest after pods have set and lost all green color. Apply at rates given in the “ANNUAL WEEDS RATE SECTION” and “PERENNIAL WEEDS RATE SECTION” of this label. Take care to avoid excessive seed shatter loss due to ground application equipment.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** Do not apply more than 3.3 quarts of this product per acre for preharvest application using ground application equipment or more than 44 fluid ounces per acre when using aerial application equipment. Allow a minimum of 7 days between application and harvest of soybean. If the preharvest application rate is greater than 22 fluid ounces per acre, do not graze or harvest hay or fodder within the application area for livestock feed within 25 days of application. If the application rate is 22 fluid ounces per acre or less, the grazing restriction is reduced to 14 days after application.

### **9.9 Sugarcane**

**TYPES OF APPLICATION:** Those listed in Section 9.0, plus Spot Application

#### **Preplant, At-Planting, Preemergence**

**USE INSTRUCTIONS:** This product may be applied in or around sugarcane fields, or in fields prior to the emergence of plant cane.

**RESTRICTIONS:** Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

#### **Spot Application**

**USE INSTRUCTIONS:** This product may be applied as a targeted spot application in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using a handheld sprayer and a spray-to-wet technique. Better control of volunteer or diseased sugarcane can be obtained when this product is applied when there are at least 7 new leaves on the targeted cane. Avoid contact of this herbicide with healthy sugarcane plants as severe damage or destruction could result.

**RESTRICTIONS:** Do not feed or graze sugarcane foliage within the application area.

#### **Selective Equipment Application**

**USE INSTRUCTIONS:** This product may be applied using a hooded sprayer for weed control in between rows of sugarcane. See additional instructions on the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**PRECAUTIONS:** Do not allow weeds within the application area to come into contact with the crop.

#### **Fallow Treatment**

**USE INSTRUCTIONS:** This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane by applying 2.5 to 3.3 quarts of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow a minimum of 7 days after application before tillage. Aerial application of up to 64 fluid ounces per acre may be made onto fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for this application in sugarcane. Read and follow label directions for all products in the tank mixture.

#### **9.9.1 Sugarcane Ripening**

**USE INSTRUCTIONS:** This product may be used as a foliar-applied plant growth regulator to hasten ripening and extend the period of high sucrose level in both low- and high-tonnage sugarcane. Most of the sucrose increase is concentrated in the top nodes of the cane stalk. To maximize sugar recovery where topping is practiced at harvest, top at the base of the fourth leaf. Consult your state sugarcane authority or local Bayer CropScience representative regarding the degree of sucrose response that can be anticipated prior to application of this product.

As a result of leaf desiccation, improved trash burn can be expected.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Apply this product at the following rates and timing according to the State in which the sugarcane is grown. Use a higher application rate within the given range when applying to sugarcane under adverse ripening conditions or to less responsive varieties.

**FLORIDA** – Apply 5 to 12 fluid ounces of this product per acre 3 to 5 weeks before harvest of LAST RATOON CANE ONLY.

**HAWAII** – Apply 9 to 21 fluid ounces of this product per acre 4 to 10 weeks before harvest.

**LOUISIANA** – Apply 4 to 12 fluid ounces of this product per acre 3 to 7 weeks before harvest of RATOON CANE ONLY.

**PUERTO RICO** – Apply 5 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

**TEXAS** – Apply 5 to 12 fluid ounces of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.

**PRECAUTIONS:** Application of this product could initiate development of shooting eyes. This product might not increase the sucrose content of sugarcane under conditions of good natural ripening. Within 2 to 3 weeks after application, this product could produce a slight yellowing to a pronounced browning and drying of leaves, and a shortening of upper internodes. Spindle death could occur.

Rainfall within 6 hours after application could reduce the effectiveness of this product.

Application to sugarcane grown for seed could result in a reduction in germination or vigor. To the extent consistent with applicable law, buyer and all users are responsible for such loss or damage in connection with the preharvest use of this product on sugarcane grown for seed.

**RESTRICTIONS:** Do not feed or graze sugarcane forage following application. Do not plant subsequent crops within 30 days after application of this product other than the following: alfalfa or other forage legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybean, squash (all types) or wheat.

Do not apply for enhanced ripening to any crops other than sugarcane. Use of this product in any manner not consistent with this label could result in injury to persons, animals or crops, or have other unintended consequences.

### 9.10 Vegetable Crops

THIS SECTION PROVIDES DIRECTIONS FOR USE THAT APPLY TO ALL VEGETABLE CROPS LISTED IN THE FOLLOWING SECTIONS. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC DIRECTIONS FOR USE, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

**TYPES OF APPLICATION:** Chemical Fallow; Preplant Fallow Beds; Preplant; At-Planting; Preemergence; Prior to Transplanting Vegetables; Hooded Sprayer in Row Middles; Shielded Sprayer in Row Middles; Wiper Application in Row Middles; Directed Application (non-bearing ginseng only); Wiper Application (carrot, rutabaga, sweet potato only); Post-Harvest

**PRECAUTIONS:** This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Remove residual product from the plastic prior to planting with a single 0.5-inch application of water, either by natural rainfall or by irrigation. Ensure that the wash water flushes off the plastic mulch and does not enter the transplant holes. Application of this product at crop emergence will result in injury or death to emerged seedlings.

Avoid contact of this herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from the plastic mulch), or fruit of crops, as severe crop injury or destruction could result. Transplanted seedlings coming into contact with weeds that are still wet with a spray solution of this product could result in significant crop injury.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Preemergence application must be made before the crop emerges from the soil to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, make hooded sprayer, shielded sprayer and wiper applications in row middles prior to vine development, otherwise severe crop injury or destruction could result.

RESTRICTIONS: Unless otherwise directed, application using selective equipment, including wiper applicators and hooded sprayers, must be made a minimum of 14 days prior to harvest. Post-harvest and fallow applications must be made a minimum of 30 days prior to the planting of any crop not listed on this label. See additional use instructions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

### 9.10.1 Brassica Vegetables

LABELED CROPS: Broccoli; Chinese broccoli (gai lan); Broccoli raab (rapini); Brussels sprouts; Cabbage; Chinese cabbage (bok choy); Chinese cabbage (napa); Chinese mustard cabbage (gai choy); Cauliflower; Cavalo broccolo; Collards; Kale; Kohlrabi; Mizuna; Mustard greens; Mustard spinach; Rape greens

### 9.10.2 Bulb Vegetables

LABELED CROPS: All cultivars, varieties and/or hybrids of Chive (including Chinese); Daylily; Elegans hosta; Fritillaria; Garlic (including great-headed, serpent); Kurrat; Leek (including lady's, wild); Onion (including Beltsville bunching, bulb, Chinese, fresh, green, macrostem, pearl, potato, tree, Welsh); Shallot

### 9.10.3 Cucurbit Vegetables and Fruits

LABELED CROPS: Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra); Melons (all); *Momordica spp* (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon); Pumpkin; Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon

RESTRICTIONS: For cantaloupe, casaba melon, crenshaw melon, cucumber, gherkin, gourds, honeydew melon, honey ball melon, mango melon, melons (all), muskmelon, Persian melon, pumpkin, squash (summer, winter), and watermelon, allow a minimum of 3 days between application and planting.

### 9.10.4 Leafy Vegetables

LABELED CROPS: Amaranth (Chinese spinach); Arugula (roquette); Beet greens; Cardoon; Celery; Chinese celery; Celtuce; Chaya; Chervil; Edible-leaved chrysanthemum; Garland chrysanthemum; Corn salad; Cress (garden, upland); Dandelion; Dock (sorrel); Dokudami; Endive (escarole); Florence fennel; Gow kee; Lettuce (head, leaf); Orach; Parsley; Purslane (garden, winter); Radicchio (red chicory); Rhubarb; Spinach; New Zealand spinach; Vine spinach; Swiss chard; Watercress (upland); Water spinach

RESTRICTIONS: For watercress, allow a minimum of 3 days between application and seeding. Do not apply this product during the period between seeding and emergence.

### 9.10.5 Fruiting Vegetables

LABELED CROPS: All cultivars, varieties and/or hybrids of Eggplant (including African, pea, scarlet); Cocona; Garden huckleberry; Goji berry; Groundcherry (*Physalis spp*); Martinyinia; Naranjilla; Okra;

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Pepino; Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); Roselle; Sunberry; Tomatillo; Tomato

RESTRICTIONS: Allow a minimum of 3 days between application and planting. For tomato and tomatillo, do not apply this product using a hooded or shielded sprayer in row-middles because of the potential for crop injury.

### 9.10.6 Legume Vegetables (Succulent or Dried)

LABELED CROPS: Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, white sweet lupin); Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); Broad bean (fava); Chickpea (garbanzo); Guar; Jackbean; Lablab bean; Lentil; Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); Pigeon pea; Soybean (immature seed); Sword bean

TYPES OF APPLICATION: Those listed in Section 9.0, plus Spot Application (dry varieties only); Preharvest (dry varieties only)

#### Spot Application (Dry Varieties Only)

USE INSTRUCTIONS: This product may be applied as a targeted spot application to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel) and milkweed in any dry legume variety listed in this section, except cowpeas or field (feed) peas. Up to 22 fluid ounces of this product per acre may be applied in dry beans, or up to 64 fluid ounces per acre in dry peas, lentils and chickpeas, in 10 to 20 gallons of water using ground application equipment, or apply a 2-percent solution using a handheld sprayer. For maximum performance, apply this product when these weeds are at or beyond the bud stage of growth.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Only one spot application may be made per year. Do not combine spot application with a preharvest broadcast application on the same crop area. Allow a minimum of 30 days between application and the planting of any crop not listed in this label. Do not feed vines and hay from the application area to livestock. Do not apply this product as a spot application in cowpeas or field (feed) peas, since these are considered to be grown only as livestock feed.

#### Preharvest (Dry Varieties Only)

USE INSTRUCTIONS: This product may be applied over the top of any dry legume variety listed in this section prior to harvest, except cowpeas or field (feed) peas. Up to 22 fluid ounces of this product per acre may be applied in dry beans, or up to 64 fluid ounces per acre in dry peas, lentils, and chickpeas, in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Preharvest application is not recommended for legumes grown for seed, as a reduction in germination or vigor may occur.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Only one preharvest application may be made per year. Do not combine a preharvest application with a spot treatment application on the same crop area. Allow a minimum of 30 days between application and the planting of any crop not listed on this label. Do not feed vines and hay from the application area to livestock. Do not make a preharvest application of this product in cowpeas or field (feed) peas since these crops are considered to be grown only as livestock feed.

### 9.10.7 Root and Tuber Vegetables

LABELED CROPS: Arracacha; Arrowroot; Chinese artichoke; Jerusalem artichoke; Beet (garden); Burdock; Canna; Carrot; Cassava (bitter and sweet); Celeriac; Chayote (root); Chervil (turnip-rooted);

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Chicory; Chufa; Dasheen (taro); Galangal; Ginger; Ginseng; Horseradish; Leren; Kava (turnip-rooted); Parsley (turnip-rooted); Parsnip; Potato; Radish; Oriental radish; Rutabaga; Salsify; Black salsify; Spanish salsify; Skirret; Sweet potato; Tanier; Turmeric; Turnip; Wasabi; Yacon; Yam bean; True yam

TYPES OF APPLICATION: Those listed in Section 9.0, plus Directed Application (non-bearing ginseng); Wiper Application (carrot, rutabaga, sweet potato)

### **Directed Application (Non-Bearing Ginseng Only)**

USE INSTRUCTIONS: This product may be applied for weed control in established non-bearing ginseng using a boom sprayer, CDA, shielded sprayer, wiper applicator, handheld or backpack wand, lance, or orchard gun. See additional use instructions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Control the application so as not to allow any contact of this product with the ginseng plant. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation could result in discoloration, stunting or destruction.

RESTRICTIONS: Application must be made a minimum of one year prior to ginseng harvest.

### **Wiper Applicator (Carrot, Rutabaga and Sweet Potato Only)**

USE INSTRUCTIONS: A 33-percent solution of this product by volume in water may be applied using a wiper applicator over the top of carrot, rutabaga and sweet potato for the control of tall weeds. See additional use instructions for wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: For carrot, a maximum of two wiper or sponge bar applications may be made a minimum of 60 days prior to harvest following the first application and 7 days prior to harvest following the second application or if only one wiper application is made over the top of the carrot crop. For rutabaga, allow a minimum of 14 days between application and harvest. For sweet potato, a maximum of five wiper or sponge bar applications may be made with a minimum of 14 days between applications and a minimum of 7 days prior to harvest.

## **9.11 Miscellaneous Crops**

LABELED CROPS: Aloe vera; Asparagus; Bamboo shoots; Globe artichoke; Okra, Peanut; Pineapple; Sugarbeet

TYPES OF APPLICATION: Those listed in Section 9.0, plus Spot Application (asparagus)

For directions for use with sugarbeet with Roundup Ready Technology, see the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label.

PRECAUTIONS: Preemergence application must be made before the crop emerges from the soil to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of crop injury. In crops with vines, apply this product in row middles using a hooded sprayer, shielded sprayer or wiper applicator prior to vine development, otherwise severe crop injury or destruction could result.

### **Spot Weed Control, Site Preparation**

USE INSTRUCTIONS: This product may be applied for spot weed control and site preparation prior to planting or transplanting crops listed in this section.

PRECAUTIONS: This product could cause crop injury when applied prior to transplanting or direct-seeding crops into plastic mulch. Remove residual product from the plastic prior to planting with a single 0.5-inch application of water, either by natural rainfall or by irrigation. Ensure that the wash water flushes off the plastic mulch and does not enter transplant holes.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

RESTRICTIONS: Allow a minimum of 21 days between residue removal and transplanting. Do not apply this product within 7 days prior to emergence of the first asparagus spears. Do not feed or graze pineapple forage from within the application area.

### Spot Application (Asparagus)

USE INSTRUCTIONS: This product may be applied immediately after cutting asparagus, but prior to the emergence of new spears.

RESTRICTIONS: Do not apply this product to more than 10 percent of the total field area to be harvested. Do not harvest asparagus within 5 days of a spot application.

### Post-Harvest in Asparagus

USE INSTRUCTIONS: This product may be applied for weed control after the last harvest of asparagus and all spears have been removed. If spears are allowed to re-grow, delay application until ferns have developed and make the application as a directed or shielded spray in order to avoid contact of this product with ferns, stems or spears. See additional use instructions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Direct contact of this product with asparagus could result in serious crop injury.

## 10.0 TREE, VINE AND SHRUB CROPS

THIS SECTION PROVIDES DIRECTIONS FOR USE THAT APPLY TO ALL TREE, VINE, AND SHRUB CROPS LISTED IN THE FOLLOWING SECTIONS. SEE THE INDIVIDUAL CROP SECTIONS FOR SPECIFIC DIRECTIONS FOR USE, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATION: Preplant (site preparation); Broadcast Spray; Selective Equipment (shielded sprayer, wiper applicator), Directed Spray and Spot Application in Middles (between rows of trees, vines or bushes) and Strips (within rows of trees, vines or bushes); Site Weed Control; Perennial Grass Suppression; Cut Stump Application

USE INSTRUCTIONS: Unless specifically prohibited in the individual crop sections that follow, this product may be applied using a boom sprayer, controlled droplet applicator (CDA), shielded sprayer, wiper applicator, handheld or backpack sprayer, lance or orchard gun, in middles (between rows of trees, vines or bushes) and strips (within rows of trees, vines or bushes), for weed control or perennial grass suppression in established tree fruit and nut groves, orchards and vineyards. It may also be used for site preparation prior to planting or transplanting these crops.

Apply 11 fluid ounces to 3.3 quarts of this product per acre as directed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label. Use a higher application rate within a given range when weeds are stressed, growing in dense populations or greater than 12 inches tall. Application may be repeated as needed up to a maximum of 7 quarts of this product (8 pounds of glyphosate acid) per acre per year. Allow a minimum of 14 days between applications.

PRECAUTIONS: Use extreme care to avoid contact of this herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of desirable trees, canes and vines. Avoid application when recent pruning wounds or other mechanical injury have occurred. Contact of this product with other than matured brown bark could result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops where the potential for crop contact is high, and then only where there is sufficient clearance. For application in strips (within rows of trees), only selective equipment (directed spray, hooded sprayer, shielded sprayer, or wiper applicator) may be used in order to minimize the potential for overspray or drift of this product onto the crop. For berry crops, hooded sprayers must be fully enclosed including top, sides, front and back. Only wiper applicators or shielded sprayers capable of preventing all contact of this product with the crop may be used. See additional use instructions and precautions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** Before using this product, read and follow the Maximum Application Rate information in the “PRODUCT INFORMATION” section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined with the use of any and all other herbicides containing glyphosate as the active ingredient, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate. Allow a minimum of 3 days between application and transplanting.

### **Middles and Strips Management**

**USE INSTRUCTIONS:** This product will control or suppress annual and perennial weeds and ground covers growing between rows (middles) and within rows (strips) of tree, vine and shrub crops listed on this label. If weeds are under drought stress, irrigate prior to application. Reduced weed control could result if weeds have been recently mowed at the time of application.

**TANK MIXTURES:** A tank mixture of this product plus Goal 2XL (EPA Reg. No. 62719-424; *oxyfluorfen*) may be applied for annual weed control between rows and within rows of a variety of tree, vine and shrub crops when weeds are stressed or growing in dense populations. Application of 11 to 22 fluid ounces of this product per acre plus an appropriate rate of Goal 2XL will control annual weeds with a maximum height or length of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd’s-purse, annual sowthistle, filaree (suppression only), horseweed/marestail, stinging nettle and common purslane (suppression only). This tank-mix will also control common cheeseweed (malva) or hairy fleabane with a maximum height or length of 3 inches.

This product may also be applied to row middles and strips in tank mixtures with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed].

---

*[Active ingredients and products to be added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; bromacil; clethodim; diuron; fluazifop-p-butyl; flumioxazin; glufosinate-ammonium; indaziflam; napropamide; norflurazon; oryzalin; oxyfluorfen; pendimethalin; penoxsulam; pyraflufen ethyl; rimsulfuron; saflufenacil; sethoxydim; simazine; thiazopyr

**Alion** (EPA Reg. No. 264-1106; *indaziflam*); **Rely 280** (EPA Reg. No. 264-829; *glufosinate-ammonium*)

Alion; Chateau Herbicide SW; Devrinol 2-XT; Devrinol 50-DF; Devrinol 50-DF Ornamental; Devrinol DF-XT Selective; Devrinol DF-XT Ornamental; Direx 4L; Dri-Clean; Fusilade II Turf & Ornamental; Fusilade DX; Goal 2XL; GoalTender; Karmex DF; Matrix FNV; Matrix SG; Orchard Master Broadleaf; Orchard Master CA Broadleaf; Pindar GT; Poast; Poast Plus; Prowl 3.3 EC; Prowl H2O; Princep 4L; Princep Caliber 90; Princep Liquid; Rely 280; Select 2 EC; Select Max Herbicide with Inside Technology; Surflan AS Agricultural; Surflan AS Specialty; Surflan Flex; Surflan XL 2G; Treevix Powered by Kixor; Venue; Visor Broadcrop]

---

Ensure that the product used is labeled for application within the crop being grown and at the maturity stage of the trees (bearing or non-bearing). It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products in the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

**RESTRICTIONS:** Do not apply these tank mixtures in Puerto Rico.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Perennial Grass Suppression

This product will suppress perennial grasses such as bahiagrass, bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass and quackgrass that are grown as ground covers in tree, vine and shrub crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 4 fluid ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 4 fluid ounces of this product per acre. Do not add ammonium sulfate to the spray mix.

For maximum performance, mow cool-season grass covers in the spring to even their height and then apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4 fluid ounces of this product in 10 to 25 gallons of water per acre 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches prior to seedhead emergence. For suppression for up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 2 to 3 fluid ounces per acre about 45 days later. Make no more than two applications per year.

For burndown of bermudagrass, apply 22 to 44 fluid ounces of this product in 3 to 20 gallons of water per acre. Make this application only if reduction of the bermudagrass stand can be tolerated. When burndown is needed prior to harvest, make the application a minimum of 21 days prior to harvest to allow sufficient time for burndown to occur.

For suppression of bermudagrass, apply 4 to 11 fluid ounces of this product per acre east of the Rocky Mountains and 11 fluid ounces west of the Rocky Mountains in a total spray volume of 3 to 20 gallons per acre no sooner than 1 to 2 weeks after full green-up. If the bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when re-growth occurs and bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, apply 4 to 7 fluid ounces of this product per acre in shaded conditions or where a lesser degree of suppression is desired.

### Cut Stump Application

Application of this product to a freshly cut tree stump may be made during site preparation or site renovation to control re-growth and re-sprouting of stumps of many tree species, some of which are listed below.

Citrus Trees: Calamondin; Chironja; Citron; Citrus hybrids; Grapefruit; Kumquat; Lemon; Lime; Mandarin (tangerine); Orange (all); Pummelo; Tangelo (ugli); Tangor

Fruit Trees: Apple; Apricot; Cherry (sweet, sour); Crabapple; Loquat; Mayhaw; Nectarine; Olive; Peach; Pear; Plum/Prune (all); Quince

Nut Trees: Almond; Beechnut; Brazil nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (hazelnut); Hickory nut; Macadamia; Pecan; Pistachio; Walnut (black, English)

**USE INSTRUCTIONS:** Cut the tree close to the soil surface and immediately apply a 50 to 100-percent (undiluted) solution of this product to the freshly cut surface using application equipment capable of covering the entire cambium. A delay in application could result in reduced performance. For optimal performance on tree stumps, cut the tree during period of active growth and full leaf expansion and apply this product.

**PRECAUTIONS:** DO NOT MAKE A CUT STUMP APPLICATION WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MIGHT BE GRAFTED TO THE ROOTS OF THE CUT STUMP, AS INJURY COULD OCCUR IN THE ADJACENT TREES. Some sprouts, stems or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indicator of a shared root system. Whether grafted or shared, injury is likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing a common root system.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### 10.1 Berry and Small Fruit Crops

**LABELED CROPS:** All cultivars, varieties and/or hybrids of Amur River grape; Aronia berry; Bayberry; Bearberry; Bilberry; Blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures de ronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora); Blueberry (highbush, lowbush); Buffaloberry; Che; Chilean guava; Chokecherry; Cloudberry; Cranberry (including highbush); Currant (black, Buffalo, red, native); Elderberry; European barberry; Gooseberry; Grape; Honeysuckle (edible); Huckleberry; Jostaberry; Juneberry (Saskatoon berry); Kiwifruit (fuzzy, hardy); Ligonberry; Maypop; Mountain pepper berries; Mulberry; Muntries; Partridgeberry; Phalsa; Pincherry; Raspberry (black, red, wild); Riberry; Salal; Schisandra berry; Sea buckthorn; Serviceberry; Strawberry

**TYPES OF APPLICATION:** Those listed in Section 10.0

**PRECAUTIONS:** To avoid damage, spray solutions of this product must not be allowed to come into contact with desirable vegetation, including green shoots, canes or foliage. In the northeast and Great Lakes regions, apply this product in grape vineyards prior to the end of the bloom stage in order to avoid crop injury, or apply using a shielded sprayer or wiper applicator. **USE THIS PRODUCT WITH EXTREME CARE AROUND RASPBERRY, AS SERIOUS CROP DAMAGE CAN OCCUR IF THIS PRODUCT COMES INTO CONTACT WITH ANY PART OF THE VINE.** To the extent consistent with applicable law, grower assumes all responsibility for crop losses resulting from misapplication of this product.

**RESTRICTIONS:** Allow a minimum of 3 days between application of this product and transplanting. Allow a minimum of 30 days between application and harvest of cranberries or the planting of any crop not listed on this label. Allow a minimum of 14 days between application and harvest for all other berry and small fruit crops listed here. Do not apply this product using selective equipment in kiwifruit.

#### **Spot Application**

**USE INSTRUCTIONS:** Spot application using a handheld sprayer or other appropriate application equipment listed in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label may be used to control weeds in berry and small fruit crops listed in this section.

For control of weeds growing in dry ditches (interior and perimeter) of cranberry production areas, drop water level to remove standing water in ditches and apply a 1- to 2-percent solution of this product with a handheld sprayer to adequately wet the vegetation only; do not spray to the point of runoff. To achieve maximum weed control in dry ditches, apply this product within 1 day after water drawdown to ensure application to actively growing weeds and allow a minimum of 2 days after application before reintroduction of water.

**RESTRICTIONS:** Allow a minimum of 30 days between spot application and harvest of cranberries. Do not apply directly to water. Use nozzles that produce medium- to large-sized droplets to minimize spray drift and avoid crop injury.

#### **Post-Harvest Application in Cranberry Bogs**

**USE INSTRUCTIONS:** This product may be applied for weed control after the harvest of berries and small fruits listed in this section. In cranberry bogs, apply this product after cranberry vines are dormant (after they have turned red) using a handheld sprayer, wiper applicator, or any other appropriate application equipment listed in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label. With a handheld sprayer, apply a 0.4- to 0.7-percent solution of this product to adequately wet the vegetation only;

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

do not spray to the point of runoff. With a handheld boom sprayer, apply 44 to 86 fluid ounces of this product per acre.

PRECAUTIONS: Even though vines appear dormant, contact of this product with desirable vegetation could result in damage or severe plant injury. Cranberry plants that are directly sprayed could be killed.

RESTRICTIONS: Apply this product only after cranberries have been harvested. Do not apply to more than 10 percent of the total bog. Allow a minimum of 6 months between post-harvest application and the next harvest of cranberries. Do not apply using aerial application equipment. Do not apply directly to water.

**10.2 Citrus Fruit Crops**

LABELED CROPS: All cultivars, varieties and/or hybrids of Calamondin; Chironja; Citron; Citrus Hybrids; Grapefruit (including Japanese summer); Kumquat; Lemon; Lime (including Australian desert lime, Australian finger lime, Australian round lime, Brown river finger lime, Mount white, New Guinea wild, Russell river, sweet, and Tahiti); Mandarin (including Mediterranean, Satsuma); Orange (all); Pummelo; Tangelo; Tangerine (Mandarin); Tangor; Uniq Fruit (ugli)

TYPES OF APPLICATION: Those listed in Section 10.0

USE INSTRUCTIONS: The following use instructions pertain to application in Florida and Texas only.

For burndown or control of the weeds listed below, apply this product at the specified rate in 3 to 30 gallons of water per acre. Where weed foliage is dense, make application in 10 to 30 gallons of water per acre.

To control goatweed, apply 44 to 64 fluid ounces of this product in 20 to 30 gallons of water per acre when target plants are actively growing. Apply 44 fluid ounces per acre when target plants are less than 8 inches tall and 64 fluid ounces per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Krovar I DF (EPA Reg. No. 5481-635; *bromacil, diuron*) or Karmex DF (EPA Reg. No. 66222-51; *diuron*) [*Alternative text: bromacil and/or diuron*] to the application mixture could improve weed control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Weed Species	Level of Perennial Weed Control at Various Application Rates (amount of this product per acre)			
	22 fl oz	44 fl oz	2 quarts	3.3 quarts
Bermudagrass	B	--	PC	C
Guinea grass				
<i>Texas and Florida Ridge</i>	B	C	C	C
<i>Florida Flatwoods</i>	–	B	C	C
Para grass	B	C	C	C
Torpedograss	S	--	PC	C

S = Suppression, PC = Partial Control, B = Burndown, C = Control

RESTRICTIONS: Allow a minimum of 1 day between application and harvest in citrus fruit crops. For citron groves, apply as directed spray only.

**10.3 Pome Fruit Crops**

LABELED CROPS: All cultivars, varieties and/or hybrids of Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear (including Asian pear); Quince (including Chinese and Japanese quince); Tejocote

TYPES OF APPLICATION: Those listed in Section 10.0

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

RESTRICTIONS: Allow a minimum of 1 day between application and harvest of pome fruit.

### 10.4 Stone Fruit Crops

LABELED CROPS: Apricot; Cherry (sweet, tart); Nectarine; Olive; Peach; Plum/Prune (all types); Plumcot

TYPES OF APPLICATION: Those listed in Section 10.0

PRECAUTIONS: Avoid application near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for a minimum of 2 years. ENSURE THAT NO PART OF A PEACH TREE IS CONTACTED WITH OVERSPRAY OR DRIFT OF THIS PRODUCT.

RESTRICTIONS: Allow a minimum of 17 days between application and harvest of stone fruit. In olive groves, apply this product as a directed spray only. Remove suckers and low-hanging limbs a minimum of 10 days prior to application.

### 10.5 Tree Nut Crops

LABELED CROPS: Almond; Beechnut; Betelnut; Brazil nut; Butternut; Cashew; Chestnut; Chinquapin; Coconut; Filbert (hazelnut); Hickory nut; Macadamia; Pecan; Pine nut; Pistachio; Walnut (black, English)

TYPES OF APPLICATION: Those listed in Section 10.0

RESTRICTIONS: Allow a minimum of 3 days between application and harvest of tree nuts, except coconut. Allow a minimum of 14 days between application and harvest of coconut.

### 10.6 Tropical and Subtropical Trees and Fruit Crops

LABELED CROPS: Ambarella; Atemoya; Avocado; Banana; Barbados cherry (acerola); Biriba; Blimbe; Breadfruit; Cacao (cocoa) bean; Canistel; Carambola (starfruit); Cherimoya; Coffee; Custard apple; Dates; Durian; Feijoa; Figs; Governor's plum; Guava; Illama; Imbe; Imbu; Jaboticaba; Jackfruit; Longan; Lychee; Mamey apple; Mango; Mangosteen; Marmaladebox (genip); Mountain papaya; Noni (Indian mulberry); Papaya; Pawpaw; Plantain; Persimmon; Pomegranate; Pulasan; Rambutan; Rose apple; Sapodilla; Sapote (black, mamey, white); Spanish lime; Soursop; Star apple; Sugar apple; Surinam cherry; Tamarind; Tea; Ti; Wax jambu

TYPES OF APPLICATION: Those listed in Section 10.0 and as a Bananacide (banana only).

RESTRICTIONS: Allow a minimum of 1 day between application and harvest in banana, coffee, guava, papaya, and plantain crops. Allow a minimum of 14 days between application and harvest of all other tropical or subtropical tree fruit listed here. In coffee and banana, delay application a minimum of 3 months after transplanting to allow the new plants to become established.

#### Bananacide (Banana Only)

USE INSTRUCTIONS: This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus, as well as non-infected banana plants, in order to establish a disease-free buffer around a plantation. Remove all fruit from the plants within the area prior to treatment. Inject 0.04 fluid ounce (1 milliliter) of this concentrated product (undiluted) for every 2 to 3 inches of pseudostem diameter of the banana plant to be controlled. Make the injection at least one foot above the ground, except for very small plants, which can be injected vertically into the top. Any subsequent re-growth must also be destroyed. Mechanically destroy all plants and mats (or units) within a 4-foot radius around a treated mat.

For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant might not show symptoms of the Banana Bunchy Top Virus for up to 125 days; therefore, it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** Do not apply more than 0.5 fluid ounce (15 milliliters) of this product per mat (or unit). Do not harvest any fruit or plant material from treated mats (or units) following injection. Do not allow livestock to consume treated plant material. Following transplant of new banana plants into treated areas, allow plants to become established for a minimum of 3 months before applying this product for weed control.

### **10.7 Vine Crops**

**LABELED CROPS:** Hops; Passion fruit

**TYPES OF APPLICATION:** Those listed in Section 10.0

**USE INSTRUCTIONS:** Apply this product for weed control only when green shoots, canes or foliage are not in the spray zone.

**RESTRICTIONS:** Allow a minimum of 14 days between application and harvest of these vine crops.

### **10.8 Miscellaneous Tree Food Crops**

**LABELED CROPS:** Cactus (all, including prickly pear and dragon fruit); Palm

**TYPES OF APPLICATION:** Those listed in Section 10.0

### **10.9 Non-Food Tree Crops**

**LABELED CROPS:** Pine; Poplar; Eucalyptus; Christmas trees; all other non-food tree crops

**TYPES OF APPLICATION:** Those listed in Section 10.0

**PRECAUTIONS:** Avoid contact of spray, drift or mist of this product with foliage or green bark of established Christmas trees and other pine trees. Desirable plants can be protected from the spray solution by using shields or coverings of impermeable materials.

**RESTRICTIONS:** DO NOT apply this product as a broadcast application over the top of plantations or tree crops.

### **Site Preparation**

**USE INSTRUCTIONS:** This product may be used for weed control prior to planting non-food tree crops.

**PRECAUTIONS:** Protect non-target plants from being sprayed with this product during site preparation application.

### **Directed Spray, Spot Application, Wiper Application**

**USE INSTRUCTIONS:** This product may be applied as a post-directed spray or spot application, or applied using a wiper applicator, around established Christmas trees, eucalyptus, poplar, and all other non-food tree crops.

## **11.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELAND**

**USE INSTRUCTIONS:** Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label for application rates of this product for specific weeds. When applied as directed, this product will control those annual and perennial grasses and broadleaf weeds listed. Application rates specified on this label for hard-to-control weeds, or those specified on separate supplemental labeling for this product, supersede rates listed in the "ANNUAL WEEDS RATE SECTION," "PERENNIAL WEEDS RATE SECTION" and "WOODY BRUSH, TREES AND VINES RATE SECTION" of this label. Additional information on hard-to-control weeds can be found on Fact Sheets published for this product.

**RESTRICTIONS:** Before using this product, read and follow the Maximum Application Rate information in the "PRODUCT INFORMATION" section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

with the use of any and all other herbicides containing glyphosate as the active ingredient, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate.

### 11.1 Alfalfa, Clover, and Other Forage Legumes

**LABELED CROPS:** Alfalfa; Clover; Kenaf; Kudzu; Lespedeza; Leucaena; Lupin; Sainfoin; Trefoil; Velvet bean; Vetch (all types)

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Spot Application; Wiper Applicator; Preharvest (except kenaf and leucaena); Stand Removal

For directions for use with alfalfa with Roundup Ready Technology, see the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label.

#### **Preplant, At-Planting, Preemergence**

**USE INSTRUCTIONS:** This product may be applied before, during or after planting crops listed in this section, but prior to crop emergence.

**RESTRICTIONS:** Remove domestic livestock before application.

#### **Spot Application, Wiper Application**

**USE INSTRUCTIONS:** This product may be applied as a targeted spot application or over the top of crops listed in this section using a wiper applicator. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label. Application may be repeated in the same area at 30-day intervals.

**RESTRICTIONS:** Make spot and wiper applications only in areas where the movement of domestic livestock can be controlled. Remove domestic livestock before application and wait a minimum of 3 days after application before grazing livestock or harvesting. Do not apply this product to more than 10 percent of the total field area at any one time.

#### **Weed Control in Dormant Alfalfa**

**USE INSTRUCTIONS:** This product will control or suppress many weeds, including quackgrass, downy brome and cheatgrass in dormant alfalfa. Apply 6 to 8 fluid ounces of this product per acre in the spring when alfalfa is dormant, after spring temperatures have warmed enough to encourage weed growth, but prior to initiation of trifoliolate leaf expansion of the alfalfa crop. Application made after expansion of the first trifoliolate leaf will cause growth reduction and reduced crop yield.

**PRECAUTIONS:** Improper application of this product to alfalfa can cause crop injury. Do not use this product on dormant alfalfa if a slight yield reduction in the first cutting cannot be tolerated. Slight discoloration of the alfalfa crop could occur, but will re-green and resume growth under moist soil conditions as effects of this product wear off.

**RESTRICTIONS:** Do not add ammonium sulfate to spray solutions of this product for application to dormant alfalfa. Do not make more than one application per year. Allow a minimum of 36 hours after application before grazing livestock or harvesting.

#### **Preharvest (Except Kenaf and Leucaena), Stand Removal**

**USE INSTRUCTIONS:** This product may be applied as a broadcast application prior to harvest (except in kenaf and leucaena) in declining stands or in any stand where severe crop injury or destruction is acceptable, or to remove an established stand of any forage legumes listed in this section. Application may be made at any time of the year to control annual and perennial weeds, including quackgrass. For control of quackgrass, apply in the spring, late-summer or fall when quackgrass is actively growing. For complete control of quackgrass, application must be followed by deep tillage. If the crop is to be harvested or grazed

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

by livestock, up to 44 fluid ounces of this product per acre may be applied in alfalfa and up to 32 fluid ounces per acre in all other legumes listed in this section. For complete removal of established stands of clover, it might be necessary to use a higher application rate, as listed in the “PERENNIAL WEEDS RATE SECTION” of this label.

**PRECAUTIONS:** This application can destroy an alfalfa stand and severely injure or destroy other legume crops listed, such as clover. Preharvest application on alfalfa grown for seed could result in a reduction in germination or vigor. To the extent consistent with applicable law, buyer and all users are responsible for such loss or damage in connection with the preharvest use of this product on alfalfa grown for seed.

**RESTRICTIONS:** Make only one application to an existing crop stand per year. Remove domestic livestock before application. Foliage within the application area can be harvested and fed to livestock according to the application rates and intervals defined in the following table. If applying at a rate greater than those listed here, do not harvest foliage for livestock feed or allow livestock to graze within the application area.

<b>Crop</b>	<b>Maximum Single Preharvest Application Rate (per acre)</b>	<b>Minimum Interval Between Application and Harvest or Livestock Grazing</b>
Alfalfa	44 fluid ounces	36 hours
All other legumes listed	32 fluid ounces	3 days

Crops listed on this label may be planted into the application area at any time; all other crops may be planted 30 days after application.

### 11.2 Conservation Reserve Program (CRP)

**TYPES OF APPLICATION:** Postemergence Weed Control in Dormant CRP Grasses; Wiper Application; Renovation (rotating out of CRP); Site Preparation

#### **Postemergence Weed Control in Dormant CRP Grasses, Wiper Application**

**USE INSTRUCTIONS:** Apply this product to suppress competitive growth and seed production of undesirable vegetation on CRP land. Application may be made using a wiper applicator to control tall weeds, or as a broadcast or targeted spot application to dormant CRP grasses. For selective weed control using broadcast application equipment, apply 5 to 8 fluid ounces of this product per acre in early-spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late-fall application may be made after desirable perennial grasses have reached dormancy.

**PRECAUTIONS:** Some stunting of CRP perennial grasses will occur if broadcast application is made when plants are not dormant.

**RESTRICTIONS:** Do not apply more than 2 quarts of this product per acre per year onto CRP land. No waiting period is required between application and grazing or harvesting for feed.

#### **Renovation (Rotating Out of CRP), Site Preparation**

**USE INSTRUCTIONS:** This product may be used to prepare CRP land for crop production. Refer to Federal, State or local use guides for CRP renovation information.

**RESTRICTIONS:** Crops listed on this label may be planted into the area at any time; all other crops may be planted 30 days after application.

### 11.3 Grass Seed and Sod Production

**LABELED CROPS:** Any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the “CEREAL AND GRAIN CROPS” section of this label

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Renovation; Removal of Established Stands; Site Preparation; Shielded Sprayer Application; Wiper Application; Spot Application; Creating Rows in Annual Ryegrass

### **Preplant, At-Planting, Preemergence, Renovation, Removal of Established Stand, Site Preparation**

USE INSTRUCTIONS: This product controls most existing vegetation for purposes of renovating turf or forage grass seed production areas, or for establishing turfgrass grown for sod. This product may be used to destroy undesirable grass vegetation when production fields are converted to alternate species or crops. Do not disturb soil or underground plant parts before application and delay tillage or renovation techniques, including vertical mowing, coring and slicing, for a minimum of 7 days after application to allow for herbicide translocation into underground plant parts.

Apply before, during, or after planting, or for renovation purposes. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. For maximum control of existing vegetation, delay planting until determining if any re-growth of underground plant parts will occur. Where repeat applications are necessary, sufficient re-growth must be attained prior to application. For optimal weed control in warm-season grasses, such as bermudagrass, summer or fall application of this product is best. Broadcast application of this product may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates of up to 3.3 quarts per acre may be used to totally remove an established stand of hard-to-kill grass species.

RESTRICTIONS: If application rate is 2 quarts of this product per acre or less, no waiting period between application and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Crops listed on this label may be planted into the area at any time; all other crops may be planted 30 days after application.

### **Selective Equipment Application**

USE INSTRUCTIONS: Apply 22 to 64 fluid ounces of this product in 10 to 20 gallons of water per acre using a shielded sprayer to control weeds between grass seed rows. Uniform planting in straight rows will aid shielded sprayer application. Best results can be obtained when the grass seed crop is small enough to easily pass by the protective shields. See additional instructions on the use of shielded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Any contact of this product on any vegetation to which application is not intended could cause damage.

### **Wiper Application**

USE INSTRUCTIONS: This product may be applied over the top of desirable grasses using a wiper applicator for the control of tall weeds. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

PRECAUTIONS: Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation could result in discoloration, stunting or destruction.

### **Spot Application**

USE INSTRUCTIONS: Apply a 1-percent solution of this product using a handheld sprayer to control weeds within established vegetation prior to heading of grasses grown for seed or to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS: This product will kill the desirable grasses along with the weeds. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted crop destruction.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### **Creating Rows in Annual Ryegrass**

USE INSTRUCTIONS: Use low-pressure nozzles or drop nozzles designed to target the application over a narrow band. Set nozzle height to establish the desired row spacing and apply 11 to 22 fluid ounces of this product per acre. Best results can be obtained when application is made before ryegrass reaches 6 inches in height. Use a higher application rate within this range when ryegrass is greater than 6 inches in height.

PRECAUTIONS: Take care not to spray or allow spray to drift outside target area in order to avoid unwanted crop destruction. To the extent consistent with applicable law, grower assumes all responsibility for crop losses resulting from misapplication of this product.

### **11.4 Pastures**

LABELED CROPS: Bahiagrass; Bermudagrass; Bluegrass; Brome; Fescue; Guinea grass; Kikuyu grass; Orchardgrass; Pangola grass; Ryegrass; Timothy; Wheatgrass and any grass (*Gramineae* family) except Corn, Sorghum, Sugarcane and those listed in the "CEREAL AND GRAIN CROPS" section of this label.

TYPES OF APPLICATION: Preplant; Preemergence; Pasture Renovation; Spot Application; Wiper Application; Postemergence Weed Control (broadcast application)

#### **Preplant, Preemergence, Pasture Renovation**

USE INSTRUCTIONS: This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be applied to control perennial pasture species listed on this label prior to re-planting.

RESTRICTIONS: If application rates total 2 quarts of this product per acre or less, no waiting period between application and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait a minimum of 8 weeks after application before grazing or harvesting. Crops listed on this label may be planted into the area at any time; all other crops may be planted 30 days after application.

#### **Spot Application, Wiper Application**

USE INSTRUCTIONS: This product may be applied in pastures as a targeted spot application or over the top of desirable grasses using a wiper applicator to control tall weeds. To achieve maximum performance, remove domestic livestock before application and wait a minimum of 7 days after application before grazing livestock or harvesting for feed. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: For spot application or use with a wiper applicator at rates of 2 quarts per acre or less, this product may be applied over the entire pasture or any portion of it. At rates above 2 quarts per acre, this product may be applied over no more than 10 percent of the total pasture at any one time. Application may be repeated in the same area at 30-day intervals.

#### **Postemergence Weed Control (Broadcast Application)**

USE INSTRUCTIONS: This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and other undesirable vegetation. For selective weed control using broadcast application equipment, apply 8 to 11 fluid ounces of this product per acre in early-spring before desirable perennial grasses break dormancy and initiate green growth. Late-fall application may be made after desirable perennial grasses have reached dormancy.

PRECAUTIONS: Some stunting of perennial grasses will occur if broadcast application is made when plants are not dormant. Higher application rates may be used for hard-to-control weeds; however, higher rates will cause stand reduction.

RESTRICTIONS: No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 2 quarts of this product per acre per year onto pasture grasses except for renovation

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

use as described on this label. If replanting is needed due to severe stand reduction, wait a minimum of 30 days after application before planting any crop not listed on this label.

### 11.5 Rangeland

TYPES OF APPLICATION: Postemergence

USE INSTRUCTIONS: This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangeland. Slight discoloration of the desirable grasses could occur, but will re-green and resume growing under moist soil conditions as effects of this product wear off.

Preventing seed production is critical to the control of invasive annual grassy weeds on rangeland. Yearly application of this product can be used to eliminate viable weed seeds in the soil after they germinate. Delay grazing of the area after application to allow desirable perennials to grow, flower and re-seed the area.

Apply 8 to 11 fluid ounces of this product per acre to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass on rangeland. Apply when most mature brome plants are in early-flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur after spring rains further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Apply this product in the fall in areas where spring moisture is normally limited and fall germination allows for good weed growth and weed seed depletion.

For control of medusahead, apply 11 fluid ounces of this product per acre at the 3-leaf stage. Delaying application beyond this stage will result in reduced or unacceptable control. Controlled burning prior to application can be useful in eliminating the thatch layer produced by slowly decaying culms. Allow new growth to occur before applying this product after a burn. Yearly application of this product is necessary to eliminate the seedbank and allow desirable perennial grasses to re-establish in medusahead-dominated rangeland.

RESTRICTIONS: Do not apply more than 2 quarts of this product per acre per year on rangeland. Do not add ammonium sulfate to the spray solution when applying this product on rangeland grasses. No waiting period between application and feeding or livestock grazing is required.

### 12.0 ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS

ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS CONTAIN A PATENTED GENE THAT PROVIDES TOLERANCE TO GLYPHOSATE, THE ACTIVE INGREDIENT IN THIS PRODUCT. THIS PRODUCT WILL CAUSE SEVERE CROP INJURY OR DESTRUCTION AND YIELD LOSS IF APPLIED TO CROPS THAT ARE NOT GLYPHOSATE TOLERANT. AVOID CONTACT OF THIS PRODUCT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS OR ANY DESIRABLE PLANTS THAT DO NOT CONTAIN A GLYPHOSATE-TOLERANCE GENE, AS SEVERE PLANT INJURY OR DESTRUCTION WILL RESULT. Information on Roundup Ready crops can be obtained from your seed supplier or Bayer CropScience representative. Roundup Ready crops must be purchased from an authorized licensed seed supplier. For information on other brands of glyphosate-tolerant crops, contact the appropriate company representative.

The directions for use in the sections that follow, or those published separately on supplemental labeling for this product, include all applications of this product that may be made onto a specified Roundup Ready or other glyphosate-tolerant crop during the complete cropping season. DO NOT combine these directions for use with the directions for use with the same crops listed in the "ANNUAL AND PERENNIAL CROPS" and "PASTURE GRASSES, FORAGE LEGUMES AND RANGELAND" sections of this label, which are intended for crops that do not contain a glyphosate-tolerance gene.

Roundup Ready seed and the method of selectively controlling weeds in a Roundup Ready crop are protected under several U.S. Patents, including 5,352,605 and 5,633,435. [*This list will be updated at the*

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

*time of printing, if necessary.*] [Alternative text: Roundup Ready seed and the method of selectively controlling weeds in a Roundup Ready crop are protected under several U.S. Patents. For a list of patents, go to [www.monsantotechnology.com](http://www.monsantotechnology.com).]

A license to use Roundup Ready seed must be obtained prior to planting. Bayer CropScience retains ownership of the gene and process technologies, and the Purchaser of the seed receives the right to use the licensed genes and technologies subject to the limited use license conditions. Seed containing a Roundup Ready trait cannot be used for research and demonstration, reverse engineering or in connection with herbicide registration. Progeny seed containing a Roundup Ready trait may not be saved for replanting or transferred to others for replanting. Contact your Authorized Bayer CropScience retailer for information on obtaining a limited use license.

**USE INSTRUCTIONS:** Refer to the “ANNUAL WEEDS RATE SECTION” and “PERENNIAL WEEDS RATE SECTION” of this label for application rates for specific weeds. When applied as directed, this product will control the annual and perennial grasses and broadleaf weeds listed. Observe the maximum application rates and crop stage timings specified for individual Roundup Ready or other glyphosate-tolerant crops in the sections that follow.

**Sprayer Preparation:** It is important that sprayer and mixing equipment be clean and free of pesticide residue before being used to apply this product over the top of glyphosate-tolerant crops. Follow the cleaning procedures specified on the label of the product(s) previously used. **THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.**

**ATTENTION:** AVOID DRIFT. USE EXTREME CARE WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS THAT DO NOT CONTAIN A GLYPHOSATE-TOLERANCE GENE.

**Ground broadcast application** – Apply this product in 5 to 20 gallons of spray solution per acre, unless otherwise directed. Select proper nozzles and spray pressure settings to avoid generating a fine mist. For best performance of this product when using ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

**Aerial application** – Unless otherwise prohibited, all applications of this product described in this section may be made using aerial application equipment, where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label and on all supplemental labeling published separately for this product. Apply this product in 3 to 15 gallons of water per acre. See the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for important information on aerial application and procedures for avoiding spray drift that could cause injury to any vegetation not intended for application. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

See the “MIXING” and “APPLICATION EQUIPMENT AND TECHNIQUES” sections of this label for additional directions and restrictions on the application of this product.

**TANK MIXTURES:** Tank mixtures of this product with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers could result in reduced weed control or crop injury when applied over the top of Roundup Ready or other glyphosate-tolerant crops. Read the label of all products used in the tank mixture prior to use to determine the potential for crop injury. Always read and follow label directions for all products in the tank mixture. Use all products according to rates and timing specified on the product label. Always predetermine the compatibility of tank-mix products together in the carrier by mixing small proportional quantities in advance. Bayer CropScience has not tested this product with all tank-mix product formulations for compatibility, antagonism or performance. To the extent consistent with applicable law, buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not specifically listed on this label or

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

on separate supplemental labeling or Fact Sheets for this product. See the “MIXING” section of this label for more information on tank mixtures.

[*Optional label text:* Unless otherwise directed, nonionic surfactant may be added to the spray solutions of this product for application to Roundup Ready or other glyphosate-tolerant crops. The addition of certain surfactants to a spray solution of this product could result in some crop response including leaf speckling or leaf necrosis due to the surfactant. Refer to the individual Roundup Ready or other glyphosate-tolerant crop sections that follow, or to separate supplemental labeling, for additional precautions or restrictions on the use of surfactants. Refer to the “MIXING” section of this label for additional information on the use of surfactants with this product.]

Ammonium sulfate may be added to spray solutions of this product for application to Roundup Ready and other glyphosate-tolerant crops. Refer to the “MIXING” section of this label for instructions on the use of ammonium sulfate.

The following use directions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, apply this product as a preplant burndown application to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morning glory, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed with multiple germination times, or suppressed (stunted) weeds, might need a second application of this product for complete control. Make second application after some re-growth has occurred and a minimum of 10 days after a previous application of this product.

Application rates of this product specified on this label for hard-to-control weeds, or those specified on separate supplemental labeling for this product, supersede rates in the “ANNUAL WEEDS RATE SECTION” and “PERENNIAL WEEDS RATE SECTION” of this label. Additional information on hard-to-control weeds can be found on Fact Sheets published for this product.

**RESTRICTIONS:** Before using this product, read and follow the Maximum Application Rate information in the “PRODUCT INFORMATION” section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined with the use of any and all other herbicides containing glyphosate, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate. When applying this product as a tank mixture with one or more products, refer to each individual tank-mix product label for restrictions and apply the tank mixture in accordance with the most restrictive statements for each product in the tank.

### 12.1 Alfalfa with Roundup Ready Technology

The directions for use of this product provided in this section are specific to alfalfa containing events J101 and J163, which includes alfalfa with Roundup Ready Technology.

**TYPES OF APPLICATION:** Preplant; At-planting; Preemergence; Postemergence (In-crop)

**USE INSTRUCTIONS:** Refer to the following table for the maximum application rates of this product.

<b>Maximum Application Rates</b>	
Combined total per year for all applications, including Preplant during year of establishment	5.3 quarts per acre
Preplant, At-planting and Preemergence single application	44 fluid ounces per acre
Combined total per year for In-crop applications on newly established and established stands	4.1 quarts per acre

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting Roundup Ready alfalfa.

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied over the top of alfalfa with Roundup Ready Technology (in-crop) from emergence until 5 days prior to cutting. To maximize crop yield and quality potential of the forage and hay, apply this product after weeds have emerged, but before alfalfa growth or re-growth interferes with spray coverage of the target weeds.

Refer to the “ANNUAL WEEDS RATE SECTION” and “PERENNIAL WEEDS RATE SECTION” of this label for application rates for specific weeds. When applied as directed, this product will control the annual and perennial grasses and broadleaf weeds listed. This product will also suppress or control the parasitic weed dodder (*Cuscuta* spp.) in alfalfa with Roundup Ready Technology. More than one application might be necessary for complete control.

PRECAUTIONS: Freezing or near freezing conditions, or large temperature swings, within 5 days after application of this product to alfalfa with Roundup Ready Technology could result in a limited, temporary crop response.

**New Stand Establishment (Seeding Year)** – Due to the biology and breeding constraints of alfalfa, up to 10 percent of the seedlings might not contain a Roundup Ready gene and will not survive after the first application of this product. To eliminate the undesirable effects of stand gaps created by this loss of plants, make a single application of at least 22 fluid ounces of this product per acre at or before the 4-trifoliate growth stage. Refer to the following table for application rates during stand establishment (seeding year).

<b>NEW STAND ESTABLISHMENT (Seeding Year)</b>	
Application Rates	
<b><i>Prior to First Cutting</i></b>	
From emergence up to 4 trifoliate leaves	22 to 44 fluid ounces per acre
From 5 trifoliate leaves up to 5 days before <i>first</i> cutting	Up to 44 fluid ounces per acre
<b><i>After First Cutting</i></b>	
In-crop application, per cutting, up to 5 days before cutting	Up to 44 fluid ounces per acre

TANK MIXTURES: Up to 44 fluid ounces of this product per acre may be applied postemergence (in-crop) over the top of alfalfa with Roundup Ready Technology in the seeding year in a tank-mix with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] after weeds have emerged, but before alfalfa growth or re-growth interferes with spray coverage of the target weeds. Ensure that the product used in the tank-mix is labeled for application postemergence (in-crop) to alfalfa. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:

clethodim; imazamox; imazethapyr; sethoxydim; quizalofop-p-ethyl

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Assure II; Poast; Poast Plus; Pursuit; Raptor; Select 2 EC; Select Max Herbicide with Inside Technology; Warrant]

---

[Optional statement: Pursuit (EPA Reg. No. 241-310; *imazethapyr*) or Raptor (EPA Reg. No. 241-379; *imazamox*) applied to seedling alfalfa could result in a temporary reduction in growth. Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury could result.]

**Established Stands (Non-seeding Year)** – Refer to the following table for directions and application rates for in-crop application to established stands of alfalfa (non-seeding year).

<b>ESTABLISHED STANDS (Non-seeding Year)</b> Application Rates	
In-crop application, per cutting, up to 5 days before cutting	Up to 44 fluid ounces per acre

**TANK MIXTURES:** This product may be applied postemergence (in-crop) over the top of established stands of alfalfa with Roundup Ready Technology in tank mixtures described below according to the growing condition of the crop. Ensure that the product used is labeled for application postemergence (in-crop) to alfalfa. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [Optional text, only required if the product information is not included in the list below: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

**Actively growing alfalfa:** For control of emerged annual grasses and broadleaf weeds when alfalfa is actively growing, this product may be applied at up to 44 fluid ounces per acre in a tank mixture with products containing one or more of the active ingredients listed below [Optional text:, or one or more of the products listed].

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:

clethodim; imazamox; imazethapyr; sethoxydim; quizalofop-p-ethyl

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Assure II; Poast; Poast Plus; Pursuit; Raptor; Select 2 EC; Select Max Herbicide with Inside Technology; Warrant]

---

[Optional statement: Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury could result.]

**Dormant Alfalfa:** For control of emerged annual grasses and broadleaf weeds when alfalfa is dormant, this product may be applied at up to 44 fluid ounces per acre in a tank mixture with products containing one or more of active ingredients listed below [Optional text:, or one or more of the products listed,] when daily temperatures remain above freezing.

---

imazamox; imazethapyr; metribuzin; pronamide; propyzamide

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Kerb 50-W; Kerb SC; Pursuit; Raptor; TriCor 4F; TriCor DF

---

[*Optional statement.* Do not include crop oil concentrate or methylated seed oil in tank mixtures of this product with Pursuit or Raptor as unsatisfactory crop injury could result.]

**PRECAUTIONS:** Where alfalfa with Roundup Ready Technology is grown with a companion or cover crop, or is over-seeded with a second species, in-crop (over-the-top) application of this product will eliminate the non-Roundup Ready (non-glyphosate-tolerant) species.

**RESTRICTIONS:** Do not exceed 44 fluid ounces per acre for any single in-crop application of this product. Sequential applications of this product must be a minimum of 7 days apart. The combined total per year for all in-crop applications in both newly established (seeding year) and established stands (non-seeding year) must not exceed 4.1 quarts (132 fluid ounces) per acre. Do not apply to frozen or snow-covered ground. Remove domestic livestock before application and wait a minimum of 5 days after application before grazing, or cutting or feeding of forage and hay.

### 12.2 Canola with Roundup Ready Technology (Spring Varieties)

The directions for use of this product provided in this section are specific to canola containing Event RT73, which includes canola with Roundup Ready Technology.

For directions for use of this product on canola with TruFlex Technology, refer to that section of this label. DO NOT combine these directions for use on canola with Roundup Ready Technology with the directions for use on canola with TruFlex Technology.

Spring canola with Roundup Ready Technology is defined as those canola varieties with Roundup Ready Technology that are seeded in the spring and harvested in the fall and do not enter a winter dormancy period.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop); Postemergence (In-crop) in Hybrid Seed Production Only

**USE INSTRUCTIONS:** Refer to the following table for the maximum application rates for this product with spring varieties of canola with Roundup Ready Technology.

Maximum Application Rates	
Total for all Preplant, At-Planting, Preemergence applications	44 fluid ounces per acre
Total for all In-crop applications from emergence to 6-leaf stage	22 fluid ounces per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

#### Preplant, At-Planting, Preemergence

**USE INSTRUCTIONS:** This product may be applied before, during or after planting spring canola with Roundup Ready Technology.

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 44 fluid ounces per acre per season.

#### Postemergence (In-crop)

**USE INSTRUCTIONS:** This product may be applied postemergence (in-crop) to spring varieties of canola with Roundup Ready Technology from emergence through the 6-leaf stage of development, unless otherwise directed. Application made during bolting or flowering could result in crop injury and yield loss. To maximize yield potential, eliminate competing weeds early.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**Single Application** – Apply 11 to 16 fluid ounces of this product per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications as this could result in temporary yellowing, delayed flowering, and/or growth reduction. Similar crop injury could result when more than 11 fluid ounces per acre is applied after the 4-leaf stage.

**Sequential Application** – Apply 11 fluid ounces of this product per acre to 1 to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential application works better for control of early emerging annual and perennial weeds, such as Canada thistle and quackgrass, or whenever more than one application is needed for adequate weed control.

**RESTRICTIONS:** No more than two in-crop (over-the-top) broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application must not exceed 22 fluid ounces of this product per acre. Allow a minimum of 60 days between application and canola harvest.

### **Postemergence (In-crop) in Hybrid Seed Production Only**

THIS POSTEMERGENCE APPLICATION IS FOR USE ONLY IN HYBRID CANOLA SEED PRODUCTION OF BOTH SPRING AND WINTER VARIETIES. DO NOT MAKE THIS APPLICATION ON CANOLA GROWN FOR FOOD OR FEED.

This product may be applied at a rate of between 11 and 22 fluid ounces per acre from emergence until pollination is complete or near completion for the control of non-glyphosate-tolerant canola pollen parental line(s) in hybrid canola seed production fields containing both a canola with Roundup Ready Technology line(s) and a non-glyphosate tolerant line(s). Sequential applications may be made for the control of non-glyphosate-tolerant pollen parental lines up to a maximum total application rate of 22 fluid ounces per acre.

**RESTRICTIONS:** Allow a minimum of 5 days between sequential applications. Maximum total application rate of this product for ALL postemergence (in-crop) applications in hybrid canola seed production fields, including application for weed control and control of non-glyphosate-tolerant canola, is 22 fluid ounces per acre.

### **12.3 Canola with Roundup Ready Technology (Winter Varieties)**

The directions for use of this product provided in this section are specific to canola containing Event RT73, which includes canola with Roundup Ready Technology.

For directions for use of this product on canola with TruFlex Technology, refer to that section of this label. DO NOT combine these directions for use on canola with Roundup Ready Technology with the directions for use on canola with TruFlex Technology.

Winter canola with Roundup Ready Technology is defined as those canola with Roundup Ready Technology that are seeded in early-fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop)

**USE INSTRUCTIONS:** Refer to the following table for the maximum application rates of this product with winter varieties of canola with Roundup Ready Technology.

<b>Maximum Application Rates</b>	
Total for all Preplant, At-Planting, Preemergence applications	44 fluid ounces per acre
Total for all In-crop applications from emergence to canopy closure or prior to bolting in the spring	44 fluid ounces per acre

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

### **Preplant, At-Planting, Preemergence**

USE INSTRUCTIONS: This product may be applied up to a maximum application rate of 44 fluid ounces per acre before, during or after planting wither canola with Roundup Ready Technology.

### **Postemergence (In-crop)**

USE INSTRUCTIONS: This product may be applied to winter varieties of canola with Roundup Ready Technology from emergence to canopy closure in the fall and prior to bolting in the spring. Application made during or after bolting could result in crop injury and yield loss. To maximize yield potential, eliminate competing weeds early.

Some weeds with multiple germination times, or suppressed (stunted) weeds, or weeds that have overwintered, might need a sequential application of this product for control. Make the second application after some re-growth has occurred and a minimum of 60 days after the initial application of this product.

**Single Application** – Apply 16 to 22 fluid ounces of this product per acre in the fall when weeds are small and actively growing. Use a higher application rate within this range when weed densities are high, when weeds have overwintered or when weeds become large and well established. Application of more than 16 fluid ounces per acre prior to the 6-leaf stage could result in reduced crop growth in the fall. Avoid spray overlaps as this could result in temporary yellowing and/or growth reduction.

**Sequential Application** – Apply 11 to 22 fluid ounces of this product per acre to 2-leaf or larger canola in the fall, followed by a second application at the same rate and a minimum interval of 60 days, but before bolting in the spring. Sequential application works best for control of early-emerging annual weeds and winter-emerging weeds, such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product will control or suppress most perennial weeds. For some perennial weeds, a sequential application might be needed to reduce competition with the crop.

RESTRICTIONS: No more than two over-the-top broadcast applications may be made from crop emergence up to the onset of bolting, and the total in-crop application must not exceed 44 fluid ounces of this product per acre. Allow a minimum of 60 days between application and harvest of canola grain. No waiting period is required between application and open grazing of livestock.

### **12.4 Canola with TruFlex Technology (Spring Varieties)**

The directions for use of this product provided in this section are specific to canola containing Event MON 88302, which includes canola with TruFlex Technology.

Spring canola with TruFlex Technology is defined as those varieties of canola with TruFlex Technology that are seeded in the spring and harvested in the fall and do not enter a period of winter dormancy.

**The directions for use provided in this section are specific to and may only be used with varieties designated as canola with TruFlex Technology.** Applications described on this label made over the top of canola that is not designated as canola with TruFlex Technology could cause serious crop injury and reduced yields. DO NOT combine these directions for use with those in the “Canola with Roundup Ready Technology (Spring Varieties)” section of this label or with any other directions for use on canola on labeling for this or any other glyphosate-containing product. Drift of this product from an application made to canola with TruFlex Technology onto adjacent fields of canola with Roundup Ready Technology could cause extensive crop injury.

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Postemergence (In-crop) in Hybrid Seed Production Only

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

USE INSTRUCTIONS: Refer to the following table for the maximum application rates of this product with spring varieties of canola with TruFlex Technology.

Maximum Application Rates	
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from emergence through harvest	44 fluid ounces per acre
Total for all In-crop applications from emergence through the 6-leaf stage	44 fluid ounces per acre
Total for all In-crop applications from the 6-leaf stage through first-flower	22 fluid ounces per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: Up to 3.3 quarts of this product may be applied before, during or after planting spring varieties of canola with TruFlex Technology.

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied postemergence (in-crop) to spring varieties of canola with TruFlex Technology from emergence through the first-flower stage of development. To maximize yield potential, eliminate competing weeds early.

For control of Canada thistle and morning glory, apply 44 fluid ounces of this product per acre no later than the 6-leaf stage of canola development. For control of wild buckwheat over 2 inches in size, make sequential applications of 22 fluid ounces followed by 22 fluid ounces of this product per acre. For control of other annual weeds, apply up to 44 fluid ounces of this product per acre no later than the 6-leaf stage or up to 22 fluid ounces after the 6-leaf stage through first-flower.

RESTRICTIONS: No more than two in-crop (over-the-top) broadcast applications may be made from crop emergence through the first-flower stage of canola development and the total in-crop application must not exceed 44 fluid ounces of this product per acre. No more than 22 fluid ounces of this product may be applied in-crop after the 6-leaf stage.

### Postemergence (In-crop) in Hybrid Seed Production Only [This section is optional on the final printed label]

THIS POSTEMERGENCE APPLICATION IS FOR USE ONLY IN HYBRID CANOLA SEED PRODUCTION OF BOTH SPRING AND WINTER VARIETIES. DO NOT MAKE THIS APPLICATION ON CANOLA GROWN FOR FOOD OR FEED.

This product may be applied at a rate of between 11 and 22 fluid ounces per acre from emergence until pollination is complete or near completion for the control of non-glyphosate-tolerant canola pollen parental line(s) in hybrid canola seed production fields containing both a canola with Roundup Ready or TruFlex Technology line(s) and a non-glyphosate-tolerant line(s). Sequential applications may be made for the control of non-glyphosate-tolerant pollen parental lines up to a maximum total application rate of 22 fluid ounces per acre.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**RESTRICTIONS:** Allow a minimum of 5 days between sequential applications. Maximum total application rate of this product for ALL postemergence (in-crop) applications in hybrid canola seed production fields, including application for weed control and control of non-glyphosate-tolerant canola, is 22 fluid ounces per acre.

### 12.5 Field Corn Hybrids with Roundup Ready 2 Technology

The directions for use of this product provided in this section are specific to field corn hybrids containing Event NK603, MON 88017 or MON 87411, which includes Roundup Ready Corn 2, field corn hybrids with Roundup Ready 2 Technology, and field corn seed products displaying the Roundup Ready 2 Technology logo.

The directions for use in this section refer only to FIELD CORN hybrids with Roundup Ready 2 Technology. For directions for use on SWEET CORN hybrids that contain Roundup Ready 2 Technology, see the “Sweet Corn Hybrids with Roundup Ready 2 Technology” section of this label.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop); Spot Application; Preharvest; Post-Harvest; Postemergence (In-crop) for Tassel Control in Roundup Hybridization Systems Only

**USE INSTRUCTIONS:** Refer to the following table for maximum application rates of this product with field corn hybrids with Roundup Ready 2 Technology

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Maximum single In-crop application rate up to 48-inch corn	32 fluid ounces per acre
Total for all In-crop applications from emergence through 48-inch corn	64 fluid ounces per acre
Maximum Preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formed) until 7 days before harvest*	22 fluid ounces per acre

\*See RESTRICTIONS for Preharvest application

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

**PRECAUTIONS:** The use of the in-crop (over-the-top) application rates described in this section on other than field corn hybrids with Roundup Ready 2 Technology could cause crop injury and reduced yields.

#### Preplant, At-Planting, Preemergence

**USE INSTRUCTIONS:** This product may be applied alone or in a tank mixture before, during or after planting field corn hybrids with Roundup Ready 2 Technology.

**TANK MIXTURES:** This product may be tank-mixed with the following products. Apply these tank mixtures in 10 to 20 gallons of water, or 10 to 60 gallons of nitrogen solution, per acre. Ensure that the product used is labeled for application prior to emergence of field corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [Optional text, only required if the product information is not included in the list below: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; dimethenamid; dimethenamid-p; flufenacet; flumetsulam; flumiclorac pentyl ester; fluthiacet-methyl; isoxaflutole; linuron; mesotrione; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; rimsulfuron; saflufenacil; simazine; thiencazabone-methyl

**Axiom DF** (EPA Reg. No. 264-766; *metribuzin, flufenacet*); **Balance Flexx** (EPA Reg. No. 264-1067; *isoxaflutole*); **Capreno** (EPA Reg. No. 264-1063; *tembotrione, thiencazabone-methyl*); **Corvus** (EPA Reg. No. 264-1066; *isoxaflutole, thiencazabone-methyl*); **Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **DiFlexx** (EPA Reg. No. 264-1173; *dicamba*); **DiFlexx DUO** (EPA Reg. No. 264-1184; *dicamba, tembotrione*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*); **Harness MAX** (EPA Reg. No. 524-636; *acetochlor, mesotrione*); **TripleFLEX II** (EPA Reg. No. 524-614; *acetochlor, clopyralid, flumetsulam*)

AAtrex 4L; AAtrex Nine-O; Acuron; Acuron Flexi; Aim EC; Aim EW; Anthem; Anthem ATZ; Anthem Flex; Anthem Maxx; Axiom DF; Balance Flexx; Banvel; Banvel 480; Bicep II MAGNUM; Bicep II MAGNUM FC; Bicep Lite II MAGNUM; Callisto; Capreno; Cinch; Cinch ATZ; Cinch ATZ Lite; Clarity; Corvus; Degree Xtra; DiFlexx; DiFlexx DUO; Distinct; Dual MAGNUM; Dual II MAGNUM; FulTime; FulTime NXT; Guardsman MAX; Harness; Harness MAX; Harness Xtra; Harness Xtra 5.6L; Hornet WDG Broadleaf Blend; Keystone; Keystone LA; Keystone LA NXT; Keystone NXT; Leadoff; Linex 4L; Lorox DF; Marksman; Me-Too-Lachlor II; Outlook; Prowl 3.3 EC; Prowl H2O; Python WDG; Resicore; Resolve DF; Resolve Q; Resolve SG; Resource; Shark EW; Shark H2O; Sharpen Powered by Kixor; Stalwart; Stalwart C; Stalwart Xtra; Surpass EC; Surpass NXT; TopNotch; TripleFLEX II; Verdict powered by Kixor; Zidua]

---

For maximum weed control, make a postemergence (in-crop) application of this product following the use of a preemergence residual product listed above.

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season. Application of 2,4-D or dicamba must be made a minimum of 7 days prior to planting corn.

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied alone or in a tank-mix over the top of field corn hybrids with Roundup Ready 2 Technology from emergence through the V8 stage (8 leaves with collars), or until corn plant height reaches 30 inches (freestanding), whichever comes first, unless otherwise directed. Use drop nozzles for optimum spray coverage and weed control when corn plant height is 24 to 30 inches. When corn plants are 30 to 48 inches tall (freestanding), apply this product using **only** ground application equipment fitted with drop nozzles aligned to avoid spraying into the whorls of the corn plants. Maximum single in-crop application rate of this product up to 48-inch field corn is 32 fluid ounces per acre. Total in-crop application of this product from corn plant emergence through 48 inches in height must not exceed 64 fluid ounces per acre.

When applied as directed, this product will control annual grasses and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Make a postemergence application of 16 to 22 fluid ounces of this product per acre before weeds exceed 4 inches in height (before they become competitive with the crop). Repeat this application before new flushes of weeds exceed 4 inches in height.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below, [*Optional text.*, or one or more of the products listed]. Ensure that the product

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

used is labeled for application postemergence (in-crop) to field corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [Optional text, only required if the product information is not included in the list below: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; flumetsulam; flumiclorac pentyl ester; foramsulfuron; halosulfuron-methyl; iodosulfuron-methyl-sodium; isoxaflutole; mesotrione; nicosulfuron; rimsulfuron; tembotrione; thiencazuron-methyl; thifensulfuron methyl; topramezone

**Capreno** (EPA Reg. No. 264-1063; *tembotrione, thiencazuron-methyl*); **Corvus** (EPA Reg. No. 264-1066; *isoxaflutole, thiencazuron-methyl*); **Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **DiFlexx** (EPA Reg. No. 264-1173; *dicamba*); **DiFlexx DUO** (EPA Reg. No. 264-1184; *dicamba, tembotrione*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness MAX** (EPA Reg. No. 524-636; *acetochlor, mesotrione*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*); **Laudis** (EPA Reg. No. 264-860; *tembotrione*); **TripleFLEX II** (EPA Reg. No. 524-614; *acetochlor, clopyralid, flumetsulam*); **Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Acuron; Acuron Flexi; Aim EC; Aim EW; Banvel; Banvel 480; Basis; Basis Blend; Callisto; Callisto Xtra; Capreno; Clarity; Corvus; Degree Xtra; DiFlexx; DiFlexx DUO; Distinct; Harness; Harness MAX; Harness Xtra; Harness Xtra 5.6L; Hornet WDG Broadleaf Blend; Impact; ImpactZ; Laudis; Marksman; Realm Q; Resicore; Resolve DF; Resolve Q; Resolve SG; Resource; Shark EW; Shark H2O; Status; Stinger; TripleFLEX II; Warrant]

---

**RESTRICTIONS:** Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage or grain.

### Preharvest

**USE INSTRUCTIONS:** Up to 22 fluid ounces of this product per acre may be applied for annual and perennial weed control prior to harvest when kernel fill is complete and the corn is physiologically mature (black layer formed) and grain moisture is 35 percent or less.

**RESTRICTIONS:** A preharvest application may be made only if the combined total of previously applied over-the-top or drop nozzle applications does not exceed 44 fluid ounces of this product per acre. Allow a minimum of 7 days between application and harvest or feeding of corn stover or grain.

### Post-Harvest

**USE INSTRUCTIONS:** This product may be applied for weed control after crop harvest. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in field corn. Read and follow label directions for all products added to the mix.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or the feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

**Postemergence (In-crop) for Tassel Control in Roundup Hybridization Systems Only** [*This section is optional on the final printed label*]

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

THIS APPLICATION IS FOR USE ONLY IN SEED PRODUCTION OF CORN HYBRIDS USING THE ROUNDUP HYBRIDIZATION SYSTEM (RHS). DO NOT MAKE THIS APPLICATION ON CORN GROWN FOR FOOD OR FEED.

The RHS designation indicates that the corn contains Bayer CropScience proprietary gene technology that allows for tassel-only susceptibility to this product. Use of this product on corn hybrids or inbreds that are not designated as RHS or as corn containing Roundup Ready 2 Technology could result in severe crop injury and yield loss.

USE INSTRUCTIONS: This product may be applied at rates of between 11 and 32 fluid ounces per acre as an over-the-top broadcast application for tassel control in RHS-based seed corn production fields from the V8 stage until either the V13 stage or 100 GDU (Growing Degree Units) before flowering.

RESTRICTIONS: Make no more than two applications of this product for tassel control. The maximum total application rate of this product for tassel control is 64 fluid ounces. The maximum combined total amount of this product that may be applied per year for both weed control and tassel control is 5.3 quarts per acre.

### 12.6 Sweet Corn Hybrids with Roundup Ready 2 Technology

The directions for use of this product provided in this section are specific to sweet corn hybrids containing Event MON 88017, which includes Roundup Ready Sweet Corn, sweet corn hybrids with Roundup Ready 2 Technology, and sweet corn seed products displaying the Roundup Ready 2 Technology logo.

The directions for use in this section apply only to SWEET CORN hybrids with Roundup Ready 2 Technology. For directions for use on FIELD CORN hybrids that contain Roundup Ready 2 Technology, see the "Field Corn Hybrids with Roundup Ready 2 Technology" section of this label.

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop)

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with sweet corn hybrids with Roundup Ready 2 Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Maximum single In-crop application rate up to 48-inch sweet corn	44 fluid ounces per acre
Total for all In-crop applications from emergence through 48-inch sweet corn	4.1 quarts per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

PRECAUTIONS: The use of the in-crop (over-the-top) applications described in this section on other than sweet corn hybrids with Roundup Ready 2 Technology could cause crop injury and reduced yields.

#### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank mixture before, during or after planting sweet corn hybrids with Roundup Ready 2 Technology.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] for maximum weed control. Ensure that the product used is labeled for application prior to emergence of sweet corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water or in 10 to 60 gallons of nitrogen solution per acre. [Optional text, only required if the product information is not included in the list below: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list :

acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; dimethenamid-p; mesotrione; metolachlor; s-metolachlor

**Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*)

AAtrex 4L; AAtrex Nine-O; Acuron; Acuron Flexi; Aim EC; Aim EW; Bicep II MAGNUM; Bicep II MAGNUM FC; Bicep Lite II MAGNUM; Cinch; Cinch ATZ; Degree Xtra; Dual MAGNUM; Dual II Magnum; FulTime; Guardsman MAX; Harness; Harness Xtra; Harness Xtra 5.6L; Keystone; Keystone LA; Keystone LA NXT; Keystone NXT; Me-Too-Lachlor II; Outlook; TopNotch]

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### Postemergence (In-crop)

USE INSTRUCTIONS: Apply this product alone or in a tank mixture over the top of sweet corn hybrids with Roundup Ready 2 Technology from emergence through the V8 stage (8 leaves with collars), or until sweet corn plant height reaches 30 inches (freestanding), whichever comes first. Use drop nozzles for optimum spray coverage and weed control when sweet corn plant height is 24 to 30 inches. When sweet corn plants are 30 to 48 inches tall (freestanding), apply this product using **only** ground application equipment fitted with drop nozzles aligned to avoid spraying into the whorls of the sweet corn plants. Avoid spraying if the crop has reached the reproductive stage. Maximum single in-crop application rate of this product up to 48-inch sweet corn is 44 fluid ounces per acre. Total in-crop application of this product from emergence through 48 inches in height must not exceed 4.1 quarts (132 fluid ounces) per acre per growing season.

When applied as directed, this product will control annual grasses and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Apply 16 to 22 fluid ounces of this product per acre before weeds exceed 4 inches in height or before they become competitive with the crop. If new flushes of weeds occur, a sequential application of 16 to 22 fluid ounces per acre may be made before weeds exceed 4 inches in height.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [Optional text:, or one or more of the products listed]. Ensure that the product used is labeled for application postemergence (in-crop) to sweet corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [Optional text, only required if the product information is not included in the list below: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:

**Laudis** (EPA Reg. No. 264-860; *tembotrione*)

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

atrazine; carfentrazone-ethyl; foramsulfuron; tembotrione; topramezone

AAtrex 4L; AAtrex Nine-O; Aim EC; Aim EW; Callisto; Callisto Xtra; Impact; Laudis]

RESTRICTIONS: Allow a minimum of 10 days between in-crop applications of this product. Do not apply atrazine in a tank-mix with this product when sweet corn plants are greater than 12 inches tall. Allow a minimum of 30 days between application of this product and harvest of sweet corn forage or grain.

### 12.7 Field Corn Hybrids with Agrisure GT Technology

The directions for use of this product provided in this section are specific to field corn hybrids containing Event GA21, which includes field corn hybrids with Agrisure GT Technology.

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Spot Application; Preharvest; Post-Harvest

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with field corn hybrids with Agrisure GT Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Maximum single In-crop application rate up to V8 stage or 30-inch corn	22 fluid ounces per acre
Total for all In-crop applications from emergence through V8 stage or 30-inch corn	44 fluid ounces per acre
Maximum Preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formed) until 7 days before harvest*	22 fluid ounces per acre

\*See RESTRICTIONS section for Preharvest application

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

PRECAUTIONS: The use of the in-crop (over-the-top) rates described in this section on other than field corn hybrids with Agrisure GT Technology could cause crop injury and reduced yields.

#### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank mixture before, during or after planting field corn hybrids with Agrisure GT Technology.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below, [*Optional text*:, or one or more of the products listed]. Apply these tank mixtures in 10 to 20 gallons of water, or 10 to 60 gallons of nitrogen solution, per acre. Ensure that the product used is labeled for application prior to emergence of field corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

[*Active ingredients and products added to the final printed label for tank-mixing may be selected*



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

from the following list:

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; dimethenamid; dimethenamid-P; flufenacet; flumetsulam; flumiclorac pentyl ester; fluthiacet-methyl; isoxaflutole; linuron; mesotrione; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; rimsulfuron; saflufenacil; tembotrione; thiencarbazone-methyl

**Axiom DF** (EPA Reg. No. 264-766; *metribuzin, flufenacet*); **Balance Flexx** (EPA Reg. No. 264-1067; *isoxaflutole*); **Capreno** (EPA Reg. No. 264-1063; *tembotrione, thiencarbazone-methyl*); **Corvus** (EPA Reg. No. 264-1066; *isoxaflutole, thiencarbazone-methyl*); **Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **DiFlexx** (EPA Reg. No. 264-1173; *dicamba*); **DiFlexx DUO** (EPA Reg. No. 264-1184; *dicamba, tembotrione*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*); **Harness MAX** (EPA Reg. No. 524-636; *acetochlor, mesotrione*); **TripleFLEX II** (EPA Reg. No. 524-614; *acetochlor, clopyralid, flumetsulam*)

AAtrex 4L; AAtrex Nine-O; Acuron; Acuron Flexi; Aim EC; Aim EW; Anthem; Anthem ATZ; Anthem Flex; Anthem Maxx; Axiom DF; Balance Flexx; Banvel; Banvel 480; Bicep II MAGNUM; Bicep II MAGNUM FC; Bicep Lite II MAGNUM; Callisto; Capreno; Cinch; Cinch ATZ; Cinch ATZ Lite; Clarity; Corvus; Degree Xtra; DiFlexx; DiFlexx DUO; Distinct; Dual MAGNUM; Dual II MAGNUM; FulTime; FulTime NXT; Guardsman MAX; Harness; Harness MAX; Harness Xtra; Harness Xtra 5.6L; Hornet WDG Broadleaf Blend; Keystone; Keystone LA; Keystone LA NXT; Keystone NXT; Leadoff; Linex 4L; Lorox DF; Marksman; Me-Too-Lachlor II; Outlook; Prowl 3.3 EC; Prowl H2O; Python WDG; Resicore; Resolve DF; Resolve Q; Resolve SG; Resource; Shark EW; Shark H2O; Sharpen Powered by Kixor; Stalwart; Stalwart C; Stalwart Xtra; Surpass EC; Surpass NXT; TopNotch; TripleFLEX II; Verdict powered by Kixor; Zidua]

For maximum weed control, make a postemergence (in-crop) application of this product following the use of a preemergence residual product listed above.

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season. Application of 2,4-D or dicamba must be made a minimum of 7 days prior to planting corn.

### **Postemergence (In-crop)**

**USE INSTRUCTIONS:** This product may be applied alone or in a tank-mix over the top of field corn hybrids with Agrisure GT Technology from emergence through the V8 stage (8 leaves with collars), or until corn plant height reaches 30 inches (freestanding), whichever comes first, unless otherwise directed. Maximum single in-crop application rate of this product is 22 fluid ounces per acre. Total in-crop application of this product from corn plant emergence through V8 stage or 30 inches in height must not exceed 44 fluid ounces per acre.

When applied as directed, this product will control annual grasses and broadleaf weeds listed on this label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Make a postemergence application of 16 to 22 fluid ounces of this product per acre before weeds exceed 4 inches in height (before they become competitive with the crop). Repeat this application before new flushes of weeds exceed 4 inches in height.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below, [*Optional text*; or one or more of the products listed]. Ensure that the product used is labeled for application postemergence (in-crop) to field corn. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the*

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

*product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; flumetsulam; flumiclorac pentyl ester; halosulfuron-methyl; isoxaflutole; mesotrione; nicosulfuron; rimsulfuron; tembotrione; thiencazone-methyl; thifensulfuron methyl; topramezone

**Capreno** (EPA Reg. No. 264-1063; *tembotrione, thiencazone-methyl*); **Corvus** (EPA Reg. No. 264-1066; *isoxaflutole, thiencazone-methyl*); **Degree Xtra** (EPA Reg. No. 524-511; *acetochlor, atrazine*); **DiFlexx** (EPA Reg. No. 264-1173; *dicamba*); **DiFlexx DUO** (EPA Reg. No. 264-1184; *dicamba, tembotrione*); **Harness** (EPA Reg. No. 524-473; *acetochlor*); **Harness MAX** (EPA Reg. No. 524-636; *acetochlor, mesotrione*); **Harness Xtra** (EPA Reg. No. 524-480; *acetochlor, atrazine*); **Harness Xtra 5.6L** (EPA Reg. No. 524-485; *acetochlor, atrazine*); **Laudis** (EPA Reg. No. 264-860; *tembotrione*); **TripleFLEX II** (EPA Reg. No. 524-614; *acetochlor, clopyralid, flumetsulam*); **Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Acuron; Acuron Flexi; Aim EC; Aim EW; Banvel; Banvel 480; Basis; Basis Blend; Callisto; Callisto Xtra; Capreno; Clarity; Corvus; Degree Xtra; DiFlexx; DiFlexx DUO; Distinct; Harness; Harness MAX; Harness Xtra; Harness Xtra 5.6L; Hornet WDG Broadleaf Blend; Impact; ImpactZ; Laudis; Marksman; Realm Q; Resicore; Resolve DF; Resolve Q; Resolve SG; Resource; Shark EW; Shark H2O; Status; Stinger; TripleFLEX II; Warrant]

---

**PRECAUTIONS:** In-crop applications of this product made alone or with the addition of other crop chemical products could result in a crop response. Contact your seed trait provider for more information.

**RESTRICTIONS:** Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage or grain.

### **Preharvest**

**USE INSTRUCTIONS:** Up to 22 fluid ounces of this product per acre may be applied for annual and perennial weed control prior to harvest when kernel fill is complete and the corn is physiologically mature (black layer formed) and grain moisture is 35 percent or less.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or feeding of corn stover or grain.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied for weed control after crop harvest. Higher application rates might be needed for control of large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in field corn. Read and follow label directions for all products added to the mix.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or the feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

### **12.8 Cotton with Roundup Ready Technology**

The directions for use of this product provided in this section are specific to cotton containing Event MON 1445, which includes cotton with Roundup Ready Technology.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop); Selective Equipment (In-crop); Preharvest

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with cotton with Roundup Ready Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking to layby	2.5 quarts per acre
Maximum Preharvest application rate	44 fluid ounces per acre
Combined total for all In-crop applications from emergence through harvest	4 quarts per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting Roundup Ready cotton.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and/or dicamba and applied prior to planting only. This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*, or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to the emergence of cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clomazone; diuron; flumioxazin; fluometuron; fomesafen; metolachlor; s-metolachlor; norflurazon; pendimethalin; prometryn; pyriithobac-sodium; saflufenacil

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Caparol 4L; Command 3ME; Cotoran 4L; Cotton Pro; Direx 4L; Dual MAGNUM; Dual II MAGNUM; Karmex DF; Prowl 3.3 EC; Prowl H2O; Reflex; Sharpen Powered by Kixor; Stalwart; Staple LX; Valor SX; Warrant; Warrant Ultra]

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied over the top of cotton with Roundup Ready Technology (in-crop) at rates of up to 22 fluid ounces per acre per application from cracking until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). NO MORE THAN TWO OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT. SEQUENTIAL OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT IN-CROP MUST BE A MINIMUM OF 10 DAYS APART AND

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

COTTON MUST HAVE AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS. Over-the-top application made after the 4-leaf (node) stage of development could result in boll loss, delayed maturity and/or yield loss.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied over the top of cotton with Roundup Ready Technology up to the 4-leaf stage. Ensure that the product used is labeled for application postemergence (in-crop) to cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clethodim; fluazifop-p-butyl; metolachlor; s-metolachlor; monosodium acid methanearsonate; pyriithiobac-sodium; quizalofop-p-ethyl; sethoxydim; trifloxysulfuron-sodium

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Assure II; Dual MAGNUM; Dual II MAGNUM; Envoke; Fusilade DX; MSMA 6 Plus; MSMA 6.6; Poast; Poast Plus; Select 2 EC; Select Max Herbicide with Inside Technology; Stalwart; Staple LX; Warrant]

---

*[Optional text:* Staple LX (EPA Reg. No. 352-613; *pyriithiobac-sodium*) could cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop) to cotton with Roundup Ready Technology.]

*[Optional text:* [Dual MAGNUM (EPA Reg. No. 100-816; *s-metolachlor*), Dual II MAGNUM (EPA Reg. No. 100-818; *s-metolachlor*), and Stalwart (EPA Reg. No. 60063-24; *metolachlor*)] applied over the top of cotton with Roundup Ready Technology could cause leaf injury in the form of necrotic spotting.]

**Salvage Treatment** – Applications of up to 22 fluid ounces of this product per acre either as an over-the-top application or as a post-directed application sprayed higher on the cotton plants and onto targeted weeds may be made after the 4-leaf stage of development only when weeds threaten to cause the loss of the crop.

IN THE STATE OF ARIZONA ONLY, up to 32 fluid ounces of this product may be applied per acre either as an over-the-top application or a post-directed application for salvage treatment.

**NOTE:** SALVAGE TREATMENT WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT MAY BE MADE PER GROWING SEASON.

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all in-crop applications from cracking to layby combined is 2.5 quarts per acre per season. Allow a minimum of 7 days between application and harvest of cotton. DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT (OTHER THAN THOSE CONTAINED IN ANY TANK-MIX PRODUCT) FOR OVER-THE-TOP APPLICATION TO COTTON WITH ROUNDUP READY TECHNOLOGY.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Selective Equipment (In-crop)

**USE INSTRUCTIONS:** This product may be applied using precision post-directed or hooded sprayers at rates of up to 22 fluid ounces per acre per application to cotton with Roundup Ready Technology through layby. At this crop stage, use post-directed application equipment to direct the spray towards the base of the cotton plants, avoiding contact of the herbicide spray with the leaves of the plant. To minimize contact, maintain a low sprayer pressure (less than 30 pounds per square inch) and place nozzles in a low position directing a horizontal spray pattern under the leaves of the cotton plant and onto targeted weeds in the row. For optimal control, apply this product while weeds are small (less than 3 inches in height). See additional use instructions in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] for in-crop application using precision post-directed or hooded sprayers. Ensure that the product used is labeled for application postemergence (in-crop) to cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; diuron; flumioxazin; fluometuron; fomesafen; linuron; metolachlor; monosodium acid methanearsonate; pendimethalin; prometryn; pyriithiobac-sodium; trifloxysulfuron-sodium

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

Aim EC; Aim EW; Caparol 4L; Cotoran 4L; Direx 4L; Envoke; MSMA 6 Plus; MSMA 6.6; Prowl 3.3 EC; Prowl H2O; Staple LX; Valor SX; Warrant; Warrant Ultra]

---

*[Optional text: Staple LX could cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop) to cotton with Roundup Ready Technology.]*

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all in-crop applications from cracking to layby combined is 2.5 quarts per acre per season. Allow a minimum of 7 days between application and harvest of cotton. NO MORE THAN TWO APPLICATIONS OF THIS PRODUCT MAY BE MADE FROM THE 5-LEAF STAGE THROUGH LAYBY. SEQUENTIAL OVER-THE-TOP OR POST-DIRECTED IN-CROP APPLICATIONS OF THIS PRODUCT MUST BE A MINIMUM OF 10 DAYS APART AND COTTON MUST HAVE AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS.

### Preharvest

**USE INSTRUCTIONS:** Up to 44 fluid ounces of this product per acre may be applied for annual and perennial weed control prior to crop harvest after 20 percent boll crack.

**NOTE:** This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

**PRECAUTIONS:** Application of this product for preharvest weed control in cotton grown for seed could result in a reduction in germination or vigor. To the extent consistent with applicable law, buyer and all

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

users are responsible for such loss or damage in connection with the preharvest use of this product on cotton with Roundup Ready Technology grown for seed.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR PREHARVEST APPLICATION TO COTTON WITH ROUNDUP READY TECHNOLOGY.

**ATTENTION:** USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF COTTON WITH ROUNDUP READY TECHNOLOGY. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN ACCORDANCE WITH THE LABEL DIRECTIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

### 12.9 Cotton with Roundup Ready Flex Technology and XtendFlex Technology

The directions for use of this product provided in this section are specific to cotton containing Event MON 88913, which includes cotton with Roundup Ready Flex Technology and cotton with XtendFlex Technology.

**The directions for use of this product provided in this section are specific to and may only be used with varieties designated as cotton with Roundup Ready Flex or XtendFlex Technology.** Applications described in this section made over the top of cotton other than cotton with Roundup Ready Flex or XtendFlex Technology will cause crop injury and reduced yields. DO NOT combine the directions for use in this section with those in the “Cotton with Roundup Ready Technology” or “Cotton with GlyTol Technology” section of this label. Drift of this product from an application made to cotton with Roundup Ready or XtendFlex Technology onto adjacent fields of post 4-leaf (node) cotton with Roundup Ready Technology could cause extensive crop injury, including boll loss, delayed maturity and/or yield loss.

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with cotton with Roundup Ready Flex or XtendFlex Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking to 60 percent open bolls	4 quarts per acre
Total for all In-crop applications between layby and 60 percent open bolls	44 fluid ounces per acre
Total for all In-crop applications from 60 percent open bolls to 7 days prior to harvest	44 fluid ounces per acre
Total for all In-crop applications from emergence through harvest	4 quarts per acre

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

#### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton with Roundup Ready Flex or XtendFlex Technology.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**TANK MIXTURES:** This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clomazone; diuron; flumioxazin; fluometuron; fomesafen; metolachlor; s-metolachlor; norflurazon; pendimethalin; prometryn; pyriithiobac-sodium; saflufenacil

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Caparol 4L; Command 3ME; Cotoran 4L; Cotton Pro; Dawn 4L; Direx; Dual MAGNUM; Dual II MAGNUM; Karmex DF; Prowl 3.3 EC; Prowl H2O; Reflex; Sharpen Powered by Kixor; Stalwart; Staple LX; Valor SX; Warrant; Warrant Ultra]

---

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### **Postemergence (In-crop)**

**USE INSTRUCTIONS:** This product may be applied to control annual grasses and broadleaf weeds listed on this label in cotton with Roundup Ready Flex or XtendFlex Technology. To maximize yield potential, eliminate competing weeds early. Many perennial weeds will be controlled or suppressed with one or more applications of this product. Make an initial application of 22 fluid ounces per acre to control or suppress 1 to 3-inch tall annual grasses and broadleaf weeds. This product may be applied postemergence to cotton with Roundup Ready Flex or XtendFlex Technology using ground application equipment at rates up to 32 fluid ounces per acre per application. In addition to broadcast application, post-directed spray equipment may be used to achieve more thorough weed coverage.

**IN THE STATES OF ARIZONA, NEW MEXICO AND TEXAS (WEST OF I-35) ONLY,** up to 44 fluid ounces of this product per acre may be applied per postemergence application using ground application equipment.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied postemergence (in-crop) over the top of cotton with Roundup Ready Flex or XtendFlex Technology.

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clethodim; fluazifop-p-butyl; metolachlor; s-metolachlor; monosodium acid methanearsonate; pyriithiobac-sodium; quizalofop-p-ethyl; sethoxydim; trifloxysulfuron-sodium

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Assure II; Dual MAGNUM; Dual II MAGNUM; Envoke; Fusilade DX; MSMA 6 Plus; MSMA 6.6; Poast; Poast Plus; Select 2 EC; Select Max Herbicide with Inside Technology; Stalwart; Staple LX; Warrant]

---

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

[*Optional text:* Staple LX (EPA Reg. No. 352-613; *pyrithiobac-sodium*) could cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop) in cotton with Roundup Ready Flex or XtendFlex Technology.]

[*Optional text:* [Dual MAGNUM (EPA Reg. No. 100-816; *s-metolachlor*), Dual II MAGNUM EPA Reg. No. 100-818; *s-metolachlor*], and Stalwart (EPA Reg. No. 60063-24; *metolachlor*)] applied over the top of cotton with Roundup Ready Flex or XtendFlex Technology could cause leaf injury in the form of necrotic spotting.]

This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text:*, or one or more of the products listed,] for in-crop application using precision post-directed or hooded sprayers.

---

[*Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; diuron; flumioxazin; fluometuron; fomesafen; linuron; metolachlor; monosodium acid methanearsonate; pendimethalin; prometryn; pyrithiobac-sodium; trifloxysulfuron-sodium

**Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Aim EC; Aim EW; Caparol 4L; Cotoran 4L; Direx 4L; Envoke; MSMA 6 Plus; MSMA 6.6; Prowl 3.3 EC; Prowl H2O; Staple LX; Valor SX; Warrant; Warrant Ultra]

---

[*Optional text:* Staple LX could cause leaf yellowing and/or leaf crinkling when applied postemergence (in-crop) in cotton with Roundup Ready Flex or XtendFlex Technology.]

Ensure that the product used is labeled for application postemergence (in-crop) to cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

**RESTRICTIONS:** The maximum single, in-crop application rate of this product to cotton with Roundup Ready Flex or XtendFlex Technology using ground application equipment is 32 fluid ounces per acre, except in Arizona, New Mexico and west Texas (west of I-35 only), where up to 44 fluid ounces per acre may be applied in a single application using ground application equipment. **In-crop application rates above 22 fluid ounces per acre made alone or with the addition of other crop chemical products containing surfactant could cause a crop response including leaf speckling or leaf necrosis.** Do not exceed a maximum single, in-crop application rate of 22 fluid ounces of this product per acre when using aerial application equipment, except in Arizona, New Mexico and west Texas (west of I-35 only), where up to 32 fluid ounces may be applied as a single application using aerial application equipment. Between layby and 60 percent open bolls, the maximum combined total application rate of this product is 44 fluid ounces per acre. The combined total for all applications of this product made from crop emergence to 60 percent open bolls must not exceed 4.0 quarts per acre.

**DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR OVER-THE-TOP APPLICATION TO COTTON WITH ROUNDUP READY FLEX OR XTENDFLEX TECHNOLOGY.**

### **Preharvest**

**USE INSTRUCTIONS:** Up to 44 fluid ounces of this product per acre may be applied to cotton with Roundup Ready Flex or XtendFlex Technology for annual and perennial weed control prior to harvest after 60 percent boll crack.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**NOTE:** This product will not enhance the performance of harvest aids when applied to cotton with Roundup Ready Flex or XtendFlex Technology.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest of cotton with Roundup Ready Flex or XtendFlex Technology. DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR PREHARVEST APPLICATION TO COTTON WITH ROUNDUP READY FLEX OR XTENDFLEX TECHNOLOGY.

**ATTENTION:** USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF COTTON WITH ROUNDUP READY FLEX OR XTENDFLEX TECHNOLOGY. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN ACCORDANCE WITH THE LABEL DIRECTIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

### 12.10 Cotton with GlyTol Technology

The directions for use of this product provided in this section are specific to cotton containing Event GHB614, which includes cotton with GlyTol Technology.

**The directions for use of this product provided in this section are specific to and may only be used with varieties designated as cotton with GlyTol Technology.** Applications described in this section made over the top of cotton other than cotton with GlyTol Technology will cause crop injury and reduced yields. DO NOT combine the directions for use in this section with those in the “Cotton with Roundup Ready Technology” or “Cotton with Roundup Ready Flex Technology” section of this label. Drift of this product from an application made to cotton with GlyTol Technology onto adjacent fields of post 4-leaf (node) cotton with Roundup Ready Technology could cause extensive crop injury, including boll loss, delayed maturity and/or yield loss.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest

**USE INSTRUCTIONS:** Refer to the following table for maximum application rates of this product with cotton with GlyTol Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking to 60 percent open bolls	4 quarts per acre
Total for all In-crop applications between layby and 60 percent open bolls	44 fluid ounces per acre
Total for all In-crop applications from 60 percent open bolls to 7 days prior to harvest	44 fluid ounces per acre
Total for all In-crop applications from emergence through harvest	4 quarts per acre

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

#### Preplant, At-Planting, Preemergence

**USE INSTRUCTIONS:** This product may be applied before, during or after planting cotton with GlyTol Technology.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**TANK MIXTURES:** This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may be tank-mixed with one or more products containing the active ingredients listed below, [*Optional text:*, or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clomazone; diuron; flumioxazin; fluometuron; fomesafen; metolachlor; s-metolachlor; norflurazon; pendimethalin; prometryn; pyriithiobac-sodium; saflufenacil

**Warrant Herbicide** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra Herbicide** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

---

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### **Postemergence (In-crop)**

**USE INSTRUCTIONS:** This product may be applied to control annual grasses and broadleaf weeds listed on this label in cotton with GlyTol Technology. To maximize yield potential, eliminate competing weeds early. Many perennial weeds will be controlled or suppressed with one or more applications of this product. Use an initial application rate of 22 fluid ounces per acre to control or suppress 1 to 3-inch tall annual grasses and broadleaf weeds. This product may be applied postemergence to cotton with GlyTol Technology using ground application equipment at rates up to 32 fluid ounces per acre per application. In addition to broadcast application, post-directed spray equipment may be used to achieve more thorough weed coverage.

IN THE STATES OF ARIZONA, NEW MEXICO AND TEXAS (WEST OF I-35) ONLY, up to 44 fluid ounces of this product per acre may be applied per postemergence application using ground application equipment.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text:*, or one or more of the products listed,] and applied postemergence (in-crop) over the top of cotton with GlyTol Technology.

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clethodim; metolachlor; s-metolachlor

**Warrant Herbicide** (EPA Reg. No. 524-591; *acetochlor*)

---

This product may be tank-mixed with products containing one or more of the active ingredients listed below, or one or more of the products listed, for in-crop application using precision post-directed or hooded sprayers.

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; diuron; flumioxazin; fluometuron; fomesafen; linuron; metolachlor; monosodium acid methanearsonate; pendimethalin; prometryn; pyriithiobac-sodium;

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

trifloxysulfuron-sodium

**Warrant Herbicide** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra Herbicide** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

---

Ensure that the product used is labeled for application postemergence (in-crop) to cotton. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix.

**RESTRICTIONS:** The maximum single, in-crop application rate of this product to cotton with GlyTol Technology using ground application equipment is 32 fluid ounces per acre, except in Arizona, New Mexico and west Texas (west of I-35 only), where up to 44 fluid ounces per acre may be applied in a single application using ground application equipment. **In-crop application rates above 22 fluid ounces per acre made alone or with the addition of other crop chemical products containing surfactant could cause a crop response including leaf speckling or leaf necrosis.** Do not exceed a maximum single, in-crop application rate of 22 fluid ounces of this product per acre when using aerial application equipment, except in Arizona, New Mexico and west Texas (west of I-35 only), where up to 32 fluid ounces may be applied as a single application using aerial application equipment. Between layby and 60 percent open bolls, the maximum combined total application rate of this product is 44 fluid ounces per acre. The combined total for all applications of this product made from crop emergence to 60 percent open bolls must not exceed 4.0 quarts per acre.

**DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR OVER-THE-TOP APPLICATION TO COTTON WITH GLYTOL TECHNOLOGY.**

### **Preharvest**

**USE INSTRUCTIONS:** Up to 44 fluid ounces of this product per acre may be applied to cotton with GlyTol Technology for annual and perennial weed control prior to crop harvest after 60 percent boll crack.

**NOTE:** This product will not enhance the performance of harvest aids when applied to cotton with GlyTol Technology.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest of cotton. **DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR PREHARVEST APPLICATION TO COTTON WITH GLYTOL TECHNOLOGY.**

**ATTENTION:** USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF COTTON WITH GLYTOL TECHNOLOGY. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN ACCORDANCE WITH THE LABEL DIRECTIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

### **12.11 Soybean with Roundup Ready Technology**

The directions for use of this product provided in this section are specific to soybean containing Event GTS 40-3-2, which includes soybean with Roundup Ready Technology.

**TYPES OF APPLICATION:** Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest; Post-Harvest

**USE INSTRUCTIONS:** Refer to the following table for maximum application rates of this product with soybean with Roundup Ready Technology.

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking through flowering (R2 stage soybean)	64 fluid ounces per acre
Maximum Preharvest application rate	22 fluid ounces per acre

See the “ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS” section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the “PRODUCT INFORMATION” section of this label for more information on Maximum Application Rates.

**Preplant, At-Planting, Preemergence**

USE INSTRUCTIONS: This product may be applied before, during or after planting soybean with Roundup Ready Technology.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*], or one or more of the products listed, and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; quizalofop-p-ethyl; saflufenacil; sulfentrazone; tribenuron methyl; trifluralin

**Axiom DF** (EPA Reg. No. 264-766; *metribuzin, flufenacet*); **Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor, fomesafen*)

Aim EC; Aim EW; Anthem; Anthem Flex; Anthem Maxx; Assure II; Authority Assist; Authority Elite; Authority First DF; Authority MAXX; Authority MTZ DF; Authority Supreme; Authority XL; Axiom DF; Blanket 4F; Boundary 6.5 EC; Cadet; Canopy; Canopy Blend; Canopy EX; Classic; Cobra; Command 3ME; Dual MAGNUM; Dual II MAGNUM; Fierce; Fierce MTZ; Fierce XLT; FirstRate; Flexstar; Fusion; Linex 4L; Lorox DF; Me-Too-Lachlor; Optill Powered by Kixor; Outlook; Phoenix; Prefix; Prowl 3.3 EC; Prowl H2O; Pursuit; Python WDG; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Sharpen Powered by Kixor; Sonic; Spartan 4F; Treflan 4L; Treflan 4 EC; TriCor 4F; TriCor DF; Valor SX; Valor XLT; Verdict powered by Kixor; Warrant; Warrant Ultra; Zidua; Zidua PRO Powered by Kixor; Zidua SC]

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Postemergence (In-crop)

**USE INSTRUCTIONS:** This product may be used to control annual grasses and broadleaf weeds in soybean with Roundup Ready Technology from emergence (cracking) through flowering (R2 stage soybean). R2 stage soybean ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). Refer to the "ANNUAL WEEDS RATE SECTION" of this label for application rates for specific annual weeds. An initial application of 22 fluid ounces of this product per acre will control or suppress most 2 to 8-inch tall weeds, which are normally found approximately 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be applied up to 44 fluid ounces per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

Application of 22 to 44 fluid ounces of this product per acre (single or multiple applications) will control or suppress perennial weeds, including bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestalk (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For optimal spray coverage and weed control, allow perennial weed species to achieve at least 6 inches of growth before applying this product.

Under adverse growing conditions, including drought, hail or wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product might be necessary to control late flushes of weeds. IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE NEEDED TO CONTROL NEW FLUSHES OF WEEDS IN THE SOYBEAN WITH ROUNDUP READY TECHNOLOGY CROP. To control giant ragweed, apply 22 fluid ounces of this product per acre when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*:, or one or more of the products listed,] and applied postemergence (in-crop) over the top of soybean with Roundup Ready Technology. Ensure that the product used is labeled for application postemergence (in-crop) to soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; acifluorfen; bentazon; chlorimuron-ethyl; clethodim; cloransulam-methyl; fluazifop-p-butyl; flumiclorac pentyl ester; fluthiacet-methyl; fomesafen; imazamox; imazethapyr; lactofen; quizalofop-p-ethyl; sethoxydim; thifensulfuron-methyl

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Arrow 2EC; Assure II; Basagran 5L; Cadet; Classic; Cobra; Extreme; FirstRate; Flexstar; Fusilade DX; Fusion; Harmony SG; Phoenix; Poast; Poast Plus; Pursuit; Raptor; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Synchrony XP; Targa; Ultra Blazer; Warrant; Warrant Ultra]

---

**PRECAUTIONS:** Some tank-mix products can cause visual soybean injury.

**RESTRICTIONS:** The combined total application of this product from crop emergence through harvest must not exceed 64 fluid ounces per acre. The maximum rate for any single in-crop application is 44 fluid ounces

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

per acre. The maximum combined total amount of this product that may be applied during flowering (R2 stage soybean) is 44 fluid ounces per acre.

### Preharvest

USE INSTRUCTIONS: Up to 22 fluid ounces of this product per acre may be applied over the top of soybean with Roundup Ready Technology for weed control prior to harvest after pods have set and lost all green color. Take care to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Allow a minimum of 14 days between application and harvest of soybean grain or feeding of soybean grain, forage or hay.

### Post-Harvest

USE INSTRUCTIONS: This product may be applied for weed control after soybean harvest. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for weed control application after harvest of soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix.

RESTRICTIONS: Application of this product must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

### 12.12 Soybean with Roundup Ready 2 Yield Technology, Roundup Ready 2 Xtend Technology and XtendFlex Technology

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest; Post-Harvest

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with soybean with Roundup Ready 2 Yield, Roundup Ready 2 Xtend and XtendFlex Technologies .

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking through flowering (R2 stage soybean)	64 fluid ounces per acre
Maximum Preharvest application rate	22 fluid ounces per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting soybean with Roundup Ready 2 Yield, Roundup Ready 2 Xtend and XtendFlex Technologies.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or dicamba and applied prior to planting only. This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of soybean. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product*

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

*information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; chlorimuron-ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; quizalofop-p-ethyl; saflufenacil; sulfentrazone; thifensulfuron; tribenuron methyl; trifluralin

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Aim EC; Aim EW; Anthem; Anthem Flex; Anthem Maxx; Assure II; Authority Assist; Authority Elite; Authority First DF; Authority MAXX; Authority MTZ DF; Authority Supreme; Authority XL; Axiom DF; Blanket 4F; Boundary 6.5 EC; Cadet; Canopy; Canopy Blend; Canopy EX; Classic; Cobra; Command 3ME; Dual MAGNUM; Dual II MAGNUM; Enlite; Envive; Fierce; Fierce XLT; FirstRate; Flexstar; Fusion; Linex 4L; Lorox DF; Me-Too-Lachlor; Optill Powered by Kixor; Outlook; Phoenix; Prowl 3.3 EC; Prowl H2O; Pursuit; Python WDG; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Sharpen Powered by Kixor; Sonic; Spartan 4F; Treflan 4L; Treflan 4 EC; TriCor 4F; TriCor DF; Valor SX; Valor XLT; Verdict powered by Kixor; Warrant; Warrant Ultra; Zidua; Zidua PRO Powered by Kixor; Zidua SC]

---

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### **Postemergence (In-crop)**

**USE INSTRUCTIONS:** This product may be used to control annual grasses and broadleaf weeds in soybean with Roundup Ready 2 Yield; Roundup Ready 2 Xtend, and XtendFlex Technologies from emergence (cracking) through flowering (R2 stage soybean). R2 stage soybean ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). Refer to the "ANNUAL WEEDS RATE SECTION" of this label for application rates for specific annual weeds. An initial application of 22 fluid ounces of this product per acre will control or suppress most 2 to 8-inch tall weeds, which are normally found approximately 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be applied up to 44 fluid ounces per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

Application of 22 to 44 fluid ounces of this product per acre (single or multiple applications) will control or suppress perennial weeds, including bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For optimal spray coverage and weed control, allow perennial weed species to achieve at least 6 inches of growth before applying this product.

Under adverse growing conditions, including drought, hail or wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product might be necessary to control late flushes of weeds. IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE SOYBEAN WITH ROUNDUP READY 2 YIELD, ROUNDUP READY 2 XTEND AND XTENDFLEX TECHNOLOGY CROP. To control giant ragweed, apply 22 fluid ounces of this product per acre when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] and applied postemergence (in-crop) over the top of soybean with Roundup Ready 2 Yield, Roundup Ready 2 Xtend and XtendFlex Technology. Ensure that the product used is labeled for application postemergence (in-crop) to soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; acifluorfen; bentazon; chlorimuron ethyl; clethodim; cloransulam-methyl; fluazifop-p-butyl; flumiclorac pentyl ester; fluthiacet-methyl; fomesafen; imazamox; imazethapyr; lactofen; quizalofop-P-ethyl; sethoxydim; thifensulfuron-methyl

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Arrow 2EC; Assure II; Basagran 5L; Cadet; Classic; Cobra; Extreme; FirstRate; Flexstar; Fusilade DX; Fusion; Harmony SG; Phoenix; Poast; Poast Plus; Pursuit; Raptor; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Synchrony XP; Targa; Ultra Blazer; Warrant; Warrant Ultra]

---

**PRECAUTIONS:** Some tank-mix products can cause visual soybean injury.

**RESTRICTIONS:** The combined total application of this product from crop emergence through harvest must not exceed 64 fluid ounces per acre. The maximum rate for any single in-crop application is 44 fluid ounces per acre. The maximum combined total amount of this product that may be applied during flowering (R2 stage soybean) is 44 fluid ounces per acre.

### **Preharvest**

**USE INSTRUCTIONS:** Up to 22 fluid ounces of this product per acre may be applied over the top of soybean with Roundup Ready 2 Yield, Roundup Ready 2 Xtend and XtendFlex Technologies for weed control prior to harvest after pods have set and lost all green color. Take care to avoid excessive seed shatter loss due to ground application equipment.

**RESTRICTIONS:** Allow a minimum of 14 days between application and harvest of soybean grain or feeding of soybean grain, forage or hay.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied for weed control after soybean harvest. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for weed control application after harvest of soybean. Read and follow label directions for all products added to the mix.

**RESTRICTIONS:** Application of this product must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

### **12.13 Soybean with GT27 Technology and Enlist E3 Technology**

The directions for use of this product provided in this section are specific to soybean containing Event FG72 and DAS44406-6, which includes soybean with GT27 Technology and soybean with Enlist E3 Technology.



## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest; Post-Harvest

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with soybean with GT27 and Enlist E3 Technologies.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking through flowering (R2 stage soybean)	64 fluid ounces per acre
Maximum Preharvest application rate	22 fluid ounces per acre

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting soybean with GT27 and Enlist E3 Technologies.

TANK MIXTURES: This product may be tank-mixed with 2,4-D or dicamba [*Alternative text*: Banvel (EPA Reg. No. 66330-276; *dicamba*) or Clarity (EPA Reg. No. 7969-137; *dicamba*)] and applied prior to planting only. This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*., or one or more of the products listed,] and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; quizalofop-p-ethyl; saflufenacil; sulfentrazone; thifensulfuron; tribenuron methyl; trifluralin

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Aim EC; Aim EW; Anthem; Anthem Flex; Anthem Maxx; Assure II; Authority Assist; Authority Elite; Authority First DF; Authority MAXX; Authority MTZ DF; Authority Supreme; Authority XL; Axiom DF; Blanket 4F; Boundary 6.5 EC; Cadet; Canopy; Canopy Blend; Canopy EX; Classic; Cobra; Command 3ME; Dual MAGNUM; Dual II MAGNUM; Enlite; Envive; Fierce; Fierce MTZ; Fierce XLT; FirstRate; Flexstar; Fusion; Linex 4L; Lorox DF; Mauler; Me-Too-Lachlor; Optill Powered by Kixor; Outlook; Phoenix; Prefix; Prowl 3.3 EC; Prowl H2O; Pursuit; Python WDG; Reflex; Resource; Select 2 EC; Select Max Herbicide with Inside Technology; Sharpen Powered by Kixor; Sonic; Spartan 4F; Treflan 4 EC; Treflan 4L; TriCor 4F; TriCor DF; Valor SX; Valor XLT; Verdict powered by Kixor; Warrant; Warrant Ultra; Zidua; Zidua PRO Powered by Kixor; Zidua SC]

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be used to control annual grasses and broadleaf weeds in soybean with GT27 and Enlist E3 Technologies from emergence (cracking) through flowering (R2 stage soybean). R2 stage soybean ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). Refer to the "ANNUAL WEEDS RATE SECTION" of this label for application rates for specific annual weeds.

An initial application of 22 fluid ounces of this product per acre will control or suppress most 2 to 8-inch tall weeds, which are normally found approximately 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be applied up to 44 fluid ounces per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

Application of 22 to 44 fluid ounces of this product per acre (single or multiple applications) will control or suppress perennial weeds, including bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before applying this product.

Under adverse growing conditions, including drought, hail or wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product might be necessary to control late flushes of weeds. IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE SOYBEAN CROP WITH GT27 AND ENLIST E3 TECHNOLOGIES. To control giant ragweed, apply 22 fluid ounces of this product per acre when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*, or one or more of the products listed,] and applied postemergence (in-crop) over the top of soybean with GT27 and Enlist E3 Technologies. Ensure that the product used is labeled for application postemergence (in-crop) to soybean. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clethodim; fomesafen

**Warrant** (EPA Reg. No. 524-591; *acetochlor*); **Warrant Ultra** (EPA Reg. No. 524-620; *acetochlor*, *fomesafen*)

Flexstar; Reflex; Select 2 EC; Select Max Herbicide with Inside Technology; Warrant; Warrant Ultra]

---

PRECAUTIONS: In-crop applications of this product made alone or with the addition of other crop chemical products could result in a crop response. Contact your seed trait provider for more information.

RESTRICTIONS: The combined total application of this product from crop emergence through harvest must not exceed 64 fluid ounces per acre. The maximum rate for any single in-crop application is 44 fluid ounces per acre. The maximum combined total amount of this product that may be applied during flowering (R2 stage soybean) is 44 fluid ounces per acre.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Preharvest

USE INSTRUCTIONS: Up to 22 fluid ounces of this product per acre may be applied to soybean with GT27 and Enlist E3 Technologies for weed control prior to harvest after pods have set and lost all green color. Take care to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Allow a minimum of 14 days between application and harvest of soybean grain or feeding of soybean grain, forage or hay.

### Post-Harvest

USE INSTRUCTIONS: This product may be applied for weed control after soybean harvest. Higher application rates might be needed to control large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for weed control application after harvest of soybean. Read and follow label directions for all products added to the mix.

RESTRICTIONS: Application of this product must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

### 12.14 Sugarbeet with Roundup Ready Technology

The directions for use of this product provided in this section are specific to sugarbeet containing Event H7-1, which includes sugarbeet with Roundup Ready Technology.

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop)

USE INSTRUCTIONS: Refer to the following table for maximum application rates of this product with sugarbeet with Roundup Ready Technology.

Maximum Application Rates	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Maximum single application rate from emergence to 8-leaf stage	32 fluid ounces per acre
Total for all applications made from emergence to 8-leaf stage	56 fluid ounces per acre
Maximum single application rate between 8-leaf stage and canopy closure	22 fluid ounces per acre
Total for all applications made between 8-leaf stage and canopy closure	44 fluid ounces per acre

See the "ROUNDUP READY AND OTHER GLYPHOSATE-TOLERANT CROPS" section of this label for information regarding the use of this product in Roundup Ready and other listed glyphosate-tolerant crops. See the "PRODUCT INFORMATION" section of this label for more information on Maximum Application Rates.

### Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting sugarbeet with Roundup Ready Technology.

TANK MIXTURES: This product may be tank-mixed with Nortron SC (EPA Reg. No. 264-613; *ethofumesate*) and applied prior to crop emergence. Ensure that the product used is labeled for application prior to emergence of sugarbeet. Read and follow label directions for all products added to the mix and follow the most restrictive directions for use and precautionary statements of each.

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### Postemergence (In-crop)

USE INSTRUCTIONS: This product may be applied over the top of sugarbeet with Roundup Ready Technology for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential, eliminate competing weeds early. Up to 4 applications of this product may be made with a minimum of 10 days between each application. This product will control or suppress most perennial weeds. For some perennial weeds, more than one application might be needed to eliminate crop competition throughout the growing season. Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label for application rates for specific weeds.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*: or one or more of the products listed,] and applied postemergence (in-crop) over the top of sugarbeet with Roundup Ready Technology. Ensure that the product used is labeled for application postemergence (in-crop) to sugarbeet. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

[*Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

acetochlor; clethodim; clopyralid; dimethenamid-P; ethofumesate; fluazifop-p-butyl; s-metolachlor; quizalofop-p-ethyl; triflusalufuron methyl

**Nortron SC** (EPA Reg. No. 264-613; *ethofumesate*); **Warrant** (EPA Reg. No. 524-591; *acetochlor*)

Assure II; Dual MAGNUM; Dual II MAGNUM; Fusilade DX; Norton SC; Outlook; Select 2 EC; Select Max Herbicide with Inside Technology; Stinger, Upbeet; Warrant]

---

RESTRICTIONS: The combined total application of this product from crop emergence through harvest must not exceed 100 fluid ounces per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 32 fluid ounces per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 22 fluid ounces per acre. Allow a minimum of 30 days between application and sugarbeet harvest.

### 13.0 FARMSTEAD USE

TYPES OF USES: Farmstead Weed Control; Trim-and-Edge; Greenhouse/Shadehouse; Chemical Mowing; Cut Stump Application; Habitat Management

USE INSTRUCTIONS: Refer to the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label for application rates for specific weeds. When applied as directed, this product will control those annual and perennial grasses and broadleaf weeds on and around a farmstead. Application rates of this product specified in the following sections, or on separate supplemental labeling or Fact Sheets published for this product, for hard-to-control weeds supersede rates in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label.

RESTRICTIONS: Before using this product, read and follow the Maximum Application Rate information in the "PRODUCT INFORMATION" section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined with the use of any and all other herbicides containing glyphosate as the active ingredient, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

### 13.1 Farmstead Weed Control, Trim-and-Edge

USE INSTRUCTIONS: This product may be used to control annual and perennial weeds, woody brush, trees and vines found on any part of the farmstead, including around building foundations and equipment storage areas, along and in fences, in dry ditches and canals, along ditch banks, driveways, farm roads, farmyards, fencerows, parking areas, rangeland, rights-of-way, shelterbelts, storage areas and prior to planting landscape ornamentals.

TANK MIXTURES: This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed], provided that the product used is labeled for these sites and uses. Refer to each individual product label for approved sites and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for all products added to the mix. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; bromacil; chlorosulfuron; dicamba; diuron; imazapic; imazapyr; metsulfuron-methyl; oryzalin; oxadiazon; pendimethalin; proflam; sulfometuron-methyl

Arsenal; Banvel; Banvel 480; Barricade 4L; Barricade 65WG; Clarity; Endurance; Escort XP; Karmex DF; Krovar I DF; Oust XP; Pendulum 3.3 EC; Pendulum 2G; Pendulum Aqua Cap; Plateau; Princep 4L; Princep Caliber 90; Princep Liquid; Ronstar 50 WSP; Ronstar Flo; Ronstar G; Sahara DG; Surflan AS Agricultural; Surflan AS Specialty; Surflan Flex; Surflan XL 2G; Telar XP; Vanquish]

---

For annual weeds, apply 22 fluid ounces of this product per acre when weeds are less than 6 inches tall, 32 fluid ounces when weeds are 6 to 12 inches tall and 44 fluid ounces when weeds are greater than 12 inches tall. For perennial weeds, apply 44 fluid ounces to 3.3 quarts per acre in a tank-mix with one of the products listed here. For application of tank mixtures using a backpack sprayer, handgun or other handheld applicator, see the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of this label for the required concentration of this product in the mix.

### 13.2 Greenhouse/Shadehouse

USE INSTRUCTIONS: This product may be used to control weeds in and around greenhouses and shadehouses.

PRECAUTIONS: Remove desirable vegetation before applying this product inside a greenhouse or shadehouse to prevent unwanted plant injury.

RESTRICTIONS: Turn air circulation fans off when applying this product inside a greenhouse or shadehouse and leave them off until the application solution has dried. Do not use this product inside residential greenhouses.

### 13.3 Chemical Mowing

USE INSTRUCTIONS: This product may be used to suppress growth of perennial grasses listed in this section along farm ditches and on any other part of the farmstead to serve as a substitute for mowing. Apply 4 fluid ounces of this product per acre to suppress Kentucky bluegrass, tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers; 11 fluid ounces to suppress bermudagrass; or 44 fluid ounces to suppress torpedo grass or para grass. Make all applications in 10 to 20 gallons of spray solution per acre.

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

PRECAUTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

### 13.4 Cut Stump Application

TYPES OF USES: Treating brush and tree stumps on any terrestrial site

USE INSTRUCTIONS: This product may be used to control re-growth and re-sprouting of many species of woody brush and trees. Cut the woody brush or tree close to the soil surface and immediately apply a 50 to 100-percent (undiluted) solution of this product to the freshly cut surface using application equipment capable of covering the entire cambium. A delay in application could result in reduced performance. For optimal performance, cut the woody brush or tree during period of active growth and full leaf expansion and apply this product. Some of the species controlled by this method of application of this product are:

Alder	Oak	Reed, giant	Tan oak
Eucalyptus	Pepper, Brazilian	Saltcedar	Willow
Madrone	Pine, Austrian	Sweetgum	

PRECAUTIONS: Do not make a cut stump application when the roots of desirable woody brush or trees might be grafted to the roots of the cut stump. Some sprouts, stems, or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indicator of a shared root system. Whether grafted or shared, injury is likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing a common root system.

### 13.5 Habitat Management

TYPES OF USES: Habitat Restoration and Maintenance; Wildlife Food Plots

#### Habitat Restoration and Maintenance

USE INSTRUCTIONS: This product may be used to control exotic and other undesirable vegetation in habitat management areas. Application may be made to allow recovery of native plant species or prior to planting desirable native species, and for similar broad-spectrum vegetation control in habitat management areas. Spot application may be used to selectively remove unwanted plants for habitat maintenance and enhancement.

#### Wildlife Food Plots

USE INSTRUCTIONS: This product may be used to eliminate annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product or native species may be allowed to repopulate the area naturally. If tillage is needed to prepare a seedbed, wait a minimum of 7 days after application before tilling.

RESTRICTIONS: There are no rotational restrictions for planting any wildlife food species or for allowing native species to repopulate the area following application of this product.

## 14.0 ANNUAL WEEDS RATE SECTION

When applying this product in water carrier volumes of between 16 and 40 gallons per acre using ground application equipment, and between 6 and 15 gallons per acre for aerial application equipment, the following application rates will control the weeds listed in the "ANNUAL WEEDS RATE TABLE" that follows.

- 22 fluid ounces per acre – grasses and broadleaf annual weeds less than 6 inches in height or circumference and vines less than 3 inches in length
- 32 fluid ounces per acre – grasses and broadleaf annual weeds 6 to 12 inches in height or circumference and vines 3 to 6 inches in length

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

- 44 fluid ounces per acre – grasses and broadleaf annual weeds greater than 12 inches in height or circumference and vines greater than 6 inches in length

**WHEN APPLYING IN WATER CARRIER VOLUMES OF BETWEEN 3 AND 15 GALLONS PER ACRE FOR GROUND APPLICATION EQUIPMENT, AND BETWEEN 3 AND 5 GALLONS PER ACRE WITH AERIAL APPLICATION EQUIPMENT, APPLY THIS PRODUCT AT THE RATES SPECIFIED FOR INDIVIDUAL WEEDS IN THE ANNUAL WEEDS RATE TABLE THAT FOLLOWS.**

Apply to actively growing annual weeds. New leaf development indicates active growth.

Annual weeds are often easiest to control when they are small. Application rates greater than those indicated in the following table might be needed to control older, mature (hardened), or otherwise hard-to-control annual weed species, even if they meet the size requirements listed. This product may be applied at rates of up to 44 fluid ounces per acre for hard-to-control annual weeds and where dense weed populations exist. Follow all precautions and restrictions, including maximum application rates and crop stage timings specified in the directions for use on specific crops, including Roundup Ready and other glyphosate-tolerant crops, and use sites listed on this label.

Maximum size refers to the maximum plant height, length of runners for vines, or circumference of rosette plants in inches.

Do not tank-mix this product with soil residual herbicides when applying at these rates, unless otherwise directed.

For control of annual weeds using a handheld controlled droplet applicator (CDA), apply a 20-percent solution of this product (25 to 26 fluid ounces per gallon of spray solution) at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 miles per hour (1 quart per acre). When using a vehicle-mounted CDA, apply the required amount of this product as indicated in the following table, in 2 to 15 gallons of water per acre.

For weeds that have been mowed, grazed or cut, allow re-growth to occur prior to application of this product.

**ANNUAL WEEDS RATE TABLE**

Weed Species	Broadcast Application Rate (fluid ounces per acre)				
	11	16	22	27	32
	Maximum Height/Length (inches)				
Ammannia, purple	3	6	12	-	18
Anoda, spurred	-	2	3	5	8
Barley	18	18 +	-	-	-
Barnyardgrass	-	3	6	7	9
Bassia, fivehook	-	-	6	-	-
Beggarweed, Florida	-	5	8	-	-
Bittercress	12	20	-	-	-
Bluegrass, annual*	10	-	-	-	-
Bluegrass, bulbous	6	-	-	-	-
Brome, downy <sup>1,2</sup>	6	12	-	-	-
Brome, Japanese	6	12	24	-	-
Browntop panicum	6	8	12	-	24
Buckwheat, wild <sup>3</sup>	-	1	2	-	-
Burcucumber	-	6	12	-	18

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Weed Species	Broadcast Application Rate (fluid ounces per acre)				
	11	16	22	27	32
	Maximum Height/Length (inches)				
Buttercup	12	20	-	-	-
Carolina geranium	-	-	4	-	9
Carpetweed	-	6	12	-	-
Cheat <sup>2</sup>	6	20	-	-	-
Chervil	20	-	-	-	-
Chickweed	-	12	18	-	-
Cocklebur	12	18	24	-	36
Copperleaf, hophornbeam	-	2	4	-	6
Copperleaf, Virginia	-	2	4	-	6
Coreopsis, plains	-	6	12	-	18
Corn, volunteer	6	12	20	-	-
Corn speedwell	12	-	-	-	-
Crabgrass	3	6	12	-	-
Crowfootgrass	-	-	6	-	12
Cutleaf evening primrose	-	-	3	-	6
Devilsclaw (unicorn plant)	-	3	6	-	-
Dwarf dandelion	12	-	-	-	-
Eastern mannagrass	8	12	-	-	-
Eclipta	-	4	8	12	-
Fall panicum	4	-	6	-	12
False dandelion	-	20	-	-	-
Falseflax, smallseed	12	-	-	-	-
Fiddleneck	-	6	12	-	-
Field pennycress	6	12	-	-	-
Filaree	-	-	6	-	12
Fleabane, annual	6	20	-	-	-
Fleabane, hairy* ( <i>Conyza bonariensis</i> )	-	-	6	-	10
Fleabane, rough	3	6	12	-	-
Florida pusley	-	-	4	-	6
Foxtail; giant, bristly, yellow	6	12	20	-	-
Foxtail, Carolina	10	-	-	-	-
Foxtail, green	12	-	-	-	-
Goatgrass, jointed	6	12	-	-	-
Goosegrass*	-	3	6	-	12
Grain sorghum (milo)	6	12	20	-	-
Groundcherry	-	3	6	-	9
Groundsel; common, cressleaf	-	6	10	-	-
Hemp sesbania	-	2	4	6	8



I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Weed Species	Broadcast Application Rate (fluid ounces per acre)				
	11	16	22	27	32
	Maximum Height/Length (inches)				
Henbit	-	-	6	-	12
Horseweed / Marestalk* ( <i>Conyza canadensis</i> )	-	6	12	-	18
Itchgrass	6	8	12	-	18
Jimsonweed	-	-	12	-	18
Johnsongrass, seedling*	6	12	18	-	24
Junglerice*	-	3	6	7	9
Knotweed	-	-	6	-	12
Kochia* <sup>4</sup>	-	3 to 6	12	-	-
Lambsquarters	-	6	12	-	20
Little barley	6	12	-	-	-
London rocket	6	-	24	-	-
Mayweed	-	2	6	12	18
Morning glory, annual ( <i>Ipomoea spp</i> )	-	-	3	-	6
Mustard; blue, tansy, tumble, wild	6	12	18	-	-
Nightshade; black, hairy	-	4	6	-	12
Oats	3	6	18	-	-
Pigweed, Palmer*	-	12	18	24	-
Pigweed species*	-	12	18	24	-
Prickly lettuce	-	6	12	-	-
Purslane	-	-	3	-	6
Ragweed; common,* giant*	-	6	12	-	18
Red rice	-	-	4	-	-
Rye, volunteer/cereal <sup>2</sup>	6	18	18 +	-	-
Ryegrass species*	-	-	6	-	12
Sandbur, field	6	12	-	-	-
Sandbur, longspine	6	12	-	-	-
Shattercane	6	12	20	-	-
Shepherd's-purse	6	12	-	-	-
Sicklepod	-	2	4	-	8
Signalgrass, broadleaf	-	3	6	7	9
Smartweed, ladythumb	-	-	6	-	9
Smartweed, Pennsylvania	-	-	6	-	9
Sowthistle, annual	-	-	6	-	12
Spanish needles	-	-	6	-	12
Speedwell, purslane	12	-	-	-	-
Sprangletop	6	12	20	-	-
Spurge; prostrate, spotted	-	6	12	-	-
Spurry, umbrella	6	-	-	-	-

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Weed Species	Broadcast Application Rate (fluid ounces per acre)				
	11	16	22	27	32
	Maximum Height/Length (inches)				
Stinkgrass	-	12	-	-	-
Sunflower*	12	18	-	-	-
Swinecress	-	5	12	-	-
Teaweed/ Prickly sida	-	2	4	-	6
Texas panicum	6	8	12	-	24
Thistle, Russian* <sup>5</sup>	-	6	12	-	-
Velvetleaf	-	-	6	-	12
Virginia pepperweed	-	18	-	-	-
Waterhemp*	-	-	6	-	12
Wheat <sup>2</sup>	6	12	18	-	-
Wheat (overwintered)	-	6	12	-	18
Wild oats	3	6	18	-	-
Wild proso millet	-	6	12	-	18
Witchgrass	-	12	-	-	-
Woolly cupgrass	-	6	12	-	-
Yellow rocket	-	12	20	-	-

- <sup>1</sup> For control of downy brome in no-till systems, apply 16 fluid ounces of this product per acre.
  - <sup>2</sup> Performance of this product is best when application is made before this weed reaches the boot stage of growth.
  - <sup>3</sup> Apply 16 fluid ounces of this product per acre to control wild buckwheat in the cotyledon to 2-leaf stage and 22 fluid ounces per acre to control 2- to 4-leaf wild buckwheat. For better control of wild buckwheat over 2 inches in size, make sequential applications of 22 fluid ounces followed by 22 fluid ounces of this product per acre.
  - <sup>4</sup> Do not apply when kochia is in the button stage.
  - <sup>5</sup> Control of Russian thistle can vary based on environmental conditions and spray coverage. If possible, apply this product in a tank mixture with 2,4-D, as described in the following section, to improve control.
- \* A glyphosate-resistant biotype has been confirmed. For additional information, refer to the “WEED RESISTANCE MANAGEMENT” section of this label. You can also visit [www.weedscience.org](http://www.weedscience.org) on the Internet or contact your Bayer CropScience representative.

**14.1 Annual Weeds—Tank Mixtures with 2,4-D, Dicamba or Tordon 22K**

Control of a broader spectrum of hard-to-control weeds can be achieved by tank-mixing this product with dicamba, 2,4-D, or Tordon 22K (EPA Reg. No. 6271-6; *picloram*). An appropriate rate of these other herbicides combined with the rate of this product specified in the “ANNUAL WEEDS RATE TABLE” will control the following weeds up to the maximum height or length indicated: 6 inches—prickly lettuce, marestail/horseweed, morning glory, kochia (in a tank-mix with dicamba only), wild buckwheat (in a tank-mix with Tordon 22K only); 12 inches—cocklebur, lambsquarters, pigweed, Russian thistle (in a tank-mix with 2,4-D only).

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

At application rates given in the “ANNUAL WEEDS RATE SECTION,” this product will control the following weeds up to a maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf. For better control of these broadleaf weeds, apply this product in a tank-mix with 2,4-D.

Ensure that the product used is labeled for application at the desired site. Follow all precautions and limitations on the tank-mix product label, including any application timing restrictions, soil restrictions, minimum re-cropping intervals and/or crop rotation restrictions. Use according to the more restrictive label requirements. Some crop injury could occur if dicamba or Tordon 22K is applied within 45 days of planting.

### **14.2 Annual Weeds—Handheld Sprayers**

For control of weeds listed in the “ANNUAL WEEDS RATE TABLE,” apply a 0.4-percent solution of this product to weeds less than 6 inches in height or runner length prior to seedhead formation in grasses or bud formation in broadleaf weeds. For control of annual weeds over 6 inches tall, or unless otherwise directed, use a 0.7-percent solution.

For control of annual weeds when using application methods that result in less than complete coverage, apply a 4-percent solution of this product.

### **14.3 Annual Weeds—Tank Mixtures for Fallow and Reduced Tillage Systems**

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

Application of 16 to 20 fluid ounces of this product per acre in a tank mixture with atrazine will control the following weeds: barnyardgrass (requires 20 fluid ounces of this product per acre for control), downy brome, green foxtail, lambsquarters, prickly lettuce, tansy mustard, pigweed, field sandbur, stinkgrass, Russian thistle, volunteer wheat and witchgrass. For control of kochia, apply 16 to 20 fluid ounces of this product per acre in a tank-mix with atrazine and dicamba. Ensure that the atrazine and dicamba products are labeled for the intended use and application site. Follow all precautions and limitations on the tank-mix product label, including any application timing restrictions, soil restrictions, minimum re-cropping intervals and/or crop rotation restrictions.

## **15.0 PERENNIAL WEEDS RATE SECTION**

Apply this product to actively growing perennial weeds. New leaf development indicates active growth. Maximum performance can be obtained when soil moisture is adequate for active weed growth.

If weeds have been recently mowed or tilled, do not apply this product until plants have resumed active growth and have reached the specified stage of growth or sufficient growth has been achieved to allow for good interception of the spray solution. For maximum performance, do not mow, cut, till, burn or disturb vegetation in the application area for a minimum of 7 days after application.

To control hard-to-control perennial weeds, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle, using a handheld sprayer, apply a 1.5-percent solution of this product. For control of perennial weeds when using application methods that result in less than complete coverage, apply a 4-percent solution of this product.

For control of perennial weeds using a handheld controlled droplet applicator (CDA), apply a 20 to 30-percent solution of this product (25 to 38 fluid ounces per gallon of applicator solution) at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mile per hour (2 to 3 quarts per acre). When using a vehicle-mounted CDA, apply the appropriate amount of this product, as indicated in the following rate table, in 2 to 15 gallons of water per acre.

This product has no soil activity and does not control emergence of perennial weeds from seed and dormant underground roots, rhizomes or tubers present in the soil at the time of application. More than one

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

application of this product might be necessary to control weeds regenerating from underground parts or seed, but must be made prior to crop emergence, except where in-crop application is allowed.

Application of this product in the fall must be made before a killing frost.

Unless otherwise directed, allow a minimum of 7 days after application before soil tillage.

**PERENNIAL WEEDS RATE TABLE**

<b>Weed Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Water Volume (gallons/acre)</b>	<b>Handheld Sprayer Concentration (% solution)</b>
<b>Alfalfa</b> Apply after the last cutting in the fall and alfalfa has re-grown to a minimum height of 6 inches. Follow with deep tillage after a minimum of 7 days after application, but before soil freeze-up.	1 – 1.5	3 – 10	1.5%
<b>Alligatorweed</b> For partial control, apply this product when most target plants are in bloom. More than one application will be needed to achieve control.	3	3 – 20	1%
<b>Anise (fennel)<sup>1</sup></b>	–	–	1 - 1.5%
<b>Bahiagrass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Bentgrass</b> For suppression in grass seed production areas using ground application equipment only. Ensure entire crown area has resumed growth prior to application in the fall. Ensure that bentgrass has at least 3 inches of growth before application. Avoid tillage prior to application. Tillage 7 to 10 days after application will help improve bentgrass control.	1	10 – 20	1.5%
<b>Bermudagrass</b> For control, apply 3.3 quarts of this product per acre when bermudagrass is actively growing and seedheads are present. More than one application might be necessary to achieve control. For partial control, apply 64 fluid ounces per acre.	2 – 3.3	3 – 20	1.5%
<b>Bermudagrass, water (knotgrass)</b> Apply 32 fluid ounces of this product in 5 to 10 gallons of water per acre when water bermudagrass is 12 to 18 inches in length. Allow a minimum of 7 days after application before tilling, flushing or flooding the field. For fall application prior to frost, till fallow fields and apply 22 fluid ounces of this product in 5 to 10 gallons of water per acre prior when water bermudagrass is 12 to 18 inches in tall. This product is not registered in California for control of water bermudagrass.	0.7 – 1	5 – 10	1.5%
<b>Bindweed, field</b> Allow maximum bindweed emergence and runner growth before applying this product. Do not apply this product when field bindweed is under drought stress as good soil moisture is necessary for active growth and efficacy of this product. For control, apply 2.5 to 3.3 quarts of this product per acre west of the Mississippi River and 2 to 2.5 quarts per acre east of the Mississippi River when bindweed is at or beyond full bloom. To maximize performance, apply this product in late-summer or fall. Fall application must be made before a killing frost. Also for control, apply 44 fluid ounces of this product plus an appropriate rate of dicamba in 10 to 20 gallons of water per acre. Do not apply this mixture using aerial application equipment. For suppression on irrigated agricultural land, Irrigate at least once to promote active bindweed growth and apply 22 to 44 fluid ounces of this product plus an appropriate rate of 2,4-D in 10 to 20 gallons of	0.4 – 3.3	3 – 20	1.5%

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<p>water per acre using ground application equipment only. Application may be made following harvest or on fallow ground in the fall when bindweed is actively growing, and the majority of runners are 12 inches or more in length.</p> <p>For suppression, apply 11 fluid ounces of this product plus a rate of 2,4-D that will provide suppression of field bindweed in 3 to 10 gallons of water per acre using ground application equipment, or in 3 to 5 gallons of water per acre using aerial application equipment. Application of this tank-mix using aerial equipment is only allowed on fallow fields and in reduced tillage systems. Delay application until maximum bindweed emergence has occurred and vines are 6 to 18 inches in length.</p> <p><b>In California only</b>, apply 22 fluid ounces to 3.3 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions.</p> <p>For suppression on irrigated land where annual tillage is performed, apply 22 fluid ounces of this product in 3 to 10 gallons of water per acre when bindweed has reached a length of 12 inches or more. and wait a minimum of 3 days after application before tillage.</p>			
<b>Bluegrass, Kentucky</b>	0.7 – 1.5	3 – 40	1.5%
<p>Apply 44 fluid ounces of this product in 10 to 40 gallons of water per acre when most plants have reached boot to early-seedhead stage of development. For partial control in pasture or hay crop renovation, apply 22 to 32 fluid ounces of this product in 3 to 10 gallons of water per acre to actively growing target plants when most have reached 4 to 12 inches in height.</p>			
<b>Blueweed, Texas</b>	2 – 3.3	3 – 40	1.5%
<p>Apply 2.5 to 3.3 quarts of this product per acre west of the Mississippi River or 2 to 2.5 quarts per acre east of the Mississippi River when plants are at or beyond full bloom. To maximize performance, apply this product in late-summer or fall. Fall application must be made before a killing frost.</p>			
<b>Brackenfern</b>	2 – 3	3 – 40	1%
<p>Make application to fully expanded fronds that are at least 18 inches long.</p>			
<b>Bromegrass, smooth</b>	0.7 – 1.5	3 – 40	1.5%
<p>Apply 44 fluid ounces of this product in 10 to 40 gallons of water per acre when most target plants have reached boot to early-seedhead stage of development. For partial control in pasture or hay crop renovation, apply 22 to 32 fluid ounces of this product in 3 to 10 gallons of water per acre to actively growing bromegrass when it has reached a height of 4 to 12 inches.</p>			
<b>Bursage, woolly-leaf</b>	–	3 – 20	1.5%
<p>For control, apply 44 fluid ounces of this product per acre in a tank-mix with an appropriate rate of dicamba when plants are producing new active growth that has been initiated by moisture for at least 2 weeks and are at or beyond flowering. For partial control, apply 22 fluid ounces of this product per acre with an appropriate rate of dicamba.</p>			
<b>Canarygrass, reed<sup>2</sup></b>	1.5 – 2	3 – 40	1.5%
<b>Cattail<sup>2</sup></b>	2 – 3.3	3 – 40	1.5%
<b>Clover; red, white<sup>1</sup></b>	2 – 3.3	3 – 20	1.5%
<p>Also for control, apply 11 to 22 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre.</p>			
<b>Cogongrass</b>	2 – 3.3	10 – 40	1.5%
<p>Apply in late-summer or fall when cogongrass is at least 18 inches tall. Due to uneven stages of growth and the dense nature of this vegetation preventing good spray coverage, more than one application might be necessary to achieve control.</p>			

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<b>Dallisgrass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Dandelion<sup>1</sup></b> Also for control, apply 11 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre.	2 – 3.3	3 – 40	1.5%
<b>Dock, curly<sup>1</sup></b> Also for control, apply 11 to 22 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre.	2 – 3.3	3 – 40	1.5%
<b>Dogbane, hemp</b> Apply when most target plants have reached the late-bud to flower stage of development. Allow weeds to re-grow to a mature stage prior to application of this product after crop harvest or mowing. To maximize performance, apply this product in late-summer or fall. For suppression, apply 11 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre using ground application equipment, and in 3 to 5 gallons of water per acre using aerial application equipment. Delay application until maximum emergence of hemp dogbane has occurred.	3	3 – 40	1.5%
<b>Fescue (except tall)<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Fescue, tall</b> Apply 64 fluid ounces of this product per acre when most tall fescue has reached boot to early-seedhead stage of development. For fall application, apply 22 fluid ounces of this product in 3 to 10 gallons of water per acre when plants have 6 to 12 inches of new growth. A sequential application of 11 fluid ounces of this product per acre will improve long-term control and will control seedlings germinating after application in the fall or the following spring.	0.7 – 2	3 – 40	1.5%
<b>Guinea grass</b> Apply when most target plants have reached the 7-leaf stage of growth. Ensure thorough coverage when using a handheld sprayer. In Texas and the ridge of Florida, apply 44 fluid ounces of this product per acre for control. In the flatwoods region of Florida, 64 fluid ounces per acre is needed for control.	1.5 – 2	3 – 40	1%
<b>Hemlock, poison</b> Apply this product using a handheld sprayer with a spray-to-wet technique. Optimum results can be obtained when thoroughly applied to target plants that are at the bud to full-bloom stage of growth.	–	–	1 - 1.5%
<b>Horsenettle<sup>1</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Horseradish</b> Apply when most plants have reached the late-bud to flower stage of growth. For maximum performance, apply in late-summer or fall.	3	3 – 40	1.5%
<b>Iceplant<sup>1</sup></b> Thorough coverage of the target weed with this product will provide best control.	–	–	1.5 - 2%
<b>Jerusalem artichoke<sup>1</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Johnsongrass</b> In annual cropping systems, apply 22 to 44 fluid ounces of this product in 3 to 10 gallons of water per acre. Apply 44 fluid ounces of this product when making application in 10 to 40 gallons of water per acre.	0.4 – 2	3 – 40	1%

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<p>On non-crop sites or in fields where annual tillage is not practiced (no-till), apply 44 to 64 fluid ounces of this product in 10 to 40 gallons of water per acre.</p> <p>For maximum performance, apply this product when most johnsongrass has reached the boot to head stage of development or in the fall prior to frost. Allow a minimum of 7 days after application before tillage. Do not tank-mix with residual herbicides when applying 22 fluid ounces of this product per acre.</p> <p>For burndown of johnsongrass, apply 11 fluid ounces of this product in 3 to 10 gallons of water per acre before plants reach a height of 12 inches and allow a minimum of 3 days after application before tillage.</p> <p>For partial control or suppression, apply a 0.7-percent solution of this product as a spot application when johnsongrass is 12 to 18 inches tall. Ensure that coverage is uniform and complete.</p>			
<b>Kikuyu grass</b>	1.5 – 2	3 – 40	1.5%
<p>Apply when most kikuyu grass is at least 8 inches tall (3- or 4-leaf stage of growth). Allow a minimum of 3 days after application before tillage.</p>			
<b>Knapweed</b>	3	3 – 40	1.5%
<p>Apply when most target plants have reached the late-bud to flower stage of growth. For maximum performance, apply this product in late-summer or fall.</p>			
<b>Lantana</b>	–	–	1%
<p>Apply at or beyond the bloom stage of growth.</p>			
<b>Lespedeza<sup>1</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Milkweed, common</b>	2	3 – 40	1.5%
<p>Apply when most plants have reached the late-bud to flower stage of growth.</p>			
<b>Muhly, wirestem</b>	0.7 – 1.5	3 – 40	1.5%
<p>Apply 22 fluid ounces of this product in 3 to 10 gallons of water per acre, or 44 fluid ounces when applying in 10 to 40 gallons of water per acre, or when applying in pasture, sod, or non-crop areas, when wirestem muhly is at least 8 inches tall. Do not till the soil between harvest and fall application, or in the fall or spring prior to spring application. Allow a minimum of 3 days after application before tillage.</p>			
<b>Mullein, common<sup>1</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Napiergrass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Nightshade, silverleaf</b>	1.5	3 – 10	1.5%
<p>For maximum performance, apply this product when at least 60 percent of the target plants have berries. Fall application must be made before a killing frost.</p>			
<b>Nutsedge; purple, yellow</b>	0.4 – 2	3 – 40	1 – 1.5%
<p>For control of nutsedge plants and immature nutlets, apply 64 fluid ounces of this product per acre or a 1- to 1.5-percent solution when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and will need repeated applications of this product after germination for long-term control.</p> <p>Sequential applications of 22 to 44 fluid ounces of this product in 3 to 10 gallons of water per acre when a majority of the nutsedge plants are in the 3- to 5-leaf stage (less than 6 inches tall) will also provide control. Repeat this application as necessary when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.</p> <p>For partial control of existing plants, apply 11 to 44 fluid ounces of this product in 3 to 40 gallons of water per acre when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat this application as needed to control subsequent emerging plants or re-growth of existing plants.</p>			

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<b>Orchardgrass</b> Apply 44 fluid ounces of this product in 10 to 40 gallons of water per acre when most plants have reached the boot to early-seedhead stage of development. For partial control in pasture or hay crop renovation, apply 22 to 32 fluid ounces of this product in 3 to 10 gallons of water per acre when orchardgrass is actively growing and has reached 4 to 12 inches in height. When going from orchardgrass sod to no-till corn, apply 22 to 32 fluid ounces of this product in 3 to 10 gallons of water per acre to orchardgrass that is a minimum of 12 inches tall for spring application and 6 inches tall for fall application. Allow a minimum of 3 days after application before planting. A sequential application of atrazine will be necessary to achieve optimum results.	0.7 – 1.5	3 – 40	1.5%
<b>Pampas grass</b> Apply this product when pampas grass is at or beyond the boot stage of growth. Thorough coverage of target plants will provide maximum control.	–	–	1 – 1.5%
<b>Para grass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Phragmites</b> For partial control, apply this product in late-summer or fall when target plants are actively growing and in full bloom. Application before or after this stage could result in reduced control. Due to the dense nature of this vegetation (which can prevent good spray coverage) and uneven stages of growth, more than one application might be necessary to achieve control. Visual symptoms of control will be slow to develop.	2 – 3.3	10 – 40	1 – 1.5%
<b>Pokeweed, common</b> Apply to actively growing target plants up to 24 inches tall.	1	3 – 40	1.5%
<b>Quackgrass</b> In annual cropping systems or in pastures and sod fields to be cultivated with deep tillage, apply 22 fluid ounces of this product in 3 to 10 gallons of water per acre or 44 fluid ounces in 10 to 40 gallons of water per acre when quackgrass is 6 to 8 inches in height. Do not tank-mix with residual herbicides when applying at the 22-fluid-ounce rate. Do not till between harvest and fall application, or in the fall or spring prior to spring application. Allow a minimum of 3 days after application before tillage. In pastures or sod fields; use a moldboard plow for maximum control of quackgrass. In pastures, sod fields or non-crop areas where deep tillage will not follow application of this product, apply 44 to 64 fluid ounces in 10 to 40 gallons of water per acre when quackgrass is greater than 8 inches tall.	0.7 – 2	3 – 40	1.5%
<b>Redvine</b> For suppression, make two applications of 16 fluid ounces of this product 7 to 14 days apart, or a single application of 44 fluid ounces, in 5 to 10 gallons of water per acre in late-September or early-October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Apply a minimum of 1 week before a killing frost.	0.5 – 1.5	5 – 10	1.5%
<b>Reed, giant</b> For maximum control, apply this product in late-summer or fall.	–	–	1.5%
<b>Ryegrass, perennial</b> In annual cropping systems, apply 22 to 44 fluid ounces of this product in 3 to 10 gallons of water per acre, or 44 fluid ounces when applying in 10 to 40 gallons of water per acre. On non-crop sites or in fields where annual tillage is not practiced (no-till), apply 44 to 64 fluid ounces of this product in 10 to 40 gallons of water per acre.	0.7 – 2	3 – 40	1%



I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
For maximum performance, apply this product when most target plants have reached the boot to head stage of growth, or in the fall prior to frost. Do not tank-mix with residual herbicides when applying this product at the 22-fluid-ounce-per-acre rate.			
<b>Smartweed, swamp<sup>1</sup></b>	2 – 3.3	3 – 40	1.5%
Also for control, apply 11 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre in late-summer or fall.			
<b>Sowthistle, perennial</b>	1.5 – 2	3 – 40	1.5%
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in late-summer or fall, allow a minimum of 4 weeks for initiation of active growth and rosette development prior to application of this product. Fall application must be made before a killing frost. Allow a minimum of 3 days after application before tillage.			
<b>Spurge, leafy</b>	–	–	1.5%
For suppression, apply 11 fluid ounces of this product in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre in late-summer or fall. If mowing has occurred, delay application until most target plants are 12 inches tall.			
<b>Starthistle, yellow</b>	1.5	10 – 40	1.5%
Best results can be obtained when application is made during the rosette, bolting or early-flower stage.			
<b>Sweet potato, wild</b>	–	–	1.5%
For partial control, apply to plants that are at or beyond the bloom stage of growth. More than one application might be needed.			
<b>Thistle, artichoke</b>	–	–	1.5%
For partial control, apply when target plants are at or beyond the bloom stage of growth. More than one application might be needed.			
<b>Thistle, Canada</b>	1.5 – 2	3 – 40	1.5%
Apply when most target plants are at or beyond the bud stage of development. After harvest, mowing, or tillage in late-summer or fall, allow a minimum of 4 weeks for initiation of active growth and rosette development before applying this product. Fall application must be made before a killing frost.			
For suppression in the spring, apply 22 fluid ounces of this product alone or 11 fluid ounces in a tank-mix with an appropriate rate of 2,4-D in 3 to 10 gallons of water per acre when rosette is a minimum of 6 inches in diameter. Application may be made as long as leaves are still green and plants are actively growing.			
Allow a minimum of 3 days after application before tillage.			
<b>Timothy<sup>2</sup></b>	1.5 – 2	3 – 40	1.5%
<b>Torpedograss</b>	2.5 – 3.3	3 – 40	1.5%
For partial control, apply when most target plants are at or beyond the seedhead stage of development. More than one application will be needed to achieve control. Fall application must be made before frost.			
<b>Trumpetcreeper</b>	1.5	5 – 10	1.5%
For partial control, apply in late-September or October when trumpetcreeper is a minimum of 18 inches tall and has been growing 45 to 60 days since the last tillage operation. Make application a minimum of 7 days before a killing frost.			
<b>Vaseygrass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%
<b>Velvetgrass<sup>2</sup></b>	2 – 3.3	3 – 20	1.5%

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Water Volume</b> (gallons/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<b>Wheatgrass, western<sup>2</sup></b>	1.5 – 2	3 – 40	1.5%

<sup>1</sup> Apply when most plants have reached the early-bud stage of growth.

<sup>2</sup> Apply when most plants have reached the early-heading stage of growth.

**16.0 WOODY BRUSH, TREES AND VINES RATE SECTION**

Apply this product during full leaf expansion, unless otherwise directed. Use a higher application rate or spray solution concentration within a given range to control larger plants or in areas of dense vegetative growth. On vines, use a higher application rate or spray solution concentration for plants that have reached the woody stage.

In most areas, maximum performance of this product on woody brush, trees and vines can be obtained when application is made in late-summer or fall after fruit formation. In arid areas, application of this product in the spring to early-summer when brush and trees are at high moisture content and flowering could provide better results.

Unless otherwise directed, make broadcast applications in 3 to 40 gallons of water per acre. Ensure thorough coverage when using handheld sprayers.

For control of woody brush, trees and vines when using application methods that result in less than complete coverage, apply a 4 to 7-percent solution of this product.

Herbicidal symptoms might not appear prior to frost or senescence following application in the fall.

Allow a minimum of 7 days after application before tillage, mowing or removal of vegetation in the application area. Repeat applications might be necessary to control plants regenerating from underground parts or seed. Some autumn color on undesirable deciduous species is acceptable when applying this product, provided no major leaf drop has occurred. Reduced performance could result if fall application is made following a frost.

**WOODY BRUSH, TREES AND VINES RATE TABLE**

<b>Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Handheld Sprayer Concentration</b> (% solution)
<b>Alder</b>	2 – 3	1%
<b>Ash<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Aspen, quaking</b>	1.5 – 2	1%
<b>Bearmat (Bearclover)<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Beech<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Birch</b>	1.5 – 2	1%
<b>Blackberry</b>	2 – 3	1%

Make application after target plants have reached full leaf maturity. For maximum performance, apply this product in late-summer or fall. Apply a 0.7-percent solution of this product after berries have set or dropped in late-fall. After leaf drop and until a killing frost or as long as stems are green, apply 2 to 2.5 quarts of this product in 10 to 40 gallons of water per acre.

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Sprayer Concentration (% solution)</b>
<b>Blackgum</b>	1.5 – 3.3	1 – 1.5%
<b>Bracken</b>	1.5 – 3.3	1 – 1.5%
<b>Broom; French, Scotch</b>	–	1 – 1.5%
<b>Buckwheat, California<sup>1,2</sup></b>	–	1 – 1.5%
<b>Cascara<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Catsclaw<sup>1</sup></b>	–	1%
<b>Ceanothus<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Chamise<sup>2</sup></b>	–	1%
<b>Cherry; bitter, black, pin</b>	1.5 – 2	1%
<b>Coyote brush</b> Make application when at least 50 percent of the new leaves are fully developed.	–	1 – 1.5%
<b>Dogwood<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Elderberry</b>	1.5 – 2	1%
<b>Elm<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Eucalyptus</b> For control of eucalyptus re-sprouts, apply when re-sprouts are 6 to 12 feet tall. Ensure complete coverage. Application to drought-stressed eucalyptus plants will result in less than optimum results.	–	1.5%
<b>Florida holly (Brazilian Peppertree)<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Gorse<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Hasardia<sup>1,2</sup></b>	–	1 – 1.5%
<b>Hawthorn</b>	1.5 – 2	1%
<b>Hazel</b>	1.5 – 2	1%
<b>Hickory<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Honeysuckle</b>	2 – 3	1%
<b>Hornbeam, American<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Kudzu</b> More than one application might be needed to achieve control.	2.5 – 3.3	1.5%
<b>Locust, black<sup>1</sup></b>	1.5 – 3	1 – 1.5%
<b>Madrone (re-sprouts)<sup>1</sup></b> Apply to re-sprouts that are 3 to 6 feet tall. For maximum performance, apply this product in the spring or early-summer.	–	1.5%
<b>Manzanita<sup>1</sup></b>	1.5 – 3.3	1 – 1.5%
<b>Maple, red</b> Apply a 1-percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 44 to 86 fluid ounces of this product per acre.	1.5 – 3	1%
<b>Maple, sugar</b>	–	1%

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Species	Broadcast Rate (quarts/acre)	Handheld Sprayer Concentration (% solution)
Apply when at least 50 percent of the new leaves are fully developed.		
<b>Monkey flower</b> <sup>1,2</sup>	–	1 – 1.5%
<b>Oak; black, white</b> <sup>1</sup>	1.5 – 3	1 – 1.5%
<b>Oak, post</b>	2 – 3	1%
<b>Oak, northern</b> Apply when at least 50 percent of the new pin leaves are fully developed	–	1%
<b>Oak, southern red</b>	1.5 – 2	1%
<b>Persimmon</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Pine</b>	1.5 – 3.3	1 – 1.5%
<b>Poison ivy/Poison oak</b> More than one application might be needed to achieve control. Application in the fall must be made before leaves lose green color.	2.5 – 3.3	1.5%
<b>Poplar, yellow</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Redbud, eastern</b>	1.5 – 3.3	1 – 1.5%
<b>Rose, multiflora</b> Make application prior to leaf deterioration by leaf-eating insects.	1.5	1%
<b>Russian olive</b> <sup>1</sup>	1.5 – 3.3	1 - 1.5%
<b>Sage, black</b> <sup>2</sup>	–	1%
<b>Sage, white</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Sagebrush, California</b> <sup>2</sup>	–	1%
<b>Salmonberry</b>	1.5 – 2	1%
<b>Saltcedar</b>	1.5 – 3.3	1 – 1.5%
<b>Sassafras</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Sourwood</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Sumac; poison, smooth, winged</b> <sup>1</sup>	1.5 – 3	1 – 1.5%
<b>Sweetgum</b>	1.5 – 2	1%
<b>Swordfern</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Tallowtree, Chinese</b> <sup>2</sup>	–	1%
<b>Tan oak (re-sprouts)</b> <sup>1</sup> Apply to re-sprouts that are less than 6 feet tall. For maximum performance, apply this product in the fall.	–	1.5%
<b>Thimbleberry</b>	1.5 – 2	1%
<b>Tobacco, tree</b> <sup>1</sup>	–	1 – 1.5%
<b>Trumpetcreeper</b>	1.5 – 2	1%
<b>Vine maple</b> <sup>1</sup>	1.5 – 3.3	1 – 1.5%
<b>Virginia creeper</b>	1.5 – 3.3	1 – 1.5%

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Species	Broadcast Rate (quarts/acre)	Handheld Sprayer Concentration (% solution)
Waxmyrtle, southern <sup>1</sup>	1.5 – 3.3	1 – 1.5%
Willow	2 – 3	1%

<sup>1</sup> Partial Control

<sup>2</sup> Thorough coverage of foliage is necessary for maximum performance of this product.

**17.0 TANK-MIX PRODUCT INFORMATION** [*this section printed only if required*]

[*This section is optional and only required if tank-mix product information is not included in the body of the label text. Products not included on the final printed label may be deleted from this list.*]

Brand Name	EPA Reg. No.	Active Ingredient(s)
AAtrex 4L	100-497	atrazine
AAtrex Nine-O	100-585	atrazine
Acuron	100-1466	atrazine, bicyclopyrone, mesotrione, s-metolachlor
Acuron Flexi	100-1568	bicyclopyrone, mesotrione, s-metolachlor
Aim EC	279-3241	carfentrazone-ethyl
Aim EW	279-3242	carfentrazone-ethyl
Alion	264-1106	indaziflam
Anthem	279-3450	fluthiacet-methyl, pyroxasulfone
Anthem ATZ	279-3449	atrazine, fluthiacet-methyl, pyroxasulfone
Anthem Flex	279-3464	carfentrazone-ethyl, pyroxasulfone
Anthem Maxx	279-3468	fluthiacet-methyl, pyroxasulfone
Arrow 2 EC	66222-60	clethodim
Arsenal	241-346	imazapyr
Assure II	352-541	quizalofop-p-ethyl
Authority Assist	279-3330	sulfentrazone, imazethapyr
Authority Elite	279-3442	s-metolachlor, sulfentrazone
Authority First DF	279-3246	sulfentrazone, cloransulam-methyl
Authority MAXX	279-9560	sulfentrazone, chlorimuron ethyl
Authority MTZ DF	279-3340	metribuzin, sulfentrazone
Authority Supreme	279-3601	pyroxasulfone, sulfentrazone
Authority XL	279-3413	sulfentrazone, chlorimuron ethyl
Axiom DF	264-766	metribuzin, flufenacet
Balance Flexx	264-1067	isoxaflutole
Banvel	66330-276	dicamba
Banvel 480	66330-421	dicamba
Barricade 4L	100-1139	proflumicet
Barricade 65WG	100-834	proflumicet
Basagran 5L	7969-112	bentazon
Basis	352-571	rimsulfuron, thifensulfuron methyl

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Brand Name</b>	<b>EPA Reg. No.</b>	<b>Active Ingredient(s)</b>
Basis Blend	352-854	rimsulfuron, thifensulfuron methyl
Bicep II MAGNUM	100-817	atrazine, s-metolachlor
Bicep II MAGNUM FC	100-817	atrazine, s-metolachlor
Bicep Lite II MAGNUM	100-827	atrazine, s-metolachlor
Blanket 4F	82534-5-55467	sulfentrazone
Boundary 6.5 EC	100-1162	metribuzin, s-metolachlor
Cadet	279-3338	fluthiacet-methyl
Callisto	100-1131	mesotrione
Callisto Xtra	100-1359	atrazine, mesotrione
Canopy	352-444	metribuzin, chlorimuron ethyl
Canopy Blend	352-886	metribuzin, chlorimuron ethyl
Canopy EX	352-635	chlorimuron ethyl, tribenuron methyl
Caparol 4L	100-620	prometryn
Capreno	264-1063	tembotrione, thiencazone-methyl
Chateau Herbicide SW	59639-99	flumioxazin
Cinch	352-625	s-metolachlor
Cinch ATZ	352-624	atrazine, s-metolachlor
Cinch ATZ Lite	352-623	atrazine, s-metolachlor
Clarity	7969-137	dicamba
Classic	352-436	chlorimuron ethyl
Cobra	59639-34	lactofen
Command 3ME	279-3158	clomazone
Corvus	264-1066	isoxaflutole, thiencazone-methyl
Cotoran 4L	66222-181	fluometuron
Cotton Pro	66222-15	prometryn
DEF 6	264-730	tribufos
Degree Xtra	524-511	acetochlor, atrazine
Devrinol 2-XT Selective	70506-301	napropamide
Devrinol 50-DF Selective	70506-36	napropamide
Devrinol 50-DF Ornamental	70506-38	napropamide
Devrinol DF-XT Selective	70506-301	napropamide
Devrinol DF-XT Ornamental	70506-38	napropamide
DiFlexx	264-1173	dicamba
DiFlexx DUO	264-1184	dicamba, tembotrione
Direx 4L	66222-54	diuron
Distinct	7969-150	dicamba, diflufenzopyr
Dri-Clean	228-260	2,4-D
Dropp	264-700	thidiazuron
Dual II MAGNUM	100-818	s-metolachlor
Dual MAGNUM	100-816	s-metolachlor
Endurance	100-834	prodiamine

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Brand Name</b>	<b>EPA Reg. No.</b>	<b>Active Ingredient(s)</b>
Enlite	352-757	chlorimuron ethyl, flumioxazin, thifensulfuron
Envive	352-756	chlorimuron ethyl, flumioxazin, thifensulfuron
Envoke	100-1132	trifloxysulfuron-sodium
Escort XP	432-1549	metsulfuron methyl
Extreme	241-405	glyphosate, imazethapyr
Fierce	59639-193	flumioxazin, pyroxasulfone
Fierce MTZ	59639-236	flumioxazin, pyroxasulfone, metribuzin
Fierce XLT	59639-194	chlorimuron ethyl, flumioxazin, pyroxasulfone
FirstRate	62719-275	cloransulam-methyl
Flexstar	100-1101	fomesafen
Folex 6 EC	5481-504	tribufos
FulTime	62719-371	acetochlor, atrazine
FulTime NXT	62719-668	acetochlor, atrazine
Fusilade DX	100-1070	fluazifop-p-butyl
Fusilade II Turf & Ornamental	100-1084	fluazifop-p-butyl
Fusion	100-1059	fluazifop-p-butyl, fenoxaprop-p-ethyl
Ginstar EC	264-634	thidiazuron, diuron
Goal 2XL	62719-424	oxyfluorfen
GoalTender	62719-447	oxyfluorfen
Guardsman MAX	7969-192	dimethenamid-p, atrazine
Harmony SG	352-633 279-9595	thifensulfuron methyl
Harness	524-473	acetochlor
Harness MAX	524-636	acetochlor, mesotrione
Harness Xtra	524-480	acetochlor, atrazine
Harness Xtra 5.6L	524-485	acetochlor, atrazine
Hornet WDG Broadleaf Blend	62719-315	clopyralid, flumetsulam
Impact	5481-524	topramezone
ImpactZ	5481-612	topramezone, atrazine
Karmex DF	66222-51	diuron
Kerb 50-W	62719-397	pronamide
Kerb SC	62719-578	pronamide
Keystone	62719-368	acetochlor, atrazine
Keystone LA	62719-479	acetochlor, atrazine
Keystone LA NXT	62719-670	acetochlor, atrazine
Keystone NXT	62719-671	acetochlor, atrazine
Krovar 1 DF	5481-635	bromacil, diuron
Laudis	264-860	tembotrione
Leadoff	352-853	rimsulfuron, thifensulfuron methyl
Linex 4L	61842-21	linuron

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Brand Name</b>	<b>EPA Reg. No.</b>	<b>Active Ingredient(s)</b>
Lorox DF	61842-23	linuron
Marksman	7969-136	dicamba, atrazine
Matrix FNV	352-671	rimsulfuron
Matrix SG	352-768	rimsulfuron
Mauler	70506-68-59639	metribuzin
Me-Too-Lachlor	19713-548	metolachlor
Me-Too-Lachlor II	19713-549	metolachlor
MSMA 6 Plus	19713-42	MSMA
MSMA 6.6	19713-41	MSMA
Nortron SC	264-613	ethofumesate
Optill Powered by Kixor	7969-280	imazethapyr, saflufenacil
Orchard Master Broadleaf	2217-703	2,4-D
Orchard Master CA Broadleaf	2217-703	2,4-D
Oust XP	432-1552	sulfometuron methyl
Outlook	7969-156	dimethenamid-P
Pendulum 2G	241-375	pendimethalin
Pendulum 3.3 EC	241-341	pendimethalin
Pendulum Aqua Cap	241-416	pendimethalin
Phoenix	59639-118	lactofen
Pindar GT	62719-611	oxyfluorfen, penoxsulam
Plateau	241-365	imazapic
Poast	7969-58	sethoxydim
Poast Plus	7969-88	sethoxydim
Prefix	100-1268	s-metolachlor, fomesafen
Princep 4L	100-526	simazine
Princep Caliber 90	100-603	simazine
Princep Liquid	100-526	simazine
Prowl 3.3 EC	241-337	pendimethalin
Prowl H2O	241-418	pendimethalin
Pursuit	241-310	imazethapyr
Python WDG	62719-277	flumetsulam
Raptor	241-379	imazamox
Realm Q	352-837	mesotrione, rimsulfuron
Reflex	100-993	fomesafen
Rely 280	264-829	glufosinate-ammonium
Resicore	62719-693	acetochlor, clopyralid, mesotrione
Resolve DF	352-556	rimsulfuron
Resolve Q	352-777	rimsulfuron, thifensulfuron methyl
Resolve SG	352-748	rimsulfuron
Resource	59639-82	flumiclorac pentyl ester



I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

<b>Brand Name</b>	<b>EPA Reg. No.</b>	<b>Active Ingredient(s)</b>
Ronstar 50 WSP	432-893	oxadiazon
Ronstar Flo	432-1465	oxadiazon
Ronstar G	432-886	oxadiazon
Sahara DG	241-372	imazapyr, diuron
Select 2 EC	59639-3	clethodim
Select Max Herbicide with Inside Technology	59639-132	clethodim
Shark EW	279-3242	carfentrazone-ethyl
Shark H2O	279-3194	carfentrazone-ethyl
Sharpen Powered by Kixor	7969-278	saflufenacil
Sonic	62719-680	sulfentrazone, cloransulam-methyl
Spartan 4F	279-3220	sulfentrazone
Stalwart	60063-24	metolachlor
Stalwart C	60063-22	metolachlor
Stalwart Xtra	60063-23	atrazine, metolachlor
Staple LX	352-613	pyrithiobac-sodium
Status	7969-242	dicamba, diflufenzopyr
Stinger	62719-73	clopyralid
Surflan AS Agricultural	70506-43	oryzalin
Surflan AS Specialty	70506-44	oryzalin
Surflan Flex	70506-308	oryzalin
Surflan Flex T&O	70506-46	oryzalin
Surflan XL 2G	70506-45	benefin, oryzalin
Surpass EC	62719-367	acetochlor
Surpass NXT	62719-672	acetochlor
Synchrony XP	352-648	thifensulfuron, chlorimuron ethyl
Targa	33906-9-81880	quizalofop-p-ethyl
Telar XP	432-1561	chlorsulfuron
TopNotch	62719-369	acetochlor
Treevix Powered by Kixor	7969-276	saflufenacil
Treflan 4 EC	5905-532	trifluralin
Treflan 4L	34704-853	trifluralin
TriCor 4F	70506-68	metribuzin
TriCor DF	70506-103	metribuzin
TripleFLEX II	524-614	acetochlor, clopyralid, flumetsulam
Ultra Blazer	70506-60	acifluorfen
Upbeet	279-9584	triflurosulfuron methyl
Valor SX	59639-99	flumioxazin
Valor XLT	59639-117	flumioxazin, chlorimuron ethyl
Vanquish	228-397	dicamba
Venue	71711-25	pyraflufen ethyl

## I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Brand Name	EPA Reg. No.	Active Ingredient(s)
Verdict Powered by Kixor	7969-279	dimethenamid-P, saflufenacil
Visor Broadcrop	89167-19-89391	metolachlor
Warrant	524-591	acetochlor
Warrant Ultra	524-620	acetochlor, fomesafen
Zidua	7969-338	pyroxasulfone
Zidua Pro Powered by Kixor	7969-365	saflufenacil, imazethapyr, pyroxasulfone
Zidua SC	7969-374	pyroxasulfone

### 18.0 LIMIT OF WARRANTY AND LIABILITY

Bayer CropScience ("Company") warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label ("Directions") and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

*[Optional statement for agricultural-use products, if applicable: For in-crop (over-the-top) uses on Roundup Ready and other glyphosate-tolerant crops, to the extent consistent with applicable law, crop safety and weed control performance are not warranted by Bayer CropScience when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.]*

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

I. DIRECTIONS FOR USE WITH FOOD AND FEED CROPS

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

---

[Alion; CROPSHIELD and Design; Degree, DiFlex, Drop, Harness, Honcho K6 and Design, Laudis, Nortron, Rely, Roundup, Roundup Original, Roundup Original 2K and Design, Roundup Original MAX and Design, Roundup PowerMAX and Design, Roundup Ready, Roundup Ready 2 Yield, Roundup Ready PLUS and Design, Roundup UltraMAX, Roundup WeatherMAX and Design, [ALTERNATE BRAND NAME and DESIGN], Rowel, RT Master II and Design, Transorb and Design, TripleFLEX, Trisorb and Design, TruFlex, VaporGrip, Warrant, Xtend, XtendFlex, and XtendMAX] are [registered] trademarks of Bayer Group.

[Optional text, if applicable: All other trademarks are the property of their respective owners.]

---

For **MEDICAL** and **TRANSPORTATION** Emergency **ONLY** Call 24 hours a day 1-800-334-7577.

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2987)

---

©[Year]

Packed for [Alternative text: Produced for] [Alternative text: Manufactured for]:

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI, 63167 USA

[Insert print plate number] [Insert barcode]

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

GLYPHOSATE

GROUP

9

HERBICIDE

**[INSERT BRAND NAME]** [Logo]

### Complete Directions for Use

EPA Reg. No. 524-539

A [Optional text: complete] broad-spectrum postemergence herbicide for industrial, turf, ornamental, forestry, roadside, utility rights-of-way, [Optional text, if applicable: select crop,] and other listed terrestrial weed control.

(For a complete list of terrestrial use sites, see the Directions for Use section of this label.)

[Optional label text: FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC AT (800) 424-9300]

[Optional label text: For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 hours a day 1-800-334-7577]

[Optional label text: For **PRODUCT USE** information Call 1-866-99BAYER (1-866-992-2937)]

[Optional statement for limited-geography product distribution: This product is not registered in all states.]

[Additional optional statements for limited-geography product distribution: For control of annual and perennial weeds in [list states and or county-level information, as appropriate, where product is registered and distributed]. [Optional text: \*County Distribution:] [Optional text: see inside for details.] [Optional text: In [list states] this product is distributed in the counties listed below:] [list counties by state]

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [Optional text: GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, [Optional text, if applicable: EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® CROPS,] AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

[Optional label statement: Roundup® – Powerful Performance at a Practical Price]

[Optional label statement: Roundup Ready PLUS™ – Weed Management Solutions (Logo)]

[Optional label statement: TrueBlue Advantage – Proven – Reliable – Supported (Logo)]

[Optional label statement: A member of the Roundup Family of Herbicides by Bayer CropScience]

[Optional label statement: A member of the Roundup Family of Agricultural Herbicides by Bayer CropScience]

[Optional label statement, if applicable: See attached labeling [Alternative text: See inside] for Complete Directions for Use [Optional text: in English and Spanish].

[Optional label statement for container labels with attached labeling: See Complete Directions for Use attached to this label [Alternative text: See attached label booklet] for complete [Optional text, as applicable: Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use [and] Limit of Warranty and Liability] [Optional text: in English and Spanish]

Read the entire label before using this product.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Use only according to label directions.

Not all products listed on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement [*Optional text, if applicable: at the end of this labeling*] [*Alternative optional text, if applicable: on side panel*] [*Alternative optional text, if statement not included on the container label: at the end of the label before buying or using. If terms are not acceptable, return at once unopened.*]

[*Insert net contents, as appropriate: NET [insert appropriate value or leave blank line for refillable container] GAL [or other appropriate unit of measure] [Alternative text: NET CONTENTS] [Alternative label statement for transport vehicles only: NET [Optional: CONTENTS]: See Bill of Lading*]

LOT [*Insert Lot number or blank space*] [*Alternative label statement for transport vehicles only: LOT: See Bill of Lading*]

[*Optional text for container labels: ATTACH COMPLETE DIRECTIONS FOR USE HERE*]

[*Optional text for container labels with attached Directions for Use labeling that will be hidden behind the attached labeling and only become visible if the labeling is removed: COMPLETE DIRECTIONS FOR USE HAVE BEEN REMOVED*]

### CONTENTS

<b>1</b>	<b>1.0</b>	<b>INGREDIENTS</b>
<b>2</b>	<b>2.0</b>	<b>IMPORTANT PHONE NUMBERS</b>
<b>3</b>	<b>3.0</b>	<b>PRECAUTIONARY STATEMENTS</b>
	3.1	Hazards to Humans and Domestic Animals
	3.2	Environmental Hazards
	3.3	Physical or Chemical Hazards
<b>4</b>	<b>4.0</b>	<b>STORAGE AND DISPOSAL</b>
<b>5</b>	<b>5.0</b>	<b>PRODUCT INFORMATION</b>
<b>6</b>	<b>6.0</b>	<b>WEED RESISTANCE MANAGEMENT</b>
	6.1	Weed Management Practices
	6.2	Management of Glyphosate-Resistant Biotypes
<b>7</b>	<b>7.0</b>	<b>MIXING</b>
	7.1	Mixing with Water
	7.2	Tank Mixtures
	7.3	Tank-Mixing Procedure
	7.4	Mixing Spray Solution Concentrations
	7.5	Surfactants [ <i>This section optional in the final printed label</i> ]
	7.6	Ammonium Sulfate [ <i>This section optional in final printed label</i> ]
	7.7	Colorants and Dyes
	7.8	Drift Reduction Additives
<b>8</b>	<b>8.0</b>	<b>APPLICATION EQUIPMENT AND TECHNIQUES</b>
	8.1	Spray Drift Management
	8.2	Aerial Application Equipment
	8.3	Ground Application Equipment
	8.4	Handheld Sprayers
	8.5	Selective Application Equipment

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

- 8.6 Injection Systems
- 8.7 Controlled Droplet Applicator (CDA)
- 9 9.0 TERRESTRIAL USE SITES**
- 10 10.0 ADDITIONAL SITE MANAGEMENT INFORMATION**
  - 10.1 Commercial, Residential and Recreational Area Management
  - 10.2 Forestry, Hardwood and Christmas Tree Management
  - 10.3 Native and Wildlife Habitat Management
  - 10.4 Ornamental and Production Nursery Management
  - 10.5 Pasture Management
  - 10.6 Railroad Management
  - 10.7 Rangeland Management
  - 10.8 Roadside Management
  - 10.9 Utility Management
- 11 11.0 CROP USES** [*This section is optional; appropriate subsections added at time of printing*]
- 12 12.0 WEEDS CONTROLLED**
  - 12.1 Weed Control, Renovation and Chemical Mowing in Turf
  - 12.2 Annual Weeds
  - 12.3 Perennial Weeds
  - 12.4 Woody Brush, Trees and Vines
- 13 13.0 TANK-MIX PRODUCT INFORMATION** [*this section printed only if required*]
- 14 14.0 LIMIT OF WARRANTY AND LIABILITY**

**1.0 INGREDIENTS**

---

ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its potassium salt..... 48.7%

OTHER INGREDIENTS: ..... 51.3%

100.0%

\*Contains 660 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 5.5 pounds per U.S. gallon, which is equivalent to 540 grams of the acid, glyphosate, per liter or 4.5 pounds per U.S. gallon (39.8% by weight).

---

[Optional label text that will be updated at the time of printing, if necessary: This product is protected by [Optional text: one or more of the following] U.S. Patent No's. 5,668,085 and 6,365,551.

[Optional label text, if applicable: Other patents pending]

[Optional label text, if applicable: No license granted under any non-U.S. patent(s)]

[Option to insert reference to a Patent Website: For a list of patents, if any, covering this product or its use, go to [www.monsantotechnology.com](http://www.monsantotechnology.com)]

---

EPA Est. [*Insert appropriate EPA establishment number: 524-IA-1; 524-LA-1; or Other; or blank space*]

[Optional text: (Establishment Number when entered here supersedes all others)]

[Alternative EPA establishment text: EPA Est. (L) 524-LA-1 or (M) 524-IA-1 Lot number prefix (L) or (M) indicate appropriate establishment number.] [*This lot number relationship with the establishment number can be expanded or changed to include additional or add new producing sites, as appropriate.*]

---

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### 2.0 IMPORTANT PHONE NUMBERS

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day **1-800-334-7577**

For **PRODUCT USE** Information Call **1-866-99BAYER (1-866-992-2937)**

### 3.0 PRECAUTIONARY STATEMENTS

#### 3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children

## CAUTION

Causes moderate eye irritation

Avoid contact with eyes, skin, or clothing

[Optional label statement, if applicable: See inside [Alternative text: See back panel] for additional] [Optional text, as applicable: Precautions [and] First Aid]

FIRST AID	
<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<ul style="list-style-type: none"><li>• Have the product container or labeling with you when calling a poison control center or doctor, or going for treatment.</li><li>• For emergency medical treatment information call toll-free 24 hours a day 1-800-334-7577.</li><li>• This product is identified as <b>[INSERT BRAND NAME]</b>, <b>EPA Registration No. 524-539</b>.</li></ul>	

[Optional text, if applicable: LABEL CONTINUED ON BACK]

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

---

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, socks and shoes, and waterproof gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If there are no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### **User Safety Recommendations:**

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### **3.2 Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.

### **3.3 Physical or Chemical Hazards**

Spray solutions of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder’s torch, lighted cigarette or other ignition source and cause serious personal injury.

---

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or on separately published supplemental labeling. Supplemental labeling for this product can be obtained from your Authorized Bayer CropScience retailer or company representative.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, socks and shoes. waterproof gloves.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

*[Optional label text for container labels with attached labeling: See complete Directions for Use attached to this label for complete [Optional text, as applicable: Agricultural [and Non-Agricultural] Use Requirements of the Worker Protection Standard, Directions for Use [and] Limit of Warranty and Liability] [Optional text, if applicable: in English and Spanish]*

## 4.0 STORAGE AND DISPOSAL

### STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. *[Optional label text, if applicable for separate Directions for Use labeling: See individual container label for additional storage conditions, if any.]*

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** *[Optional label text, if applicable for separate Directions for Use labeling: See label attached to the container for handling and disposal instructions and refilling limitations.]*

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID CONTAINERS OF LESS THAN 1-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill the *[this]* container.

*[Alternative container statement: Nonrefillable container. Do not reuse the [this] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [this] container. Contact your state regulatory agency to determine allowable practices in your state.]*

Triple rinse the *[this]* container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once properly rinsed, some plastic [*Optional text: agricultural*] pesticide containers can be taken to a container collection site or picked up for recycling. [*Alternative container disposal statement: Then offer the container for recycling, if available.*]

[*Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].*]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 2.5-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 1-GALLON, BUT EQUAL TO OR LESS THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse the [*this*] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [*this*] container. Contact your state regulatory agency to determine allowable practices in your state.

[*Alternative container statement: Nonrefillable container. Do not reuse or refill the [this] container.*]

Triple rinse or pressure rinse (or equivalent) the [*this*] container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [*Optional text: agricultural*] pesticide containers can be taken to a container collection site or picked up for recycling. [*Alternative container disposal statement: Then offer the container for recycling, if available.*]

[*Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].*]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[*Insert UN packaging certification, if applicable*]

[*Optional packaging text: Pull at scored cutouts for easier opening*]

[*Optional packaging text: THIS SIDE UP*] [*Optional graphic of two arrows pointing upward*]

[*Optional packaging text: Liquid*] [*Optional graphic of a liquid droplet*]

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 30-GALLON CONTAINER AND OTHER NONREFILLABLE CONTAINERS OF GREATER THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill the [this] container.

[Alternative container statement: Nonrefillable container. Do not reuse the [this] container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the [this] container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse or pressure rinse (or equivalent) the container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank. [Optional label text: For containers not equipped with pumping systems,] Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth for 30 seconds, ensuring at least one complete revolution. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or mix-tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

[Alternative or additional triple rinsing instructions for large containers equipped with pumping systems: [Optional label text: For containers equipped with pumping systems,] Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.]

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a collection site or picked up for recycling. [Alternative container disposal statement: Then offer the container for recycling, if available.]

[Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the [this] container, if available. If no recycling information is available on the [this] container, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)] to find the nearest recycling location.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].]

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-866-99BAYER (1-866-992-2937)]

[Optional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) [www.schuetz.net/ticket](http://www.schuetz.net/ticket); Schuetz ticket service]

[Optional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only.]

[Optional container disposal statement: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA –

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Clarkston, Michigan – USA]

### [CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR ALL REFILLABLE CONTAINERS, EXCEPT TRANSPORT VEHICLES]

Refillable container. Refill the [this] container with pesticide only. Do not reuse the [this] container for any other purpose.

Cleaning the [this] container before refilling is the responsibility of the refiller. Cleaning the [this] container before final disposal is the responsibility of the person disposing of the container.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix-tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Then offer the container for recycling, if available.

[Optional container disposal statement: Some container manufacturers offer container recycling. See additional information regarding manufacturer recycling programs attached to the [this] container, if available. If no recycling information is available on the [this] container, contact your chemical dealer [Optional text: or Bayer CropScience at 1-866-99BAYER (1-866-992-2937)] to find the nearest recycling location.]

[Optional container disposal statement: IBC EMPTY? – FREE CALL – 1-888-SCHUETZ (1-888-724-8389) [www.schuetz.net/ticket](http://www.schuetz.net/ticket); Schuetz ticket service]

[Optional container disposal statement: FREE IBC PICKUP] [For continental USA and Canada only.]

[Optional container disposal statement: RETURNnet SYSTEM – To return empty IBC's Email or Call – [www.returnnetsystem.com](http://www.returnnetsystem.com) – 1-888-758-SHIP – United States and Canada (1-888-758-7447 – IBCNA – Clarkston, Michigan – USA]

[Optional container disposal statement: To obtain information about recycling refillable containers, contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937)].

[Optional container disposal statement: Return Properly Rinsed Container for Recycling – Call 1-866-99BAYER (1-866-992-2937)]

[Optional container label statements for the CUBE refillable packaging system only:

#### **CUBE** Refillable Delivery System

##### FEATURES INCLUDE:

- Automatic Venting
- Heavy duty one-way 2-inch camloc ball valve with protective shield door
- Complete coated steel protective enclosure
- Durable 4-way plastic pallet

Lift door to access one-way valve]

### [CONTAINER HANDLING AND DISPOSAL STATEMENT FOR ALL TRANSPORT VEHICLES, AS DEFINED IN 40 CFR 156.3]

Emptied container retains vapor and product residue. Observe all precautions stated on this label until the container is cleaned, reconditioned or destroyed. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, and worn-out threads and closures. Clean thoroughly before reuse for transportation of a material of different composition or before retiring this transport vehicle from service.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

[Additional label statement for transport vehicles only: FOR BULK PESTICIDE TRANSPORT ONLY]

[Additional label statement for transport vehicles only: THIS LABEL FOR USE WITH TRANSPORT VEHICLES ONLY]

[STORAGE AND DISPOSAL FOR THE NONREFILLABLE SMART-PAK CONTAINER – A COMPOSITE PACKAGING CONSISTING OF A LIGHTWEIGHT PLASTIC CONTAINER INSIDE RIGID CARDBOARD PACKAGING. ONCE IT IS FILLED, THE LIGHTWEIGHT PLASTIC CONTAINER IS NOT STORED, TRANSPORTED OR SOLD WITHOUT THE OUTER CARDBOARD PACKAGING.]

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. Do not store the [this] container unprotected from the weather, especially precipitation, for extended periods of time. DO NOT REMOVE THE PLASTIC CONTAINER FROM THE PACKAGE UNTIL IT HAS BEEN EMPTIED AND PROPERLY RINSED.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the [this] container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not reuse or refill the [this] container.

Triple rinse or pressure rinse (or equivalent) the inner plastic container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the inner plastic container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into the application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing or collect rinsate for later use or disposal. Insert pressure rinsing nozzle through the bottom of the container, ensuring that you puncture the inner plastic container, and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

After rinsing, open the outer cardboard packaging and remove the inner plastic container.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a collection site or picked up for recycling. Recycle the cardboard separately. [Alternative container disposal statement: Then offer the outer cardboard packaging and inner plastic container for recycling, if available.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or call 1-866-99BAYER (1-866-992-2937)].]

If recycling is not available, dispose of each component in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed inner plastic container and disposing both the outer cardboard packaging and inner plastic container in a sanitary landfill.

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

[The inner plastic container of the Smart-Pak must be labeled with the following information]

[INSERT BRAND NAME] [Logo optional]

ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its potassium salt..... 48.8%

OTHER INGREDIENTS: ..... 51.2%  
100.0%

\*Contains 660 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 5.5 pounds per U.S. gallon, which is equivalent to 540 grams of the acid, glyphosate, per liter or 4.5 pounds per U.S. gallon (39.8% by weight).

Keep out of reach of children

## CAUTION

EPA Reg. No. 524-539

**UNLESS THIS PLASTIC CONTAINER IS EMPTY AND HAS BEEN PROPERLY RINSED FOR DISPOSAL, IT IS NOT TO BE REMOVED FROM ITS OUTER CARDBOARD PACKAGING. FOR MORE INFORMATION, CALL 1-899-99BAYER (1-866-992-2937).**

[Optional Smart-Pak container label statements:

- Protect this package from precipitation until emptied and properly rinsed.
- Use the cap to cut the foil seal.
- Invert cap over foil seal and rotate clockwise to cut the foil seal. Remove foil seal.
- East to Handle
- Uses less plastic than ordinary [Alternative text: typical] jugs
- Produces less plastic waster than ordinary [Alternative text: typical] jugs
- Compact Design
- Space Efficient
- Stackable
- Easy to Use
- Easy to Pour
- Easy to Recycle]

[CONTAINER HANDLING AND DISPOSAL STATEMENT FOR ALL TRANSPORT VEHICLES, AS DEFINED IN 40 CFR 156.3]

Emptied container retains vapor and product residue. Observe all precautions stated on this label until the container is cleaned, reconditioned or destroyed. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, and worn-out threads and closures. Clean thoroughly before reuse for transportation of a material of different composition or before retiring this transport vehicle from service.

[Additional label statement for transport vehicles only: FOR BULK PESTICIDE TRANSPORT ONLY]

[Additional label statement for transport vehicles only: THIS LABEL FOR USE WITH TRANSPORT VEHICLES ONLY]

[STORAGE AND DISPOSAL FOR THE NONREFILLABLE SMART-PAK CONTAINER – A COMPOSITE PACKAGING CONSISTING OF A LIGHTWEIGHT PLASTIC CONTAINER INSIDE RIGID CARDBOARD PACKAGING. ONCE IT IS FILLED, THE LIGHTWEIGHT PLASTIC CONTAINER IS NOT STORED, TRANSPORTED OR SOLD WITHOUT THE OUTER CARDBOARD PACKAGING.]

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. Do not store the [this] container unprotected from the weather, especially precipitation, for extended periods of time. DO NOT REMOVE THE PLASTIC CONTAINER FROM THE PACKAGE UNTIL IT HAS BEEN EMPTIED AND PROPERLY RINSED.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the [this] container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not reuse or refill the [this] container.

Triple rinse or pressure rinse (or equivalent) the inner plastic container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Fill the inner plastic container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into the application equipment or mix-tank, or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix-tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix-tank while rinsing or collect rinsate for later use or disposal. Insert pressure rinsing nozzle through the bottom of the container, ensuring that you puncture the inner plastic container, and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

After rinsing, open the outer cardboard packaging and remove the inner plastic container.

Once properly rinsed, some plastic [Optional text: agricultural] pesticide containers can be taken to a collection site or picked up for recycling. Recycle the cardboard separately. [Alternative container disposal statement: Then offer the outer cardboard packaging and inner plastic container for recycling, if available.]

[Optional container disposal statement: To find the nearest collection site, contact your chemical dealer [Optional text: or call 1-866-99BAYER (1-866-992-2937)].]

If recycling is not available, dispose of each component in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed inner plastic container and disposing both the outer cardboard packaging and inner plastic container in a sanitary landfill.

[The inner plastic container of the Smart-Pak must be labeled with the following information]

[INSERT BRAND NAME] [Logo optional]

ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl) glycine, in the form of its potassium salt..... 48.8%

OTHER INGREDIENTS: ..... 51.2%

100.0%

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

\*Contains 660 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 5.5 pounds per U.S. gallon, which is equivalent to 540 grams of the acid, glyphosate, per liter or 4.5 pounds per U.S. gallon (39.8% by weight).

Keep out of reach of children

### CAUTION

EPA Reg. No. 524-539

**UNLESS THIS PLASTIC CONTAINER IS EMPTY AND HAS BEEN PROPERLY RINSED FOR DISPOSAL, IT IS NOT TO BE REMOVED FROM ITS OUTER CARDBOARD PACKAGING. FOR MORE INFORMATION, CALL 1-899-99BAYER (1-866-992-2937).**

[Optional Smart-Pak container label statements:

- Protect this package from precipitation until emptied and properly rinsed.
- Use the cap to cut the foil seal.
- Invert cap over foil seal and rotate clockwise to cut the foil seal. Remove foil seal.
- Easy to Handle
- Uses less plastic than ordinary [Alternative text: typical] jugs
- Produces less plastic waste than ordinary [Alternative text: typical] jugs
- Compact Design
- Space Efficient
- Stackable
- Easy to Use
- Easy to Pour
- Easy to Recycle]

## 5.0 PRODUCT INFORMATION

**Product Description:** This product is a postemergence, systemic herbicide with no soil residual activity. This product provides broad-spectrum control of many annual and perennial weeds, woody brush, trees and vines. It is formulated as a water-soluble liquid containing surfactant that may be applied using standard and specialized pesticide application equipment after dilution and thorough mixing with water or other carriers according to label directions.

[Optional label text: Do not add [Optional label text: surfactants, additives containing surfactants,] buffering agents or pH adjusting agents to the spray solution when [INSERT BRAND NAME] is the only pesticide being applied unless otherwise directed. See the "MIXING" section of this label for instructions regarding other additives.]

[Optional label text: No additional surfactant in the spray solution is needed. This includes additives containing surfactants, buffering agents or pH adjusting agents when [INSERT BRAND NAME] is the only pesticide used unless otherwise directed.]

**Mechanism of Action:** The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

**No Soil Activity:** This product binds tightly to soil particles and does not provide residual weed control. Weeds must be emerged at the time of application to be controlled by foliar application of this product. Weed seeds in the soil will not be affected by this product and will continue to germinate. Unattached plant rhizomes and rootstocks beneath the soil surface will also not be affected by this product.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

**Stage of Weeds:** Annual weeds are easiest to control when they are small. Performance of this product on most perennial weeds is best when applied at late growth stages approaching maturity. Refer to the “WEEDS CONTROLLED” section of this label for more information on the control of specific weeds.

**Cultural Considerations:** Reduced weed control could result when this product is applied to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to re-grow prior to application. Always use a higher application rate of this product within a given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area. Reduced weed control could also result when this product is applied to weeds that show signs of disease or insect damage, are covered with dust, or are surviving under poor growing conditions.

**Spray Coverage:** For best results with this product, spray coverage must be uniform and complete. Do not spray foliage to the point of runoff.

**Rainfastness:** Rainfall within 4 hours of application could wash this product off of the foliage and a second application might then be needed to achieve adequate weed control. Refer to specific use sections of this label for additional information on the minimum intervals required before re-application of this product.

**Time to Symptoms:** This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of aboveground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds, effects might not be visible for 7 or more days after application. Extremely cool or cloudy weather following application could slow activity of this product and delay development of visual symptoms.

**Maximum Application Rates:** The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowable application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or in a tank mixture, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalents) does not exceed the maximum allowed. See the “INGREDIENTS” section of this label for necessary product information.

Unless otherwise specified on this label, the combined total of all applications of this product on a site must not exceed 7 quarts (8 pounds of glyphosate acid) per acre per year.

Unless otherwise specified on this label in the sections that follow, the maximum single application rate of this product is 7 quarts (8 pounds of glyphosate acid) per acre. Unless otherwise specified on this label, multiple applications of this product made be made onto a site up to the maximum annual application rate of 7 quarts (8 pounds of glyphosate acid) per acre per year. Allow a minimum of 14 days between applications.

The following table provides the glyphosate application rate (pounds of glyphosate acid equivalents per acre) when this product is applied at the application rates indicated (fluid ounces or quarts of this product per acre).

<b>Application Rate of [INSERT BRAND NAME] (amount of product per acre)</b>	<b>Application Rate of Glyphosate Acid Equivalents (ae) (pounds of ae per acre)</b>
11 fluid ounces	0.39
16 fluid ounces	0.56
22 fluid ounces	0.77
36 fluid ounces	1.27

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

44 fluid ounces	1.55
64 fluid ounces	2.25
2.7 quarts	3.0
3.3 quarts	3.7
4.1 quarts	4.6
7 quarts	8

To determine the Glyphosate acid equivalents (pounds of Glyphosate ae per acre) for application rates of **[INSERT BRAND NAME]** not listed here, multiply the application rate (fluid ounces per acre) by 0.0352.

$$\text{Application Rate} \times 0.0352 = \text{Glyphosate (ae) per acre}$$

### 6.0 WEED RESISTANCE MANAGEMENT

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mechanism of action classification system of the Weed Science Society of America. Any weed population can contain plants that are naturally resistant to Group 9 herbicides. Weeds resistant to Group 9 herbicides can be effectively managed by using another herbicide from a different Group (either alone or in a mixture according to label directions), by using other cultural or mechanical methods of weed control, or a combination of the two. Consult your local company representative, state cooperative extension agent, professional consultant or other qualified authority to determine appropriate actions for controlling specific resistant weeds.

#### 6.1 Weed Management Practices

Resistant populations arise when rare individual plants are uncontrolled by a normal dose of a given herbicide under normal environmental conditions. In the absence of other control measures these individuals survive, produce seed, and eventually become the dominant biotype in the field through continuous selection. The best means of reducing this selection is to use diverse weed control practices such as multiple herbicides with different mechanisms of action, and often in combination with various mechanical and cultural practices.

Suspected herbicide resistance can be identified by these factors:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially when control is achieved on adjacent weeds
- A spreading patch of non-controlled plants of a particular species
- Surviving plants mixed with controlled individuals of the same species

To minimize the occurrence of glyphosate-resistant biotypes, including those resistant to glyphosate, implement the following weed management practice options that are practical to your situation. These management practices are applicable to reduce the spread of suspected or confirmed resistant biotypes (managing existing resistant biotypes) and to reduce the potential for selecting for resistance in new species (proactive resistance management).

- Use a diversified approach toward weed management focused on preventing weed seed production and reducing the number of weed seeds in the soil.
- Plant crops into fields that are as weed-free as possible and then keep them as weed-free as possible.
- Plant seed that is as weed-free as possible.
- Scout fields and application sites routinely, before and after herbicide application.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

- Use multiple herbicide mechanisms of action that are effective against the most troublesome weeds on your application site and against those with known resistance.
- Apply herbicides at application rates listed on the label when weeds are within the size range indicated on the label.
- Emphasize cultural practices that suppress weeds by using crop competitiveness.
- Use mechanical and biological weed management practices, where appropriate.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules.
- Manage weed seed at harvest and after harvest to prevent a buildup of the weed seedbank

### 6.2 Management of Glyphosate-Resistant Biotypes

Appropriate testing is needed to determine if a weed is resistant to glyphosate. Call 1-866-99BAYER (1-866-992-2937) or contact your Bayer CropScience representative to report any incidence of non-performance of this product against a particular weed species. To determine if resistance in any particular weed biotype has been confirmed in your area, or for additional information on glyphosate-resistant biotypes, go to [www.weedscience.org](http://www.weedscience.org).

Glyphosate-resistant weeds can be controlled or managed by applying this product in combination with residual preemergence herbicides and/or other postemergence herbicides labeled for control of the targeted weed in the crop being grown or on the site of application. For more information, see the "WEEDS CONTROLLED" section of this label.

Since the occurrence of resistant weeds is difficult to detect before use, to the extent consistent with applicable law, Bayer CropScience accepts no liability for any losses that result from the failure of this product to control resistant weeds.

### 7.0 MIXING

Spray solutions of this product may be mixed, stored and applied using clean stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

A 50-mesh nozzle screen or line strainer on the spray equipment is adequate.

Clean sprayer parts promptly after using this product by thoroughly flushing with water.

#### 7.1 Mixing with Water

PERFORMANCE OF THIS PRODUCT CAN BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS OR DITCHES THAT IS VISIBLY MUDDY OR MURKY.

This product mixes readily with water. Mix spray solutions of this product as follows. Begin filling the mixing tank or spray tank with clean water. Add the required amount of this product near the end of the filling process and mix gently. Foaming of the spray solution can occur during mixing. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, add an appropriate anti-foam or defoaming agent to the spray solution.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### 7.2 Tank Mixtures

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control in the soil, a broader weed control spectrum or an alternate mechanism of action.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers could result in reduced weed control. Bayer CropScience has not tested all tank-mix product formulations for compatibility, antagonism or reduction in product performance. To the extent consistent with applicable law, buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label, or on separate Supplemental Labeling or Fact Sheets published for this product.

When a tank-mix with a generic active ingredient, such as 2,4-D or dicamba, or any other product or material, is listed on this label, it is the responsibility of the pesticide user to ensure that the intended use is included on the label of the product added to the mix.

Bayer CropScience has not tested all tank-mix product formulations for compatibility, antagonism or reduction in product performance. Mixing this product with herbicides or other materials not specified on this label could result in reduced performance of this product. To the extent consistent with applicable law, buyer and all users are responsible for any loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label, or on separate supplemental labeling or Fact Sheets published for this product.

Refer to all individual product labels, supplemental labeling and Fact Sheets for all products in the tank mixture and observe all precautions and limitations on the label, including any application timing and soil restrictions. Users must follow the most restrictive directions for use and precautionary statements for each product in the tank mixture.

This product may be applied at any rate listed on this label in a tank mixture with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed], to provide preemergence and/or improved postemergence control of weeds listed on the label of the tank-mix product. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list*

2,4-D; atrazine; dicamba; bromacil; diuron; imazapyr; metsulfuron methyl; oryzalin; pendimethalin, proflaminate; simazine; sulfosulfuron; trichlopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Banvel; Banvel 480; Barricade 4L; Barricade 65WG; Certainty® Turf; Chopper Gen2; Crossbow; Endurance; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XLT Specialty; Gallery SC; Gallery 75 Dry Flowable Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Goal 2XL; GoalTender; Habitat; Hyvar X; Hyvar X-L; Karmex DF; Krenite S Brush Control Agent; Krovar I DF; Landmark XP; Oust Extra; Oust XP; Outrider; Plateau; Poast; Poast Plus; Ronstar 50 WSP; Ronstar Flo; Ronstar G; Sahara DG; Spike 20P Specialty; Spike 80 DF Specialty; Stalker; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Telar XP; Tordon 101 Mixture Specialty; Tordon 22K Specialty; Tordon K Herbicide Specialty; Transline Specialty; Vanquish; Velpar DF CU; Velpar DF VU; Velpar L CU; Velpar L; Velpar L VU]

---

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

When used in combination as described on this label, and to the extent consistent with applicable law, the liability of Bayer CropScience [*Alternative text:* Bayer Environmental Science] shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Bayer CropScience [*Alternative text:* Bayer Environmental Science] product in such combination use.

### 7.3 Tank-Mixing Procedure

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Add individual tank-mix components to the tank in the following order: wettable powders, flowables, emulsifiable concentrates, drift reduction additives, water soluble liquids (this product) [*Optional text:* , surfactants]. Ensure that the tank-mix products are well mixed in the spray solution before adding this product.

Mix only the quantity of spray solution that will be applied that day. Application of tank-mix solutions that are allowed to stand overnight could result in reduced weed control.

Maintain gentle agitation at all times until the contents of the tank are sprayed out. If the spray mixture is allowed to settle, agitate thoroughly to resuspend the mixture before resuming application.

Keep by-pass line on or near the bottom of the tank to minimize foaming.

A 50-mesh nozzle screen or line strainer on the spray equipment is adequate.

### 7.4 Mixing Spray Solution Concentrations

All reference throughout this label to concentration of this product in a spray solution is on a percentage-of-volume basis.

Prepare the desired volume of spray solution at a given concentration by mixing the amount of this product indicated in the following table in water.

Desired Volume of Spray Solution	Amount of [INSERT BRAND NAME] to Achieve Indicated Concentration in Spray Solution (percent by volume)					
	0.4%	0.7%	1%	1.5%	4%	7%
1 gallon	0.5 fl oz	1 fl oz	1.3 fl oz	2 fl oz	5 fl oz	9 fl oz
25 gallons	13 fl oz	22 fl oz	1 qt	1.5 qts	4 qts	7 qts
100 gallons	1.6 qts	2.8 qts	1 gal	1.5 gals	4 gals	7 gals

2 tablespoons = 1 fluid ounce (fl oz)

For filling backpack and pump-up sprayers, consider mixing the appropriate amount of this product with water in a larger container and then filling the sprayer from the larger container.

### 7.5 Surfactants [*this section is optional in the final printed label*]

Although not always required, nonionic surfactants that are labeled for use with herbicides may be added to spray solutions of this product. Do not reduce application rate or concentration of this product when applying in spray solutions containing additional surfactant.

[*Optional text:* Additional surfactant can increase the performance of this product at water carrier volumes above 30 gallons per acre or at application rates below 16 fluid ounces of product per acre.]

[*Optional text:* Use a surfactant concentration of 0.25 to 0.5 percent (1 to 2 quarts per 100 gallons of spray solution) when adding surfactant that contains at least 70 percent active ingredient, or a 1-percent

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

surfactant concentration (4 quarts per 100 gallons of spray solution) when adding surfactant that contains less than 70 percent active ingredient.]

Read and follow the directions for use and observe all precautionary statements and other information on the surfactant label.

[*Optional label statement:* DO NOT add buffering agents or pH adjusting agents to the spray solution when [INSERT BRAND NAME] is the only pesticide product being applied.]

### 7.6 Ammonium Sulfate [*This section optional in final printed label*]

Unless otherwise directed, the addition of 1 to 2 percent dry ammonium sulfate by weight (8.5 to 17 pounds per 100 gallons of water), could increase the performance of this product on annual and perennial weeds, particularly under hard water conditions, drought conditions or when tank-mixed with certain residual herbicides. An equivalent amount of a liquid formulation of ammonium sulfate may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water promptly after use to reduce corrosion.

When using ammonium sulfate, apply this product at rates or spray solution concentrations as directed on this label; lower the application rates or concentration could result in reduced weed control.

### 7.7 Colorants and Dyes

Colorants and marking dyes may be added to spray solutions of this product; however, they can reduce the performance of this product. Use colorants and dyes according to the manufacturer's directions.

### 7.8 Drift Reduction Additives

Drift reduction additives may be used with all application equipment types, except wiper applicators, sponge bars and controlled droplet applicators (CDA). However, use of drift reduction additives can affect spray coverage, which could reduce the performance of this product. When a drift reduction additive is used, read and follow all directions for use, precautions, limitations and other information on the product label.

## 8.0 APPLICATION EQUIPMENT AND TECHNIQUES

This product may be applied with the following equipment:

**Aerial Application Equipment**—fixed-wing and helicopter [*Alternative text:* helicopter only] [*Optional text:* (Aerial application allowed by helicopter only in) [*list states where aerial application is allowed by helicopter only, if applicable*)]

**Ground Application Equipment**—boom or boomless systems, pull-type sprayers, floaters, pick-up sprayers, spray coupes and other ground broadcast application equipment

**Handheld Sprayers**—backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other handheld and motorized spray equipment used to direct the spray onto unwanted foliage

\*This product is not registered in California or Arizona for use in mistblowers.

**Selective Application Equipment**—recirculating sprayer, shielded and hooded sprayers, wiper applicator, sponge bar, single or hollow stem injectors, tree injector, spray bottle

**Injection Systems**—aerial or ground injection sprayers

**Controlled Droplet Applicator (CDA)**—handheld or boom-mounted applicators that produce a spray pattern consisting of a narrow range of droplet sizes

APPLY THIS PRODUCT USING PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF ACCURATELY DELIVERING DESIRED VOLUMES.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Do not apply this product through any type of irrigation system.

### **8.1 Spray Drift Management**

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, [*Optional text:* GREEN] STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, [*Optional text, if applicable:* EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY CROPS], AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation, as even small quantities of this product can cause severe damage or destruction to the crop, plants or other vegetation on which application was not intended.

AVOID DRIFT. USE EXTREME CARE TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHEN APPLYING THIS PRODUCT.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and grower are responsible for considering all these factors when making decisions regarding the application of this product.

The likelihood of injury occurring as the result of spray drift while applying this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or generation of fine particles (mist) that are likely to drift.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFERS MUST BE MAINTAINED.

AVOID APPLYING THIS PRODUCT AT EXCESSIVE SPEED OR SPRAYER PRESSURE.

### **8.2 Aerial Application Equipment**

Unless otherwise prohibited, all broadcast applications of this product described on this label may be made using aerial application equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this label and on separate supplemental labeling published for this product.

DO NOT APPLY THIS PRODUCT USING AERIAL APPLICATION EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED ON THIS LABEL OR ON SEPARATELY PUBLISHED SUPPLEMENTAL LABELING FOR THIS PRODUCT.

[*Optional voluntary label restriction:* Aerial application of this product may be made using helicopters only.]

[*Optional voluntary label restriction:* Aerial application of this product may be made using helicopters only in] [*List states where aerial application is permitted by helicopter only, if applicable*].

FOR SPECIFIC USE INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS RELATED TO THE AERIAL APPLICATION OF THIS PRODUCT IN CALIFORNIA OR SPECIFIC COUNTIES THEREIN, REFER TO THE LIMITATIONS ON AERIAL APPLICATION IN THAT STATE OR COUNTY PRESENTED IN THIS SECTION.

Apply this product at the rate specified on this label in 3 to 25 gallons of water per acre, unless otherwise directed. Use a larger spray volume within this range where weeds, brush, trees and vines are dense or form multiple canopy layers.

Avoid direct application to any body of water.

Drift control additives may be used.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

### **Aircraft Maintenance**

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES COULD RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 can help prevent corrosion.

### **AERIAL SPRAY DRIFT MANAGEMENT**

The following drift management requirements must be followed to minimize off-target drift movement during aerial application. [*Optional label text, if appropriate:* These requirements do not apply to forestry applications.]

1. The distance of the outermost nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the [*Optional text, if application using fixed-wing aircraft allowed:* wingspan or] rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be followed.

### **Importance of Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly or under unfavorable environmental conditions, such as in windy, high temperature with low humidity, and/or inversion conditions as described below.

### **Controlling Droplet Size**

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Operate sprayer at a pressure towards the lower end of the range listed for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provides uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the [*Optional text, if application using fixed-wing aircraft allowed:* wingspan or] rotor length can further reduce drift without reducing swath width.
- **Application height:** Application must be made at a height of 10 feet or less above the top of the largest plants, unless a greater height is required for aircraft safety. Making the application at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### Swath Adjustment

When an application is made in the presence of a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase the swath adjustment distance with increasing drift potential (higher wind, smaller droplets, etc.).

### Wind

Drift potential is lowest at wind speeds of between 2 and 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Avoid application when wind speeds are below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making an application in low relative humidity, set application equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Do not apply this product during a temperature inversion as drift potential is high under these conditions. Temperature inversions restrict vertical air mixing, which causes small droplets to remain suspended in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from a sensitive area).

**State Specific Limitations on Aerial Application** [*This section is optional, to be used only if it is NOT stated on the label that this product is not registered for use in California, and only if required by the State*]

---

### LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY

---

[*Optional voluntary restriction:* Aerial application of this product in California may be made by helicopter only.]

DO NOT apply this product using aerial application equipment in residential areas.

AVOID DRIFT – DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT OF THIS PRODUCT ONTO ANY VEGETATION TO WHICH APPLICATION WAS NOT INTENDED CAN CAUSE DAMAGE. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, USE PROPER AERIAL APPLICATION EQUIPMENT FITTED WITH APPROPRIATE NOZZLES AND MAINTAIN ADEQUATE BUFFERS.

Follow the directions below when making an aerial application near non-target crops, desirable annual vegetation, or desirable perennial vegetation after bud break and before total leaf drop.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

1. Do not apply this product within 100 feet of all desirable vegetation or non-target crops.
2. If winds are blowing up to 5 miles per hour TOWARD desirable vegetation or non-target crops, do not apply this product within 500 feet of the desirable vegetation or crops.
3. If winds are blowing between 5 and 10 miles per hour TOWARD desirable vegetation or non-target crops, a buffer zone greater than 500 feet might be needed to protect the desirable vegetation or crops.
4. Do not apply this product using aerial application equipment when winds are blowing in excess of 10 miles per hour
5. Do not apply this product using aerial application equipment when inversion conditions exist.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be applied in California using aerial application equipment. Tank mixtures of this product with 2,4-D amine formulations may be applied by air in California on fallow fields and in reduced tillage systems [*Optional text, if applicable:* , and for alfalfa and pasture renovation applications] only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

---

### **ADDITIONAL LIMITATIONS ON AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY**

---

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

- North: Fresno County line
- South: Fresno County line
- East: State Highway 99
- West: Fresno County line

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

#### **Written Directions**

Written directions MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. These written directions MUST state the proximity of surrounding crops and that conditions of each manufacturer's product label and this label have been satisfied.

#### **Aerial Applicator Training and Equipment**

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

**Applications at Night** – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

For additional information on the proper aerial application of this product in Fresno County, call 1-866-99BAYER (1-866-992-2937).

---

### 8.3 Ground Application Equipment

Apply this product at an appropriate rate specified on this label in 3 to 40 gallons [*Alternative volume*: 10 to 60 gallons] of water per acre when making a broadcast application using ground application equipment, unless otherwise directed on this label or on separate supplemental labeling or Fact Sheets published for this product. As the weed density increases, increase the spray volume towards the upper end of this range to ensure complete coverage. Use nozzles that will avoid generating a fine mist. For best performance of this product with ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

### 8.4 Handheld Sprayers

When using a handheld sprayer, apply spray solutions of this product uniformly and completely to the foliage of target weeds using a coarse droplet spectrum and a spray-to-wet technique; do not spray to the point of runoff. For the appropriate concentration of this product in the spray solution and timing of application to control specific weeds, woody brush, trees and vines, refer to the “WEEDS CONTROLLED” section of this label

For control of annual weeds, make application when weeds are small and prior to seedhead or bud formation. For control of perennial weeds, woody brush, trees and vines, make application after flowering and before fall color and leaf drop.

When making a low-volume directed spray application to annual and perennial weeds, woody brush, trees and vines using a handheld sprayer, ensure that at least 50 to 75 percent of the foliage or the top one-half of each unwanted plant is sprayed. If a straight stream nozzle is used, start the application at the top of the targeted plant and spray from top to bottom in a lateral zig-zag motion. To ensure uniform and complete coverage, spray both sides of large or tall woody brush, trees and vines, or when foliage is thick and dense, or where there are multiple sprouts. For best performance of this product on woody brush, trees and vines, apply to actively growing vegetation after full leaf expansion and flowering, prior to fall color and leaf drop.

The following table summarizes various methods of foliar application using a backpack sprayer with a spray-to-wet or low-volume directed spray technique and high-volume sprayer application using handheld application equipment for control or partial control of herbaceous weeds, woody brush, trees and vines listed in the “WEEDS CONTROLLED” section of this label.

Method of Application	Spray Solution Concentration	Spray Volume
Handgun or Backpack Sprayer	1.5% by volume	Spray-to-wet technique
Low-Volume Directed Spray (Backpack)	4 to 7% by volume	15 to 25 gallons/acre
Modified High-Volume Spray	1.5 to 3% by volume	40 to 60 gallons/acre

Low-volume directed spray application with a backpack sprayer works best when applying to weeds and brush less than 10 feet tall. For taller weeds and brush, a high-volume handgun can be modified by reducing the nozzle size and spray pressure to produce a modified high-volume directed spray application.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### 8.5 Selective Application Equipment

Selective application equipment allows this product to be applied to weeds growing near a crop or other desirable vegetation without killing the desirable vegetation. Selective application equipment must be capable of preventing all contact of the herbicide solution with the desirable vegetation and operated without spray mist escape, leakage or dripping of the herbicide solution.

**AVOID CONTACT OF THIS HERBICIDE WITH DESIRABLE VEGETATION.** Contact of this product with desirable vegetation could result in unwanted plant damage or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

This product may be diluted with water and applied using a recirculating sprayer, shielded sprayer, hooded sprayer, wiper applicator or sponge bar to weeds listed on this label growing on any terrestrial non-food or non-feed crop site listed on this label, where feasible. This product may also be used with sprayers equipped with optical weed sensor technology. Other selective equipment that may be used to deliver or apply this product are single and hollow stem injectors, tree injectors, wiper applicators for cut stem and cut stump applications, and spray or squirt bottles for cut stem, cut stump and frill applications to control large stem weeds, brush, trees and vines listed on this label.

#### **Recirculating Sprayer**

A recirculating sprayer directs the spray solution onto weeds growing above desirable vegetation, while spray solution that is not intercepted by weeds is collected and returned to the spray tank for reapplication. A recirculating sprayer may be used to apply spray solutions of this product to weeds listed on this label in any terrestrial non-crop site described on this label.

#### **Shielded and Hooded Sprayers**

A shielded sprayer directs the herbicide solution onto targeted weeds while using an impervious material, or shield, to protect desirable vegetation from coming into contact with the herbicide spray. To provide maximum protection for desirable vegetation, keep shields properly adjusted and use nozzles that provide uniform coverage within the application area.

A hooded sprayer is a type of shielded sprayer where the spray pattern is fully enclosed, including the top, sides, front and back, thereby shielding desirable vegetation from the spray solution.

This product may be diluted with water and applied using a shielded or hooded sprayer to weeds listed on this label growing on any terrestrial non-crop site described on this label, where feasible.

Properly adjust the hood to protect desirable vegetation. Ensure that the hood is capable of completely enclosing the spray pattern.

A hooded sprayer must be configured and operated in a manner that minimizes bouncing and avoids raising the hood up off the ground surface at any time. If the hood is raised, spray particles can escape and come into contact with desirable vegetation, causing damage to or destruction of the desirable vegetation. Avoid operating this equipment on rough or sloping terrain where the spray hood is likely to rise up off the ground surface.

Use hoods designed to minimize excessive dripping or runoff down the inside of the hood, such as a single, low pressure, low drift, flat-fan nozzle with an 80- to 95-degree spray angle positioned at the top center of the hood, with a spray volume of 20 to 30 gallons per acre.

The following procedures will help reduce the potential for injury to desirable vegetation when using a hooded sprayer:

- Operate the sprayer with the hood on the ground or skimming across the ground surface.
- Leave at least an 8-inch untreated strip over the drill row. (For example, if a crop row width is 38 inches, use a sprayer hood with a maximum width of 30 inches.)

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

- Operate at a ground speed no greater than 5 miles per hour to minimize bouncing of the hooded sprayer.
- Apply when wind speed is 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the application area.

Injury to a crop or other desirable vegetation can occur when application is made to foliage of weeds that come into direct contact with the desirable vegetation. Do not apply this product when leaves of desirable vegetation are growing in direct contact with weeds. Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting or destruction.

### **Wiper Applicator**

A wiper applicator is a device that physically wipes this product or solutions of this product directly onto the unwanted vegetation or cut stump. Any handheld device that is capable of physically wiping this product or solutions of this product directly onto the target weed or cut stump, such as a paint brush, may be used.

A mechanical wiper applicator, such as a rope wick or sponge bar that can be driven through a field over the top of desirable vegetation to control tall weeds growing above the desirable vegetation, must be designed, maintained and operated to prevent the herbicide solution from coming into contact with desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling onto desirable vegetation can result in discoloration, stunting, or destruction. Avoid leakage or dripping onto desirable vegetation. To protect the desirable vegetation, adjust the height of the applicator to ensure that the wiper contact point is a minimum of 2 inches above the desired vegetation.

Weeds that do not come into contact with the herbicide solution will not be affected. The more weed foliage exposed above the desirable vegetation the better mechanical wiper applicators work. Better results can be obtained when weeds are a minimum of 6 inches above the desired vegetation. Poor contact can also occur when weeds are growing in dense clumps, when operating in areas of severe weed infestation or when weed height varies dramatically. In these situations, more than one application of this product might be necessary to achieve desired weed control.

For optimal results, operate wiper applicators at a ground speed of no greater than 5 miles per hour. Performance in areas of heavy weed infestation can be improved by reducing speed, which will provide more time for re-saturation of the wiper with the herbicide solution and more contact time of the wiper with the weed. Better weed control using a wiper applicator can also be obtained when two applications are made traveling in opposite directions across the field.

Keep wiping surfaces clean.

Be aware that on sloping ground the herbicide solution can migrate to one side, causing dripping at the lower end and drying of the wiper surface at the upper end of the applicator.

Do not apply this product with a wiper applicator when weeds are wet.

Do not add surfactant to the herbicide solution when using a wiper applicator.

With **Rope Wick or Sponge Wick Applicators**, apply solutions ranging from 25 to 70 percent of this product by volume in water.

With **Panel Applicators**, apply solutions ranging from 25 to 100 percent (undiluted) of this product by volume in water.

Mix only the amount of this product that will be used during a 1-day period, as reduced product performance can result from the use of solutions held in storage.

Clean wiper parts promptly after using this product by thoroughly flushing with water.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### Single and Hollow Stem Injectors

Control of certain weeds listed in the “WEEDS CONTROLLED” section can be obtained by injecting this concentrated product or solutions of this product directly in or onto the target weed. Ensure that the handheld injector being used for this application is capable of accurately delivering the volume specified on the label. When making stem injections, the combined total use of this product must not exceed 7 quarts per acre per year. At 5 milliliters of concentrated (undiluted) product per stem, 7 quarts will treat approximately 1300 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

#### 8.6 Injection Systems

This product may be used in aerial and ground injection spray systems as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this concentrated product with the undiluted concentrate of other products when using injection systems, unless otherwise directed.

#### 8.7 Controlled Droplet Applicator (CDA)

The amount of this product applied per acre using a controlled droplet application (CDA) must be no less than the rate specified on this label for application using conventional broadcast application equipment.

A controlled droplet applicator produces a spray pattern that is not easily visible. Use extreme care to avoid spray or drift from coming into contact with foliage or any other green tissue of desirable vegetation, as plant damage or destruction could result.

### 9.0 TERRESTRIAL USE SITES

This product may be used according to the directions described on this label to control annual and perennial weeds, woody brush, trees and vines listed on this label on any terrestrial site described on this label.

This product may be used to control weeds, woody brush, trees and vines on maintained landscapes, on improved and unimproved land, on lawns and turf and around ornamentals on industrial, commercial and residential sites, including [*Select all uses that are applicable to the final printed labeling*: airports, apartment complexes, chaparrals, ditch banks, driveways, dry ditches, dry canals, farmsteads, fencerows, forestry sites, golf courses, greenhouses, lumber yards, manufacturing sites, municipal sites, natural areas, nurseries, office complexes, ornamental beds, parks, parking areas, pastures, petroleum tank farms, pumping installations, railroads, rangeland, recreational areas, residential areas, roadsides, schools, shadehouses, sod and turfgrass seed farms, sports complexes, storage areas, substations, utility rights-of-way, utility sites, warehouse areas, wildlife food plots and wildlife management areas.]

This product may be used for non-selective control of unwanted vegetation on any site listed on this label for trim-and-edge application around objects, including around building foundations, equipment storage areas and trees, along and in fences, and to eliminate unwanted vegetation growing in and around established shrub beds and ornamental plantings. This product may also be used for complete elimination of vegetation from a terrestrial site prior to planting ornamentals, flowers, or turfgrass (sod or seed), and prior to land development, including prior to beginning construction projects or the laying of asphalt or other road material. Application of this product may be repeated, as needed, to maintain bare ground, up to a total application of 7 quarts per acre per year.

This product may be used for establishment and maintenance of fuel breaks, for establishing fire perimeters and black lines, along fire roads and to facilitate prescribed burning practices on any site described on this label.

[*Optional label text*: This product may also be used for weed control or growth regulation on] [*Optional list of any terrestrial uses that are included on this Master Label, including, but not limited to*: Christmas tree farms, farmsteads, production nurseries, sod farms and turfgrass seed farms.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Unless otherwise directed, application of this product may be made according to the directions for use in the sections that follow to any of these sites using any method of application described on this label to control any weeds, woody brush, trees and vines listed in the “WEEDS CONTROLLED” section of this label.

**RESTRICTIONS:** Before using this product, read and follow the Maximum Application Rate information in the “PRODUCT INFORMATION” section of this label. Observe these and all other maximum application rates stated throughout this label. Maximum application rates apply to the use of this product combined with the use of any and all other herbicides containing glyphosate, whether applied separately or as mixtures. Calculate the application rates (glyphosate acid equivalents) and ensure that the total use of this and other glyphosate-containing products does not exceed the stated maximum rate.

### 10.0 ADDITIONAL SITE MANAGEMENT INFORMATION

The following sections contain additional use information specifically related to certain use sites. Unless otherwise directed, any application of this product described in the “WEEDS CONTROLLED” section or any other section of this label may be made on the use sites described in the sections that follow, where applicable, using any method of application described on this label that is appropriate.

#### 10.1 Commercial, Residential and Recreational Area Management

All applications of this product described on this label may be used in commercial, residential and recreational areas, including parks, schools and athletic fields, using any method of application described on this label, including spot treatment of unwanted vegetation, trim-and-edge application around trees, fences, walking paths, buildings, sidewalks, nature trails and other objects in these areas, to eliminate unwanted weeds growing in established shrub and ornamental beds, for turf management and renovation, and to eliminate vegetation from a site prior to development, including prior to planting an area to ornamentals, flowers or turfgrass (sod or seed), or beginning construction projects.

#### 10.2 Forestry, Hardwood and Christmas Tree Management

This product may be used for control or partial control of woody brush, trees, and herbaceous weeds on any tree site, including forestry settings, Christmas tree plantations, and silvicultural and production nursery sites, using any method of application described on this label. See the “WEEDS CONTROLLED” section of this label for application rates and specific use directions.

##### **Weed Management, Site Preparation**

This product may be used to control or partially control undesirable woody brush, trees, vines and herbaceous weeds listed on this label for preparing sites prior to planting any tree species, including Christmas trees, eucalyptus trees and hybrid tree cultivars, and for controlling weeds around established trees, [*Optional text:* for the release of conifer and hardwood trees,] establishing wildlife openings and maintaining roads on any tree site.

**TANK MIXTURES:** This product may be applied in a tank-mix with products containing one or more of the active ingredients listed below, [*Optional text:*, or one or more of the products listed,] to increase the spectrum of vegetation controlled. Any application rate of this product listed on this label may be used in a tank-mix with the following products for tree site management, including site preparation, provided that the product is labeled for the use on the site of application and prior to planting the desired species. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

imazapyr; metsulfuron methyl; sulfometuron methyl; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Chopper; Chopper Gen2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Landmark XP; Oust Extra; Oust XP

---

For control or partial control of dense stands of vegetation or of hard-to-control woody brush, trees and vines, apply these products at a rate or spray solution concentration towards the higher end of the given range.

### **Conifer Release, Mid-Rotation Conifer Release, Hardwood Release, Timber Stand Improvement**

This product may be applied as a directed spray using a handheld sprayer or using any selective application equipment described on this label to control woody and herbaceous weeds and other undesirable understory vegetation below the tree crop canopy in conifer plantations, hardwood sites, Christmas tree plantations and silvicultural and ornamental nurseries to facilitate the release and growth of conifer and hardwood trees.

This product may also be applied using ground broadcast equipment or as a directed spray application for mid-rotation release under the canopy of pines, other conifers and hardwoods.

**PRECAUTIONS:** Avoid contact of spray drift, mist or drips with foliage, green bark or non-woody surface roots of desirable plant species. Use application techniques that prevent or minimize contact of this product with foliage of desired trees or other plants through direct contact or off-target spray movement.

*[Optional label text:* RESTRICTIONS: Do not apply this product as an over-the-top broadcast application for conifer or hardwood release, unless otherwise directed on this label or on separate supplemental labeling for this product.]

### **Conifer Release – Broadcast Application** *[This section is optional in the final printed label]*

This product may be broadly applied over the top of conifer tree species listed in this section after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring for control, partial control or suppression of herbaceous weeds and hardwoods listed in the “WEEDS CONTROLLED” section of this label to facilitate the release of these desirable tree species in a forestry, plantation or nursery setting. Unless otherwise directed, make this application only where conifers have been established for a minimum of one growing season.

**PRECAUTIONS:** Conifer injury can occur if this product is applied over the top of conifers at rates higher than prescribed on this label, where spray applications overlap, if application is made when conifers are actively growing, or when they are growing under stress from drought, flood, improper planting or insect, animal or disease damage.

### Conifer Release Outside the Southeastern United States

For release of the following conifer species growing for a minimum of one growing season in most areas outside the southeastern United States, apply 22 to 44 fluid ounces of this product per acre as a broadcast application over the top of the conifer trees.

- Douglas fir
- Hemlock
- California redwood
- Fir species
- Pines\*
- Spruce

\* Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Apply 22 to 36 fluid ounces of this product for release of Douglas fir, pine and spruce that have been established for only one growing season (except in California).

For release of spruce (*Picea* spp.) in Maine, Michigan, Minnesota, New Hampshire and Wisconsin, up to 2 quarts of this product may be applied after formation of final resting buds in the fall for control of woody brush and tree species.

**PRECAUTIONS:** Ensure that the conifers are well hardened off before application of this product. [*Optional text, if adding surfactant to spray solutions of this product is allowed:* The addition of nonionic surfactants to spray solutions of this product when making an over-the-top conifer release application could cause conifer injury.]

### Conifer Release in the Southeastern United States

For release of the following conifer species established for more than one growing season in the southeastern United States, apply 32 to 54 fluid ounces of this product per acre in the fall as a broadcast application over the top of the trees. For release of these species after only one growing season, apply only 22 fluid ounces of this product per acre.

- Eastern white pine
- Longleaf pine
- Slash pine
- Loblolly pine
- Shortleaf pine
- Virginia pine

**TANK MIXTURES:** This product may be applied for conifer release in a tank-mix with products containing one or more of the active ingredients listed below [*Optional text;* or one or more of the products listed,] to provide a broader spectrum of postemergence weed control and for residual control of weeds listed on the label of those products. Only apply products that are labeled for application over the top of conifer species when tank-mixing with this product. Refer to the individual product labels for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

atrazine; imazapyr; metsulfuron methyl; sulfometuron methyl

Arsenal; Oust Extra; Oust XP

---

For release of Douglas fir established for a minimum of one growing season prior to bud swell in early-spring, apply 22 fluid ounces of this product per acre in a tank-mix with an appropriate rate of atrazine.

For herbaceous release of loblolly pine, Virginia pine and longleaf pine in the spring and early-summer, apply 11 to 16 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra (EPA Reg. No. 432-1557; *sulfometuron methyl, metsulfuron methyl*) or Oust XP (EPA Reg. No. 432-1552; *sulfometuron methyl*).

### Late-Summer and Fall after Resting Bud Formation

For release of jack pine, white spruce and white pine, apply 22 to 44 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP that will not harm these conifer species.

For release of Douglas fir, apply 22 to 32 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal.

For release of balsam fir and red spruce, apply 44 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal (EPA Reg. No. 241-346; *imazapyr*) or Arsenal Herbicide Applicators Concentrate (EPA Reg. No. 241-299; *imazapyr*).

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### 10.3 Native and Wildlife Habitat Management

This product may be used to control exotic and other undesirable vegetation in wildlife habitat and natural areas, including riparian and estuarine areas, rangeland and wildlife refuges. Application may be made to allow recovery of native plant species or prior to planting desirable native species, and for similar broad-spectrum vegetation control. Spot, cut stump, cut stem, stem injection, wiper and all other methods of application listed on this label may be used to selectively remove unwanted plants for habitat management and enhancement.

This product may also be used to eliminate annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product or native species may be allowed to repopulate the area naturally. If tillage is needed to prepare a seedbed, wait a minimum of 7 days after application before tillage to allow translocation of this herbicide into underground plant parts.

### 10.4 Ornamental and Production Nursery Management

All uses of this product described on this label may be used in a plant nursery setting using any method of application described.

This product may be used to clear an area of unwanted vegetation prior to planting any ornamental plant, tree, shrub or other plants.

This product may also be used to control weeds growing around established woody ornamental species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew, and to trim and edge around potted plants and other objects in a plant nursery.

**PRECAUTIONS:** Protect desirable plants from the spray solution using shields or coverings made of waterproof material. Take care to avoid contact of spray, drift or mist with foliage, green stems or immature bark of established ornamental species.

#### **Greenhouse/Shadehouse**

This product may be used to control weeds growing in and around greenhouses and shadehouses.

**RESTRICTIONS:** Desirable vegetation must not be present during application in a greenhouse. Turn air circulation fans off before applying this product inside a greenhouse or shadehouse and leave them off until the application solution has dried.

### 10.5 Pasture Management

The use of this product in pastures includes use on bahiagrass, bermudagrass, bluegrass, brome, fescue, guinea grass, kikuyu grass, orchardgrass, pangola grass, ryegrass, Timothy, and wheatgrass.

#### **Preplant, Preemergence, Pasture Renovation**

This product may be applied prior to planting or emergence of forage or perennial grasses. Refer to the "WEEDS CONTROLLED" section of this label for application rates of this product for control of specific weeds.

**RESTRICTIONS:** If the total application rate of this product is 2 quarts per acre or less, no waiting period between application and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait a minimum of 8 weeks after application before grazing or harvesting.

#### **Spot Application, Wiper Application**

This product may be applied in pastures as a spot treatment or over the top of desirable grasses using a wiper applicator to control taller growing weeds. For better weed control, remove domestic livestock before application to allow for sufficient plant growth and wait a minimum of 7 days after application before grazing

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

livestock or harvesting for feed. See additional instructions on the use of wiper applicators in the “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label.

**RESTRICTIONS:** For spot treatment or use with a wiper applicator at rates of 2 quarts per acre or less, this product may be applied over the entire pasture or any portion of it. At rates greater than 2 quarts per acre, this product may be applied over no more than 10 percent of the total pasture at any one time. Application may be repeated in the same area at 30-day intervals.

### **Weed Suppression in Dormant Pastures**

This product may be applied in dormant pastures to suppress competitive growth and seed production of annual weeds and other undesirable vegetation. Apply 8 to 11 fluid ounces of this product per acre using broadcast application equipment on pastures in late-fall after desirable perennial grasses have reached dormancy or in late-winter before desirable perennial grasses break dormancy and initiate green growth.

**PRECAUTIONS:** Higher application rates may be used for hard-to-control weeds; however, higher rates can cause stand reduction. Some stunting of perennial grasses can occur if broadcast application is made when they are not dormant.

**RESTRICTIONS:** No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 2 quarts of this product per acre per year onto pasture grasses except for renovation. If reseeding is needed due to severe stand reduction, no waiting period is required after application of this product before seeding the pasture grasses listed at the beginning of this section; for all other pasture grasses, wait a minimum of 30 days after application before seeding.

### **10.6 Railroad Management**

All uses of this product described in the “WEEDS CONTROLLED” or any other section of this label may be used on railroad sites using any method of application described.

Application of this product along railroad rights-of-way may be made in up to 80 gallons of spray solution per acre.

### **Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment**

This product may be used to maintain bare ground on railroad ballast and shoulders and reduce the need for mowing and mechanical brush removal along railroad rights-of-way. Application of this product may be repeated as weeds continue to emerge in order to maintain bare ground, up to a maximum total application rate of 7 quarts per acre per year.

**TANK MIXTURES:** This product may be applied in a tank mixture with products containing one or more of the active ingredients listed below [*Optional text*: or one or more of the products listed], to broaden the spectrum of control of woody brush, trees and vines for bare ground, ballast and shoulder, crossing and spot applications, and other brush, tree and vine control on railroad sites, provided that the product used is labeled for the application being made. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for each product in the mix. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below*: For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diquat; diuron; hexazinone; imazapyr; metsulfuron methyl; pelargonic acid; simazine; sulfometuron methyl; sulfosulfuron; tebuthiuron; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Chopper; Chopper Gen2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Habitat; Hyvar X; Hyvar X-L; Krovar I DF; Oust Extra; Oust XP; Outrider®; Princep 4L; Princep Caliber 90; Princep Liquid; Sahara DG; Scythe; Stalker; Spike 20P Specialty; Spike 80DF Specialty; Telar XP; Transline Specialty; Vastlan Specialty; Velpar DF CU; Velpar DF VU; Velpar L CU; Velpar L VU]

---

### Brush, Tree and Vine Control

This product may be used to control woody brush, trees and vines along railroad rights-of-way. Apply 2.5 to 7 quarts of this product in up to 80 gallons of spray solution per acre as a broadcast application using either a boom or boomless sprayer. Apply a 0.7- to 1.5-percent solution of this product when using high-volume application equipment with a spray-to-wet technique, or a 4- to 7-percent solution when using a low-volume directed spray for spot application.

**TANK MIXTURES:** This product may be applied in a tank-mix with products containing one or more of the active ingredients listed below, [*Optional text*], or one or more of the products listed,] to broaden the spectrum of control of woody brush, trees and vines along railroad rights-of-way, provided that the product is labeled for use on these sites. Refer to the individual label of all products used in the tank mixture for approved use sites and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the “TANK-MIX PRODUCT INFORMATION” section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

chlorsulfuron; clopyralid; dicamba; fosamine; hexazinone; imazapyr; metsulfuron methyl; picloram; triclopyr

Arsenal; Arsenal Herbicide Applicator's Concentrate; Chopper; Chopper Gen2; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Habitat; Krenite S Brush Control Agent; Stalker; Telar XP; Tordon 101 Mixture Specialty; Tordon 22K Specialty; Tordon K Herbicide Specialty; Transline Specialty; Vanquish; Vastlan Specialty; Velpar DF CU; Velpar DF VU; Velpar L; Velpar L CU; Velpar L VU]

---

### Weed Control in Dormant and Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds in dormant and actively growing bermudagrass along railroad rights-of-way. See the “WEEDS CONTROLLED” section of this label for directions for use of this product for weed control in grasses.

#### 10.7 Rangeland Management

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangeland. Slight discoloration of the desirable grasses could occur, but will re-green and resume growing under moist soil conditions as effects of this product wear off.

Preventing seed production is critical to the control of invasive annual grassy weeds on rangeland. Yearly application of this product to eliminate invasive annual weeds before they produce seed will help eliminate

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

viable weed seeds from the soil. Delay grazing of the area after application to allow desirable perennials to grow, flower and re-seed the area.

**Bromus Control:** A broadcast application of 5 to 11 fluid ounces of this product per acre will control or suppress downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*), cheatgrass (*Bromus secalinus*), cereal rye and jointed goatgrass on rangeland. For maximum performance, apply this product when most brome plants are in early-flower and before the plants, including seedheads, turn color. Allow for secondary weed flushes to occur after spring rains to further deplete the seed reserve in the soil and encourage perennial grass conversion on weedy sites. Apply this product in the fall in areas where spring moisture is normally limited and fall germination allows for good weed growth and weed seed depletion.

**Medusahead Control:** To control or suppress medusahead, apply 11 fluid ounces of this product per acre at the 3-leaf stage. Delaying application beyond this stage will result in reduced or unacceptable control. Controlled burning prior to application of this product will eliminate the thatch layer produced by slowly decaying culms. Allow new weed growth to occur before applying this product after a burn. Repeat this application annually to eliminate medusahead seeds in the soil and to allow desirable perennial grasses to repopulate the area.

**RESTRICTIONS:** Do not apply more than 2 quarts of this product per acre per year on rangeland. Do not use ammonium sulfate when applying this product on rangeland grasses. No waiting period between application of this product and feeding or livestock grazing is required.

### 10.8 Roadside Management

All uses of this product described on this label may be used for weed management along roadways, including weed control in dormant and active bermudagrass and bahiagrass, weed control along shoulders and under and around guardrails, signposts and other objects along the road, using any method of application described on this label.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed], for shoulder, guardrail, spot application and for maintaining bare ground, provided that the product used is labeled for use on these sites. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diuron; fosamine; hexazinone; imazapic; imazapyr; metsulfuron methyl; oryzalin; oxadiazon; pendimethalin; picloram; proflam; prodiamine; simazine; sulfometuron; sulfosulfuron; triclopyr

AAtrex 4L; AAtrex Nine-O; Banvel; Barricade 65WG; Chopper; Chopper Gen2; Crossbow; Direx 4L; Escort XP; Endurance; Formula 40; Gallery 75 Dry Flowable Specialty; Gallery SC; Garlon 4 Specialty; Garlon XRT; Hyvar X; Karmex DF; Krenite S Brush Control Agent; Krovar I DF; Landmark XP; Oust Extra; Oust XP; Outrider; Pendulum 3.3 EC; Pendulum AquaCap; Plateau; Poast; Poast Plus; Princep 4L; Ronstar 50 WSP; Ronstar Flo; Ronstar G; Sahara DG; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Telar XP; Tordon K Herbicide Specialty; Vanquish; Vastlan Specialty; Velpar DF CU; Velpar DF VU; Velpar L CU; Velpar L VU; Weedar 64]

---

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### 10.9 Utility Management

This product may be used along electrical power, pipeline and telephone rights-of-way, and on all sites associated with these utility rights-of-way, including substations, access roads and rail lines, and along similar rights-of-way that run in conjunction with utilities, for spot application on unwanted vegetation, side-trimming, trim-and-edge application around objects, weed control prior to planting a utility site to ornamentals, flowers, or turfgrass (sod or seed), turf management, to eliminate unwanted weeds growing in established shrub or ornamental beds, to prepare or establish wildlife openings and to eliminate vegetation prior to or beginning construction projects. Application of this product may be repeated as needed to maintain bare ground as weeds continue to emerge, up to a maximum application rate of 7 quarts per acre per year.

**TANK MIXTURES:** This product may be tank-mixed with products containing one or more of the active ingredients listed below [*Optional text:*, or one or more of the products listed], for use on utility sites, provided that the product used is labeled for use on these sites. Refer to the individual label of all products used in the tank mixture for approved sites and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to the mix. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

2,4-D; atrazine; bromacil; chlorsulfuron; clopyralid; dicamba; diuron; fosamine; hexazinone; imazapic; imazapyr; metsulfuron methyl; oryzalin; pendimethalin; prodiamine; simazine; sulfometuron methyl; sulfosulfuron; triclopyr

AAtrex 4L; AAtrex Nine-O; Arsenal Herbicide Applicators Concentrate; Endurance; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XLT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Hyvar XL; Krenite S Brush Control Agent; Krovar I DF; Oust Extra; Oust XP; Outrider®; Plateau; Sahara DG; Surflan AS Agricultural; Surflan AS Specialty; Surflan Flex; Surflan Flex T&O; Surflan XL 2G; Telar XP; Transline Specialty; Vanquish; Vastlan Specialty; Velpar DF CU; Velpar DF VU; Velpar L CU; Velpar L VU; Weedar 64]

---

To broaden the spectrum of weed control when side trimming, apply this product in a tank-mix with a product containing triclopyr.

Ensure that any product containing triclopyr is thoroughly mixed with water in the spray tank according to label directions before adding this product to the spray mixture. Maintain continuous agitation when adding this product in order to avoid tank-mix incompatibility problems.

### 11.0 CROP USES [*This section is optional in the final label*]

*[Any crop use listed in the Directions for Use with Food Crops section of this Master Label may be included on this sub-label for terrestrial use]*

### 12.0 WEEDS CONTROLLED

Read the entire label before proceeding to use this product.

Always use a higher application rate or spray solution concentration of this product within a given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Poor weed control could be realized if application is made to weeds covered with dust. For weeds that have been mowed, grazed or cut, allow re-growth to occur prior to application of this product.

Refer to the sections that follow for application rates and timing of application for the control of annual and perennial weeds, woody brush, trees and vines.

### 12.1 Weed Control, Renovation and Chemical Mowing in Turf

The use of this product described in this section may be applied to turfgrass growing on any terrestrial site listed on this label. Ensure that any tank-mix product applied with this product is labeled for the intended use and on the site of application.

#### Weed Control in Dormant Bermudagrass and Bahiagrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass prior to spring green-up in areas where these turfgrasses are desirable ground covers and some temporary injury or discoloration can be tolerated.

Apply 5 to 44 fluid ounces of this product in 10 to 40 gallons of water per acre when bermudagrass and bahiagrass are dormant and prior to spring green-up.

Application of more than 11 fluid ounces of this product per acre on highly maintained bermudagrass and bahiagrass turf, such as golf courses and lawns, could result in injury or delayed green-up in the spring.

For residual weed control in dormant bermudagrass and bahiagrass, this product may be tank-mixed with Outrider (EPA Reg. No. 59639-223; *sulfosulfuron*), Oust Extra (EPA Reg. No. 432-1557; *sulfometuron methyl, metsulfuron methyl*), or Oust XP (EPA Reg. No. 432-1552; *sulfometuron methyl*) herbicides. Apply 5 to 44 fluid ounces of this product in a tank-mix with an appropriate rate of Outrider, Oust Extra or Oust XP herbicide in 10 to 40 gallons of water per acre. To avoid delays in green-up and minimize injury, apply no more than 1 ounce of Oust Extra or Oust XP herbicide per acre on bermudagrass and no more than 0.5 ounce on bahiagrass, and avoid application when these grasses are in a semi-dormant condition.

DO NOT apply this product in a tank-mix with Outrider, Oust Extra or Oust XP herbicides on highly maintained bermudagrass and bahiagrass turf, such as on golf courses and lawns.

#### Weed Control in Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds in actively growing bermudagrass. Some bermudagrass injury could result from the application of this product, but the bermudagrass will recover under moist conditions once the effects of the product wear off. Use only on well-established bermudagrass where some temporary injury or discoloration can be tolerated.

Apply 11 to 32 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Use a lower application rate within this range when controlling annual weeds less than 4 inches tall (or runner length) and increase the rate towards the upper end of the range as weeds increase in size or as they approach flower or seedhead formation. At these application rates, this product will provide partial control of the following perennial weeds in actively growing bermudagrass:

- Bahiagrass
- Fescue, tall
- Trumpetcreeper
- Bluestem, silver
- Johnsongrass
- Vaseygrass

**PRECAUTIONS:** Applying more than 11 fluid ounces of this product per acre on highly maintained bermudagrass, such as on golf courses and lawns, could cause unacceptable turf injury and discoloration.

For a broader weed control spectrum in actively growing bermudagrass, this product may be tank-mixed with Outrider, Oust Extra or Oust XP herbicides. Apply these tank-mixtures only on well-established bermudagrass where some temporary injury or discoloration can be tolerated. Make no more than one application of this product in these tank mixtures in the same season, otherwise the bermudagrass could be severely injured.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Apply 5 to 22 fluid ounces of this product per acre in a tank-mix with Outrider herbicide for control or partial control of johnsongrass and other weeds listed on the Outrider herbicide label. Apply both products at a rate toward the upper end of the given ranges to control annual or perennial weeds greater than 6 inches tall.

Apply 11 to 22 fluid ounces of this product per acre in a tank-mix with Oust Extra or Oust XP herbicide to increase the spectrum of weed control to include weeds listed on those labels. Use a lower application rate of each product within the given ranges to control annual weeds listed on the labels that are less than 4 inches tall (or runner length) and increase the rates toward the upper end of the ranges as annual weeds increase in size and approach the flower or seedhead stage. This tank-mix will provide partial control of the following perennial weeds in actively growing bermudagrass:

- Bahiagrass
- Dallisgrass
- Fescue, tall
- Trumpetcreeper
- Bluestem, silver
- Dock, curly
- Johnsongrass
- Vaseygrass
- Broomsedge
- Dogfennel
- Poorjoe
- Vervain, blue

**PRECAUTIONS:** Apply these tank mixtures only on well-established bermudagrass where some temporary injury or discoloration can be tolerated.

**RESTRICTIONS:** DO NOT apply this product in a tank mixture with Outrider or Oust herbicides on highly maintained bermudagrass, such as on golf courses and lawns.

### **Weed Control in Actively Growing Bahiagrass**

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4 fluid ounces of this product in 10 to 40 gallons of water per acre 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches prior to seedhead emergence.

For growth suppression of bahiagrass for up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 1.5 to 3 fluid ounces per acre about 45 days later. Make no more than two growth suppression applications per year.

For broad spectrum weed control in actively growing bahiagrass, this product may be tank-mixed with Outrider, Oust Extra, or Oust XP herbicides.

Apply 1.5 to 3.5 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Outrider herbicide to control perennial weeds or annual weeds greater than 4 inches in height.

Apply 4 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP herbicide 1 to 2 weeks following an initial spring mowing to increase the spectrum of weed control in actively growing bahiagrass. Make this application only once per year.

**PRECAUTIONS:** Apply these tank mixtures only on well-established bahiagrass where some temporary injury or discoloration can be tolerated.

### **Turf Renovation**

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding until after determining if any re-growth of underground plant parts will occur. Where repeat applications are necessary, sufficient re-growth must be attained prior to re-application of this product. Summer or fall application of this product is the optimal timing for control of warm-season grasses, such as bermudagrass. For managed turfgrass, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray solution.

This product has no residual soil activity and will not affect plants, seed or sod planted back into the area after application.



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

A handheld sprayer may be used for spot application onto unwanted vegetation growing in existing turfgrass. Broadcast application or spot treatment using a handheld sprayer may be used to control sod remnants or other unwanted vegetation after sod is harvested.

**PRECAUTIONS:** Do not disturb soil or underground plant parts before application of this product and delay tillage and renovation techniques, such as vertical mowing, coring or slicing, a minimum of 7 days after application to allow translocation of the herbicide into underground plant parts.

**RESTRICTIONS:** If application rates total 2 quarts of this product per acre or less, no waiting period between application and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

### **Chemical Mowing**

This product may be used to suppress growth of perennial and annual grasses listed in this section to serve as a substitute for mowing.

Perennial Grasses – apply 4 fluid ounces of this product per acre to suppress growth of Kentucky bluegrass, or 5 fluid ounces to suppress tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass in 10 to 40 gallons of spray solution per acre after grasses have greened up to at least 75 percent green color in the spring, or 7 to 10 days after mowing when sufficient re-growth has occurred to provide a desirable height for growth regulation. Use chemical mowing only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Annual Grasses – apply 3 to 4 fluid ounces of this product in 10 to 40 gallons of spray solution per acre to suppress growth of some annual grasses, such as annual ryegrass, wild barley and wild oats when actively growing in coarse turf on roadsides or other industrial areas and before seedheads are in the boot stage of development. This application could injure the desired annual grasses.

**PRECAUTIONS:** Use this product for chemical mowing only in areas where some temporary injury or discoloration of perennial and annual grasses can be tolerated.

### **12.2 Annual Weeds**

Annual weeds are easiest to control with this product when they are small and actively growing. New leaf development indicates active growth.

To control or partially control the annual weeds listed in this section when they are less than 6 inches in height or runner length and actively growing, apply 22 fluid ounces of this product per acre. If they are over 6 inches in height or runner length or slowly growing under stressed conditions, increase the application rate to 1 to 2.7 quarts per acre, depending on weed height and severity of the poor growing conditions.

For application using a handheld sprayer with a spray-to-wet technique, apply a 0.4-percent solution of this product to annual weeds less than 6 inches in height or runner length prior to seedhead formation in grasses or bud formation in broadleaf weeds. To control annual weeds over 6 inches tall, or for even smaller weeds growing under stressed conditions, apply a 0.7- to 1.5-percent solution. Apply the maximum concentration of this product within this range to hard-to-control weeds or to control weeds over 24 inches tall.

[*Optional text:* For control of annual weeds using a handheld controlled droplet applicator (CDA), apply a 15-percent solution of this product (19 to 20 fluid ounces of this product per gallon of spray solution) at a flow rate of 2 fluid ounces of spray solution per minute and a walking speed of 1.5 miles per hour (1 quart of spray solution per acre). When using a vehicle-mounted CDA, apply the appropriate amount of this product in 2 to 15 gallons of water per acre.]

To maximize performance of this product, do not mow, cut, till, burn or disturb vegetation in the application area for a minimum of 3 days after application.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

This product has no residual soil activity and does not control emergence of new annual weeds from seed. Repeated application of this product will be needed to control weeds that continue to emerge.

### **ANNUAL WEED SPECIES**

---

Anoda, spurred  
Balsam apple<sup>1</sup>  
Barley  
Barley, little  
Barnyardgrass  
Bassia, fivehook  
Bittercress  
Bluegrass, annual\*  
Bluegrass, bulbous  
Brome, downy  
Brome, Japanese  
Broomsedge  
Buttercup  
Castor bean<sup>2</sup>  
Cheatgrass  
Cheeseweed (*Malva parviflora*)  
Chervil  
Chickweed  
Cocklebur  
Copperleaf, hophornbeam  
Copperleaf, Virginia  
Coreopsis, plains/tickseed  
Corn  
Crabgrass  
Cupgrass, woolly  
Dwarf dandelion  
Eclipta  
False dandelion  
Falseflax, smallseed  
Fiddleneck  
Filaree  
Fleabane, annual  
Fleabane, hairy\* (*Conyza bonariensis*)  
Fleabane, rough  
Foxtail  
Foxtail, Carolina  
Geranium, Carolina  
Goatgrass, jointed  
Goosegrass\*  
Groundsel, common  
Henbit  
Horseweed / Marestalk\* (*Conyza canadensis*)  
Itchgrass  
Johnsongrass, seedling\*  
Junglerice\*  
Knotweed  
Kochia\*

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Lambsquarters  
Lettuce, prickly  
Mannagrass, eastern  
Mayweed  
Medusahead  
Morning glory (*Ipomoea spp*)  
Mustard, blue  
Mustard, tansy  
Mustard, tumble  
Mustard, wild  
Nightshade, black  
Oats  
Panicum, browntop  
Panicum, fall  
Panicum, Texas  
Pennycress, field  
Pepperweed, Virginia  
Pigweed\*  
Puncturevine  
Purslane, common  
Pusley, Florida  
Ragweed, common\*  
Ragweed, giant\*  
Rice, red  
Rocket, London  
Rocket, yellow  
Rye  
Ryegrass\*  
Sandbur, field  
Sesbania, hemp  
Shattercane  
Shepherd's-purse  
Sicklepod  
Signalgrass, broadleaf  
Smartweed, ladythumb  
Smartweed, Pennsylvania  
Sorghum, grain (milo)  
Sowthistle, annual  
Spanish needles<sup>3</sup>  
Speedwell, corn  
Speedwell, purslane  
Sprangletop  
Spurge, annual  
Spurge, prostrate  
Spurge, spotted  
Spurry, umbrella  
Starthistle, yellow  
Stinkgrass  
Sunflower\*  
Teaweed / Prickly sida  
Thistle, Russian\*

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Velvetleaf  
Wheat  
Wild oats  
Witchgrass

---

- <sup>1</sup> For control of balsam apple, apply this product using handheld equipment only.
  - <sup>2</sup> Control of castor bean can also be achieved by injecting 4 milliliters of this concentrated (undiluted) product per plant into the lower portion of the main stem.
  - <sup>3</sup> For control of Spanish needles, apply 44 fluid ounces of this product per acre.
- \* A glyphosate-resistant biotype has been confirmed. For additional information, refer to the “WEED RESISTANCE MANAGEMENT” section of this label. You can also visit [www.weedscience.org](http://www.weedscience.org) or contact your Bayer CropScience representative.

### 12.3 Perennial Weeds

More effective control of perennial weeds can be obtained when this product is applied to target weeds that are small and actively growing. New leaf development indicates active growth. If application must be made to larger weeds or to weeds that are slowly growing under stressful conditions, apply this product at a rate or spray solution concentration towards the upper end of the specified range.

If weeds have been mowed or tilled, do not apply this product until plants have resumed active growth and have reached the specified stage of growth or sufficient growth has been achieved to allow for good interception of the spray solution. To maximize performance of this product, do not mow, cut, till, burn or disturb vegetation in the application area for a minimum of 7 days after application.

For control of perennial weeds listed on this label using backpack or handheld equipment and a low-volume application technique, apply a 4- to 7-percent solution of this product over the crown of the target plant to cover 50 percent of the upper plant foliage.

For control of perennial weeds using a handheld controlled droplet applicator (CDA), apply a 15 to 30-percent solution of this product (19 to 38 fluid ounces of this product per gallon of spray solution) at a flow rate of 2 fluid ounces of spray solution per minute and a walking speed of 0.75 mile per hour (2 to 4 quarts of spray solution per acre). When using a vehicle-mounted CDA, apply the required amount of this product, as indicated in the following table, in 2 to 15 gallons of water per acre.

Application of this product in the fall must be made before a killing frost.

This product has no soil activity and does not control emergence of perennial weeds from seed and dormant underground roots, rhizomes or tubers present in the soil at the time of application. Repeated application of this product will be necessary for continued control of weeds that emerge after application.

**PERENNIAL WEEDS RATE TABLE**

<b>Perennial Weed Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Sprayer Concentration (% Solution)</b>
<b>Alfalfa<sup>1</sup></b>	1 – 1.5	1.5
<b>Alligatorweed<sup>1</sup></b> Apply this product when most of the target plants are in bloom. More than one application will be needed to achieve control.	3	1
<b>Anise (fennel)</b>	1.3 – 2.7	1 – 1.5

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Perennial Weed Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Sprayer Concentration (% Solution)</b>
<b>Artichoke, Jerusalem</b>	2 - 3.3	1.5
<b>Bahiagrass</b>	2 – 3.3	1.5
<b>Beachgrass, European</b> ( <i>Ammophila arenaria</i> )	–	3.5
<p>Apply a 3.5-percent solution of this product using a spray-to-wet technique or an 8-percent solution using a low-volume application technique. More effective control can be obtained when application is made onto target weeds that are actively growing at the boot through the full-heading stage of development. Make application prior to the loss of more than 50 percent of green leaf color in the fall. Monitor application site and re-apply this product to any target weeds that were missed, if necessary, before re-seeding the area with desirable vegetation.</p> <p>For selective control of European beachgrass, apply a 33.3-percent solution of this product during period of active growth using a wiper applicator. Maximizing the amount of individual leaf tissue contacted by the wiper applicator or making a second pass through the field in the opposite direction will improve control. Avoid contact of the herbicide solution with desirable vegetation.</p>		
<b>Bentgrass<sup>1</sup></b>	1	1.5
<p>This product alone will provide only partial control of bentgrass (<i>Agrostis spp.</i>). For better control, apply 1.6 to 2.2 quarts of this product in a tank-mix with an appropriate rate of clethodim, fluazifop-P-butyl, fenoxaprop-P-butyl, or sethoxydim in a spray volume of 20 to 40 gallons per acre using broadcast application equipment. When using a handheld sprayer, apply this product at a concentration of 1.5 fluid ounces per gallon in a tank-mix with an appropriate concentration of one or more of these active ingredients. More than one application might be needed for complete control.</p>		
<b>Bermudagrass</b>	3.3	1.5
<p>Make application when seedheads are present.</p>		
<b>Bermudagrass, water (knotgrass)</b>	1	1.5
<b>Bindweed, field</b>	2.7 – 3.3	1.5
<p>For control, apply 2.7 to 3.3 quarts of this product per acre as a broadcast application west of the Mississippi River and 2 to 2.7 quarts per acre east of the Mississippi River when bindweed is at or beyond full bloom. For maximum performance, apply in late-summer or fall.</p>		
<b>Bittersweet, Oriental</b>	2	1.5
<p>For control, apply this product as a broadcast application in 30 to 40 gallons of spray solution per acre. To maximize performance, apply this product in late-summer or fall while leaves are still green and after fruit formation. When using handheld application equipment and a spray-to-wet technique, ensure complete coverage of the target plant with the spray solution.</p>		
<b>Bluegrass, Kentucky</b>	1.5	1.5
<p>Apply when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.</p>		
<b>Blueweed, Texas</b>	2.7 – 3.3	1.5
<p>Apply 2.7 to 3.3 quarts of this product per acre west of the Mississippi River and 2.3 to 3 quarts per acre east of the Mississippi River when most target plants are at or beyond full bloom. For maximum performance apply in late-summer or fall.</p>		
<b>Brackenfern</b>	2 – 3	1
<p>Apply to fully expanded fronds that are at least 18 inches long</p>		

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Perennial Weed Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Sprayer Concentration (% Solution)</b>
<b>Bromegrass, smooth</b> Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.	1.5	1.5
<b>Bursage, woolly-leaf</b>	–	1.5
<b>Canarygrass, reed</b> Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.	1.5 – 2	1.5
<b>Cattail</b> Apply this product when target plants are actively growing and are at or beyond the early to full bloom stage of development. For maximum performance, apply this product during the summer or fall months.	2 – 3.3	1.5
<b>Clover; red, white</b>	2 – 3.3	1.5
<b>Cogongrass</b> Apply this product in late-summer or fall when cogongrass is at least 18 inches tall and actively growing. Due to uneven stages of growth and the dense nature of cogongrass vegetation, more than one application might be necessary to achieve control.	2 – 3.3	1.5
<b>Dallisgrass</b>	2 – 3.3	1.5
<b>Dandelion</b>	2 – 3.3	1.5
<b>Dock, curly</b>	2 – 3.3	1.5
<b>Dogbane, hemp</b> Apply this product when most target plants have reached the late-bud to flower stage of growth. For maximum performance, make application in late-summer or fall.	2.5	1.5
<b>Fescue (except tall)</b>	3	1.5
<b>Fescue, tall</b> Apply this product when most target plants have reached the boot to head stage of growth. If applied prior to the boot stage, less than desirable control might be obtained.	2	1.5
<b>Guinea grass</b> Apply this product when most target plants have reached at least the 7-leaf growth stage.	2	1
<b>Hemlock, poison</b> Control can also be achieved by injecting 5 milliliters of a 5-percent solution of this product using a handheld injection device in one leaf cane per plant, 12 inches above the root crown. <sup>2</sup>	1.3 – 2.7	1 – 1.5
<b>Hogweed, giant</b> Inject 5 milliliters of a 5-percent solution of this product into one leaf cane per plant, 12 inches above the root crown. <sup>2</sup>	–	–
<b>Horsenettle</b>	2 – 3.3	1.5
<b>Horseradish</b> Apply this product when most target plants have reached the late-bud to flower stage of development. For maximum performance, apply in late-summer or fall.	3	1.5
<b>Horsetail, field</b>	–	–

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Perennial Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Handheld Sprayer Concentration</b> (% Solution)
Inject 0.5 milliliter of this product per stem directly into the plant stem, one segment above the root crown. <sup>2</sup>		
<b>Iceplant</b>	1.3	1.5 – 2
<b>Ivy; cape, German</b>	1.3 – 2.7	1 – 1.5
<b>Jerusalem artichoke</b>	2 – 3.3	1.5
<b>Johnsongrass</b> Apply this product when most target plants have reached the boot to head stage of development or before plants have turned brown in the fall. When applied prior to the boot stage, reduced control can result.	1.3 – 2	1
<b>Kikuyu grass</b>	1.5 – 2	1.5
<b>Knapweed</b> Apply this product when most target plants have reached the late-bud to flower stage of growth. For maximum performance, make application in late-summer or fall.	3	1.5
<b>Knotweed; Bohemian, giant, Japanese</b> Apply 2.7 quarts of this product per acre as a broadcast application in 3 to 40 gallons of spray solution. For application using a backpack sprayer and a spray-to-wet technique, apply a 2-percent solution of this product. For maximum performance, do not disturb vegetation in the application area for a minimum of 7 days after application.  Control can also be achieved by cutting stems cleanly just below the 2nd or 3rd node above the ground and immediately apply 0.36 fluid ounce (10 milliliters) of a 50-percent solution of this product in water into the "well" or remaining internode. Ensure that the upper plant material that was removed is gathered and properly discarded to prevent new plants from propagating from sprouting buds. Use of a bio-barrier, such as cardboard, plywood or plastic sheeting, will help guard against the spread of plant material. The combined total application rate of this product must not exceed 6 quarts per acre. <sup>2</sup>  Control can also be achieved by injecting 5 milliliters of this concentrated (undiluted) product per stem into the second or third internode using a handheld injection device. <sup>2</sup>	2.7	2
<b>Lantana</b> Apply this product when most target plants are at or beyond the bloom stage of growth. Use the higher spray solution concentration on plants that have reached the woody stage of growth.	–	1
<b>Lespedeza</b>	2 – 3.3	1.5
<b>Loosestrife, purple</b> Apply this product when most target plants are at or beyond the bloom stage of growth. Maximum performance can be achieved when this product is applied during summer or fall months. Fall application must be made before a killing frost.	1.75	1 – 1.5
<b>Lotus, American</b> Apply this product when most target plants are at or beyond the bloom stage of growth. Maximum performance can be achieved when application is made during summer or fall months. Fall application must be made before a killing frost. More than one application of this product might be necessary to control re-growth of underground plant parts and seeds.	1.75	0.75
<b>Milkweed, common</b> Apply this product when most target plants have reached the late-bud to flower stage of growth.	2	1.5
<b>Muhly, wirestem</b>	1.5	1.5

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Perennial Weed Species</b>	<b>Broadcast Rate</b> (quarts/acre)	<b>Handheld Sprayer Concentration</b> (% Solution)
Make application when most target plants are at least 8 inches in height (3 to 4-leaf stage of development) and actively growing.		
<b>Mullein, common</b>	2 – 3.3	1.5
<b>Napiergrass</b>	2 – 3.3	1.5
<b>Nightshade, silverleaf</b>	1.5	1.5
Apply 2.7 to 3.3 quarts of this product per acre as a broadcast application west of the Mississippi River and 2 to 2.7 quarts per acre east of the Mississippi River when most target plants are at or beyond full bloom. Maximum performance can be obtained when application is made in late-summer or fall after berries have formed.		
<b>Nutsedge; purple, yellow</b>	2	1 – 1.5
Apply this product to control existing nutsedge plants and attached immature nutlets when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and will require repeated application of this product for long-term control.		
<b>Orchardgrass</b>	1.5	1.5
Make application when most target plants have reached the boot to head stage of development. When applied prior to the boot stage, less than acceptable control could be obtained. In the fall, make application before plants have turned brown.		
<b>Pampas grass</b>	2 – 3.3	1 – 1.5
<b>Para grass</b>	2 – 3.3	1.5
More than one application of this product will be needed to achieve complete control. Allow plants to re-grow to the 7- to 10-leaf stage before making next application.		
<b>Pepperweed, perennial</b>	2.7	1.5
<b>Phragmites<sup>1</sup></b>	2 – 3.3	1 – 1.5
For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 3.3 quarts of this product per acre as a broadcast application or a 1.5-percent solution using a handheld sprayer. In other areas of the U.S., apply 1.75 to 2.7 quarts per acre as a broadcast application or, for partial control, apply a 0.75-percent solution using a handheld sprayer. For maximum performance, make application in late-summer or fall when plants are actively growing and in full bloom. Due to the dense nature of this vegetation (which can prevent good spray coverage) and uneven stages of growth, more than one application of this product might be necessary to achieve control. Visual symptoms of control will be slow to develop.		
<b>Pokeweed, common</b>	1	1.5
Apply to actively growing target plants up to 24 inches tall.		
<b>Quackgrass</b>	1.3 – 2	1.5
Apply this product when most target plants are at least 8 inches in height (3- to 4-leaf stage of development) and actively growing.		
<b>Redvine<sup>1</sup></b>	1.5	1.5
<b>Reed; common, giant</b>	2.7 – 3.3	1.5
For maximum performance, apply this product in late-summer or fall. Control can also be achieved by injecting 5 milliliters of this concentrated product (undiluted) directly into the second or third internode using a handheld injection device. <sup>2</sup>		



II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Perennial Weed Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Sprayer Concentration (% Solution)</b>
<b>Ryegrass, perennial</b> Apply this product when most target plants have reached the boot to head stage of growth. When applied prior to the boot stage, reduced control can result. In the fall, make application before ryegrass turns brown.	1.5 – 2	1
<b>Smartweed, swamp</b>	2 – 3.3	1.5
<b>Spatterdock</b> Make application when most target plants are in full bloom. For maximum performance, make application in the summer or fall.	2.7	0.75
<b>Sowthistle, perennial</b>	1.5 – 2	1.5
<b>Spurge, leafy<sup>1</sup></b>	–	1.5
<b>Starthistle, yellow</b>	1.5	1.5
<b>Sweet potato, wild<sup>1</sup></b> Make application when most target plants are at or beyond the bloom stage of growth. More than one application will be needed to achieve control.	–	1.5
<b>Thistle, artichoke</b> Make application when target plants are at or beyond the bud stage of growth.	1.3 – 2	1 – 1.5
<b>Thistle, Canada</b> Make application when target plants are at or beyond the bud stage of growth. Control can also be achieved by stem-injection. Cut 8 to 9 of tallest plants in a clump at bud stage. Push a cavity needle into the stem center and then slowly remove it as you inject 0.5 milliliter of this concentrated (undiluted) product into the stem. <sup>2</sup>	1.5 – 2	1.5
<b>Timothy</b> Make application when most target plants have reached the boot to head stage of development. If application is made prior to the boot stage, reduced control can result. In the fall, make application before plants turn brown.	1.5 – 2	1.5
<b>Torpedograss<sup>1</sup></b>	2.7 – 3.3	1.5
<b>Trumpet creeper<sup>1</sup></b>	1.5 – 2	1.5
<b>Tules, common</b> Make application to target plants at or beyond the seedhead stage of development. Visual symptoms will be slow to appear and might not appear for 3 or more weeks after application.	–	1.5
<b>Vaseygrass</b>	2 – 3.3	1.5
<b>Velvetgrass</b>	2 – 3.3	1.5
<b>Wheatgrass, western</b> Make application when most target plants have reached the boot to head stage of development. Application made prior to the boot stage could result in reduced control. In the fall, make application before plants turn brown.	1.5 – 2	1.5

<sup>1</sup> Partial control only

<sup>2</sup> When using stem injection, the combined total use of this product must not exceed 7 quarts per acre per year. At 5 milliliters of concentrated (undiluted) product per stem, 7 quarts of this product will treat

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

approximately 1300 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

### 12.4 Woody Brush, Trees and Vines

Apply this product to brush, trees and vines that are actively growing after full leaf expansion, unless otherwise directed. Apply this product at a rate or spray solution concentration towards the upper end of a given range to control larger brush and trees, and/or for application in areas of dense vegetative growth, or for the control of vines that have reached the woody stage of growth.

In most areas, maximum performance of this product on woody brush, trees and vines can be obtained when application is made in late-summer or fall after fruit formation. In arid areas, application of this product in the spring to early-summer when brush and trees are at high moisture content and flowering could provide better results. Poor control can be expected when this product is applied to drought-stressed brush and trees.

When applying this product using a spray-to-wet technique with a handheld sprayer to control harder-to-control woody brush and trees, use a 1.5-percent solution of this product.

Some autumn color on undesirable deciduous species on brush and trees is acceptable when applying this product in the fall, provided no major leaf drop has occurred. Reduced performance of this product could result if application is made after a frost. Symptoms might not appear prior to frost or senescence following a fall application.

Repeat applications of this product might be required to control plants regenerating from underground parts or from seed.

For maximum performance, allow 7 or more days after application of this product before mowing, cutting, tilling, burning or removal of woody brush, trees or vines from that application site. Additional applications of this product will be needed to control brush and trees regenerating from underground parts or seed.

**TANK MIXTURES:** This product may be applied at any rate stated on this label in a tank mixture with products containing one or more of the active ingredients listed below [*Optional text*; or one or more of the products listed,] to increase the spectrum of control of herbaceous weeds, woody brush, trees and vines. For control of herbaceous weeds, apply the tank-mix product at the lower end of the given application rate or spray solution concentration range. For control of dense stands of vegetation or for hard-to-control woody brush, trees and vines, increase the application rate or spray solution concentration of the tank-mix product towards the higher end of the range. Refer to the individual product labels for approved uses and application rates. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Read and follow label directions for each product in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of any product in the mix. [*Optional text, only required if the product information is not included in the list below:* For more information on the products listed below, see the "TANK-MIX PRODUCT INFORMATION" section of this label.]

---

*[Active ingredients and products added to the final printed label for tank-mixing may be selected from the following list:*

imazapyr; metsulfuron methyl; triclopyr

Arsenal; Arsenal Herbicide Applicators Concentrate; Escort XP; Forestry Garlon 4 Specialty; Forestry Garlon XRT Specialty; Garlon 3A Specialty; Garlon 4 Specialty; Garlon 4 Ultra Specialty; Vastlan Specialty]

---

Ensure that any product containing triclopyr is thoroughly mixed with water in the spray tank according to label directions before adding this product to the spray mixture.

## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

### Cut Stump Application

This product may be used to control re-growth and re-sprouting of woody brush and trees on any site listed on this label.

Cut the woody brush or tree close to the soil surface and immediately apply a 50 to 100-percent (undiluted) solution of this product to the freshly-cut surface using an applicator capable of applying this product to the entire cambium. A delay in application could result in reduced performance. For best results, cut the woody brush or tree during period of active growth and full leaf expansion and apply this product.

For control of the tree of heaven (*Ailanthus altissima*), cut the tree close to the soil surface and immediately apply a 50-percent solution of this product (16 fluid ounces per quart of solution) and 10 percent Arsenal herbicide (3 to 4 fluid ounces per quart of solution) in water to the freshly-cut surface.

DO NOT MAKE A CUT STUMP APPLICATION WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MIGHT BE GRAFTED TO THE ROOTS OF THE CUT STUMP, AS INJURY TO THE ADJACENT TREES COULD OCCUR. Some sprouts, stems, or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indicator of a shared root system. Whether grafted or shared, injury is likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing a common root system.

### Woody Brush and Tree Injection and Frill Application

This product may be used to control woody brush and trees listed in this section by injection or frill application on any site listed on this label.

Inject or apply the equivalent of 1 milliliter (0.04 fluid ounce) of this concentrated product for every 2 to 3 inches of trunk diameter at breast height (DBH). If injecting this product into the woody brush or tree, use equipment capable of penetrating into the living plant tissue under the bark.

For frill application, apply a 50 to 100-percent (undiluted) solution of this product in water to either a continuous frill around the tree or to cuts evenly spaced around the tree below all branches. As tree diameter increases, better results can be achieved by applying this product to a continuous frill or to more closely spaced cuttings. Avoid application techniques that allow runoff of this product to occur from frilled or cut areas. In species that freely exude sap, make the frill or cuts at an oblique angle to produce a cupping effect and apply this concentrated product undiluted. For maximum performance, make this application during period of active growth and after full leaf expansion.

### Modified High-Volume and Low-Volume Backpack Application

For control and partial control of woody bush, trees and vines listed on this label when using a backpack sprayer or other handheld equipment and a directed low-volume foliar application technique, apply a 4 to 8-percent solution of this product evenly over the plant crown to cover 50 percent of the upper foliage of undesirable woody brush, trees and vines.

**WOODY BRUSH, TREES AND VINES RATE TABLE**

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% Solution)
Alder	2 – 3	1
Ash <sup>1</sup>	1.5 – 3.3	1 – 1.5
Aspen, quaking	1.5 – 2	1
Bearclover (Bearmat) <sup>1</sup>	1.5 – 3.3	1 – 1.5

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Spray-to-Wet Concentration (% Solution)</b>
<b>Beech<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Birch</b>	1.5 – 2	1
<b>Blackberry</b>	2 – 3	1
<b>Blackgum</b>	1.5 – 3.3	1 – 1.5
<b>Bracken</b>	1.5 – 3.3	1 – 1.5
<b>Broom; French, Scotch</b>	1.3 – 3.3	1 – 1.5
<b>Buckwheat, California<sup>1</sup></b>	1.3 – 2.5	1 – 1.5
<b>Cascara<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Castor bean</b>	–	–
Also for control, inject 4 milliliters of this concentrated (undiluted) product per plant directly into the lower portion of the main stem using a handheld injection device. <sup>2</sup>		
<b>Catsclaw<sup>1</sup></b>	–	1
For partial control, apply this product when at least 50 percent of the new leaves are fully developed.		
<b>Ceanothus<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Chamise<sup>1</sup></b>	1.3 – 3.3	1
<b>Cherry; bitter, black, pin</b>	1.5 – 2	1
<b>Coyote brush</b>	2 – 2.7	1 – 1.5
For control, apply this product when at least 50 percent of the new leaves are fully developed.		
<b>Deerweed</b>	1.3 - 3.3	1
<b>Dogwood<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Elderberry</b>	1.5 – 2	1
<b>Elm*</b>	1.5 – 3.3	1 – 1.5
<b>Eucalyptus</b>	–	1.5
For control of eucalyptus re-sprouts, apply this product using a handheld sprayer when re-sprouts are 6 to 12 feet tall. Ensure complete coverage.		
<b>Gallberry</b>	1.5 – 3.3	1 – 1.5
<b>Gorse<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Hackberry, western</b>	1.5 – 3.3	1 – 1.5
<b>Hasardia<sup>1</sup></b>	1.3 – 2.5	1 – 1.5
<b>Hawthorn</b>	1.5 – 2	1
<b>Hazel</b>	1.5 – 2	1
<b>Hickory<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Honeysuckle</b>	2 – 3	1
<b>Hornbeam, American<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Ivy, poison</b>	2.5 – 3.3	1.5

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

<b>Species</b>	<b>Broadcast Rate (quarts/acre)</b>	<b>Handheld Spray-to-Wet Concentration (% Solution)</b>
<b>Kudzu</b>	2.5 – 3.3	1.5
<b>Locust, black<sup>1</sup></b>	1.5 – 2.5	1 – 1.5
<b>Madrone (re-sprouts)<sup>1</sup></b>	–	1.5
<b>Manzanita<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Maple, red</b> For control, apply a 1-percent solution of this product using a handheld sprayer when leaves are fully developed. For partial control, apply 1.5 to 3 quarts per acre as a broadcast application.	1.5 – 3	1
<b>Maple, sugar</b> For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.	–	1
<b>Maple, vine<sup>1</sup></b>	1.3 – 3.3	1
<b>Monkey flower<sup>1</sup></b>	1.3 – 2.7	1 – 1.5
<b>Oak; black, white<sup>1</sup></b>	1.5 – 3	1 – 1.5
<b>Oak; northern, pin</b> For control, apply this product when at least 50 percent of the new leaves are fully developed.	1.3 – 2.7	1
<b>Oak, poison</b> Repeat applications might be necessary to maintain control. Application in the fall must be made before leaves lose green color.	2.5 – 3.3	1.5
<b>Oak, post</b>	2 – 3	1
<b>Oak, red</b> For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.	–	1
<b>Oak, scrub<sup>1</sup></b>	1.3 – 2.7	1
<b>Oak, southern red</b>	1.5 – 2	1
<b>Orange, Osage</b>	1.2 – 3.3	1 – 1.5
<b>Peppertree, Brazilian (Florida holly)<sup>1</sup></b>	1.3 – 3.3	1 – 1.5
<b>Persimmon<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Pine</b>	1.5 – 3.3	1 – 1.5
<b>Poplar, yellow<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Redbud, eastern</b>	1.5 – 3.3	1 – 1.5
<b>Rose, multiflora</b> Make application prior to leaf deterioration by leaf-feeding insects.	1.5	1
<b>Russian olive<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Sage, black</b>	1.3 – 2.7	1
<b>Sage, white<sup>1</sup></b>	1.5 – 2.7	1 – 1.5
<b>Sagebrush, California</b>	1.3 – 2.7	1
<b>Salmonberry</b>	1.5 – 2	1

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% Solution)
<b>Saltcedar<sup>1</sup></b> For partial control, apply a 1 to 1.5-percent solution of this product using a handheld sprayer or 1.5 to 3.3 quarts per acre as a broadcast application. For control, apply a 1 to 1.5-percent solution of this product in a tank-mix with an appropriate concentration of imazapyr [ <i>Optional alternative text:</i> Arsenal herbicide (EPA Reg. No. 241-346; <i>imazapyr</i> ) or Arsenal Herbicide Applicator's Concentrate (EPA Reg. No. 241-299; <i>imazapyr</i> )] using a handheld sprayer. For control using broadcast application, apply 1.3 quarts of this product per acre in a tank-mix with an appropriate rate of imazapyr to plants less than 6 feet tall. To control saltcedar greater than 6 feet tall using broadcast application, apply 2.75 quarts of this product per acre in a tank-mix with a higher application rate of imazapyr.	1.5 – 3.3	1 – 1.5
<b>Sassafras<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Sourwood<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Sumac; laurel, poison, smooth, sugarbush, winged<sup>1</sup></b>	1.5 – 3	1 – 1.5
<b>Sweetgum</b>	1.5 – 2	1
<b>Swordfern<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Tallowtree, Chinese</b>	–	1
<b>Tan oak (re-sprouts)<sup>1</sup></b>	–	1.5
<b>Thimbleberry</b>	1.5 – 2	1
<b>Tobacco, tree<sup>1</sup></b>	1.5 – 2.5	1 – 1.5
<b>Toyon<sup>1</sup></b>	–	1.5
<b>Trumpetcreeper</b>	1.5 – 2	1
<b>Virginia creeper</b>	1.5 – 3.3	1 – 1.5
<b>Waxmyrtle, southern<sup>1</sup></b>	1.5 – 3.3	1 – 1.5
<b>Willow</b>	2 – 3	1
<b>Yerba santa<sup>1</sup></b>	–	1.5

<sup>1</sup> Partial control

<sup>2</sup> When using stem injection, the combined total use of this product must not exceed 7 quarts per acre per year. At 4 milliliters of concentrated (undiluted) product per stem, 7 quarts of this product will treat approximately 1650 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

**13.0 TANK-MIX PRODUCT INFORMATION** [*this section printed only if required*]

[*This section is optional and only required if tank-mix product information is not included in the body of the label text. Products not included on the final printed label may be deleted from this list.*]

Brand Name	EPA Reg. No.	Active Ingredient(s)
AAtrex 4L	100-497	atrazine
AAtrex Nine-O	100-585	atrazine
Arsenal	241-346	imazapyr

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Brand Name	EPA Reg. No.	Active Ingredient(s)
Arsenal Herbicide Applicator's Concentrate	241-299	imazapyr
Banvel	66330-276	dicamba
Banvel 480	66330-421	dicamba
Barricade 4FL	100-1139	prodiamine
Barricade 65WG	100-834	prodiamine
Certainty Turf	59639-226	sulfosulfuron
Chopper	241-296	imazapyr
Chopper Gen2	241-430	imazapyr
Crossbow	62719-260	2,4-D, triclopyr
Direx 4L	66222-54	diuron
Endurance	100-834	prodiamine
Escort XP	432-1549	metsulfuron methyl
Forestry Garlon 4 Specialty	62719-40	triclopyr
Forestry Garlon XRT Specialty	62719-553	triclopyr
Formula 40	228-357	2,4-D
Garlon 3A Specialty	62719-37	triclopyr
Garlon 4 Specialty	62719-40	triclopyr
Garlon 4 Ultra Specialty	62719-527	triclopyr
Gallery 75 Dry Flowable Specialty	62719-145	isoxaben
Gallery SC	62719-658	isoxaben
Garlon XRT	62719-553	triclopyr
Goal 2XL	62719-424	oxyfluorfen
GoalTender	62719-447	oxyfluorfen
Habitat	241-426	imazapyr
Hyvar X	432-1546	bromacil
Hyvar X-L	432-1548	bromacil
Karmex DF	66222-51	diuron
Krenite S Brush Control Agent	42750-247	fosamine
Krovar 1 DF	5481-635	bromacil, diuron
Landmark XP	432-1560	sulfometuron methyl, chlorsulfuron
Oust Extra	432-1557	sulfometuron methyl, metsulfuron methyl
Oust XP	432-1552	sulfometuron methyl
Outrider	59639-223	sulfosulfuron
Pendulum 3.3 EC	241-341	pendimethalin
Pendulum AquaCap	241-416	pendimethalin
Plateau	241-365	imazapic
Poast	7969-58	sethoxydim
Poast Plus	7969-88	sethoxydim
Princep 4L	100-526	simazine
Princep Caliber 90	100-603	simazine

II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

Brand Name	EPA Reg. No.	Active Ingredient(s)
Princep Liquid	100-526	simazine
Ronstar 50 WSP	432-893	oxadiazon
Ronstar Flo	432-1465	oxadiazon
Ronstar G	432-886	oxadiazon
Sahara DG	241-372	imazapyr, diuron
Scythe	10163-325	pelargonic acid
Spike 20P Specialty	62719-121	tebuthiuron
Spike 80 DF Specialty	62719-107	tebuthiuron
Stalker	241-398	imazapyr
Surflan AS Agricultural	70506-43	oryzalin
Surflan AS Specialty	70506-44	oryzalin
Surflan Flex T&O	70506-308	oryzalin
Surflan XL 2G	70506-45	benefin, oryzalin
Telar XP	352-654 432-1561	chlorsulfuron
Tordon 101 Mixture Specialty	62719-5	2,4-D, picloram
Tordon 22K Specialty	62719-6	picloram
Tordon K Herbicide Specialty	62719-17	picloram
Transline Specialty	62719-259	clopyralid
Vanquish	228-397	dicamba
Vastlan Specialty	62719-687	triclopyr
Velpar DF CU	61842-48	hexazinone
Velpar DF VU	432-1576	hexazinone
Velpar L CU	61842-47	hexazinone
Velpar L VU	432-1573	hexazinone
Weedar 64	71368-1	2,4-D

**14.0 LIMIT OF WARRANTY AND LIABILITY**

Bayer CropScience (“Company”) warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label (“Directions”) and shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or



## II. DIRECTIONS FOR USE ON INDUSTRIAL, TURF AND ORNAMENTAL SITES

application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

[*Optional statement for agricultural-use products, if applicable:* For in-crop (over-the-top) uses on Roundup Ready and other glyphosate-tolerant crops, to the extent consistent with applicable law, crop safety and weed control performance are not warranted by Bayer CropScience when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.]

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

---

[Honcho K6 and Design, Roundup, Roundup Original, Roundup Original 2K and Design, Roundup Original MAX and Design, Trisorb and Design, and TRUEBLUE ADVANTAGE PROVEN RELIABLE SUPPORTED and Design] are [registered] trademarks of Bayer Group.

[*Optional text, if applicable:* All other trademarks are the property of their respective owners.]

---

For **MEDICAL** and **TRANSPORTATION** Emergency **ONLY** Call 24 hours a day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2987)

---

©[Year]

Packed for [*Alternative text:* Produced for] [*Alternative text:* Manufactured for]:

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

[*Insert print plate number*]

[*Insert barcode*]

Roundup Original 2K Herbicide (pending) 10/19/2023, 10/25/2023, 10/26/2023

# SUPPLEMENTAL LABELING

This supplemental label expires on May 1, 2025 and must not be used or distributed after this date.


READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Primary Brand Name:

**Roundup Original® 2K Herbicide**

Alternate Brand Names:

Honcho® K6 Herbicide  
Roundup Original® MAX Herbicide  
RT Master® II Herbicide



EPA Reg. No. 524-539

---

## USE WITH AGRISURE® GT CORN

---

Keep out of reach of children

### CAUTION

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

---

#### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

Read the label affixed to the [INSERT BRAND NAME] container before applying.

Use of [INSERT BRAND NAME] according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the [INSERT BRAND NAME] container.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® AND OTHER GLYPHOSATE-TOLERANT CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

See "PRODUCT INFORMATION" and "MIXING" sections of the [INSERT BRAND NAME] labeling for essential product information.

---

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Spot Treatment; Preharvest; Post-Harvest

<b>Maximum Application Rates</b>	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Maximum single In-crop application rate up to V8 stage or 30-inch corn	22 fluid ounces per acre
Total for all In-crop applications from emergence through V8 stage or 30-inch corn	44 fluid ounces per acre
Maximum Preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest	22 fluid ounces per acre

### **Preplant, At-Planting, Preemergence**

**USE INSTRUCTIONS:** This product may be applied alone or in a tank mixture before, during or after planting Agrisure GT corn.

**TANK MIXTURES:** This product may be tank-mixed with one or more products containing the active ingredients listed below. Apply these tank mixtures in 10 to 20 gallons of water, or 10 to 60 gallons of nitrogen solution, per acre. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. Read and follow all label directions. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

2,4-D; acetochlor; atrazine; bicyclopyrone; carfentrazone-ethyl; clopyralid; dicamba; diflufenzopyr; dimethenamid; dimethenamid-P; flufenacet; flumetsulam; flumiclorac pentyl ester; fluthiacet-methyl; isoxaflutole; linuron; mesotrione; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; rimsulfuron; saflufenacil; simazine; tembotrione; thiencazone-methyl

---

**RESTRICTIONS:** Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season. Application of 2,4-D or dicamba must be made a minimum of 7 days prior to planting corn.

**NOTE:** For maximum weed control, make a postemergence (in-crop) application of this product following the use of a preemergence residual product listed above.

### **Postemergence (In-crop)**

**USE INSTRUCTIONS:** This product may be applied alone or in a tank-mix over the top of Agrisure GT corn from emergence through the V8 stage (8 leaves with collars), or until corn plant height reaches 30 inches (freestanding), whichever comes first, unless otherwise directed. Maximum single in-crop application rate of this product is 22 fluid ounces per acre. Total in-crop application of this product from corn plant emergence through the V8 stage or 30 inches in height must not exceed 44 fluid ounces per acre.

When applied as directed, this product will control annual grasses and broadleaf weeds listed on the Honcho K6 Herbicide label. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of this product. Make a postemergence application of 16 to 22 fluid ounces of this product per acre before weeds exceed 4 inches in height (before they become competitive with the crop). Repeat this application before new flushes of weeds exceed 4 inches in height.

**TANK MIXTURES:** This product may be tank-mixed with a product containing the active ingredient acetochlor or atrazine. Ensure that the product used is labeled for application postemergence (in-crop) to field corn. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. Read and follow label directions for all products in the tank mixture. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**PRECAUTIONS:** In-crop applications made alone or with the addition of other crop chemical products could result in a crop response. Contact your seed trait provider for more information.

**RESTRICTIONS:** Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage or grain.

### **Preharvest**

**USE INSTRUCTIONS:** Up to 22 fluid ounces of this product per acre may be applied for annual and perennial weed control prior to harvest when kernel fill is complete, and the corn is physiologically mature (black layer formed) and grain moisture is 35 percent or less.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or feeding of corn stover or grain.

### **Post-Harvest**

**USE INSTRUCTIONS:** This product may be applied for weed control after crop harvest. Higher rates might be needed for control of large weeds that were growing in the field at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Ensure that the product used is labeled for post-harvest application in field corn. Read and follow label directions for all products in the tank mixture.

**RESTRICTIONS:** Allow a minimum of 7 days between application and harvest or the feeding of vegetation within the application area. Application must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

---

Read the "Limit of Warranty and Liability" on the [INSERT BRAND NAME] product labeling before using. These terms apply to this supplemental labeling. If these terms are not acceptable, return the product unopened at once.

---

[INSERT BRAND NAME] and Roundup Ready are [registered] trademarks of Bayer CropScience LP. Agrisure is a [registered] trademark of a Syngenta Group Company.

©[YEAR]

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

[INSERT DATE]

[PRINT PLATE NUMBER]

## SUPPLEMENTAL LABELING

This supplemental label expires on May 1, 2025 and must not be used or distributed after this date.


READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Primary Brand Name:

**Roundup Original® 2K Herbicide**

Alternate Brand Names:

Honcho® K6 Herbicide  
Roundup Original MAX® Herbicide  
RT Master® II Herbicide



EPA Reg. No. 524-539

---

### USE WITH ENLIST E3™ SOYBEAN

---

Keep out of reach of children

## CAUTION

In case of an emergency involving this product, call collect, day or night, (314) 694-4000.

---

#### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

Read the label affixed to the [INSERT BRAND NAME] container of before applying.

Use of [INSERT BRAND NAME] according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the [INSERT BRAND NAME] container.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® AND OTHER GLYPHOSATE-TOLERANT CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

See "PRODUCT INFORMATION" and "MIXING" sections of the [INSERT BRAND NAME] labeling for essential product information.

---

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest; Post-Harvest

<b>Maximum Application Rates</b>	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking through flowering (R2 stage soybean)	64 fluid ounces per acre
Maximum Preharvest application rate	22 fluid ounces per acre

### **Preplant, At-Planting, Preemergence**

USE INSTRUCTIONS: This product may be applied before, during or after planting Enlist E3 soybean.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and applied prior to planting only. This product may be tank-mixed with one or more products containing the active ingredients listed below and applied prior to crop emergence. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. Read and follow all label directions. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

acetochlor; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; saflufenicil; quizalofop-p-ethyl; saflufenacil; sulfentrazone; thifensulfuron; tribenuron methyl; trifluralin

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### **Postemergence (In-crop)**

USE INSTRUCTIONS: This product may be used to control annual grasses and broadleaf weeds in Enlist E3 soybean from emergence (cracking) through flowering (R2 stage soybean). R2 stage soybean ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). Refer to the "ANNUAL WEEDS RATE SECTION" of the [INSERT BRAND NAME] label for application rates for specific annual weeds.

An initial application of 22 fluid ounces of this product per acre will control or suppress most 2 to 8-inch tall weeds, which are normally found approximately 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be applied up to 44 fluid ounces per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

Application of 22 to 44 fluid ounces of this product per acre (single or multiple applications) will control or suppress perennial weeds, including bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestalk (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For maximum performance, allow perennial weed species to achieve at least 6 inches of growth before applying this product.

Under adverse growing conditions, including drought, hail or wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product might be necessary to control late flushes of weeds.

IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE ENLIST E3 SOYBEAN CROP. To control giant ragweed, apply 22 fluid ounces of this product per acre when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

PRECAUTIONS: In-crop applications made alone or with the addition of other crop chemical products could result in a crop response. Contact your seed trait provider for more information.

RESTRICTIONS: The combined total application of this product from crop emergence through harvest must not exceed 64 fluid ounces per acre. The maximum rate for any single in-crop application is 44 fluid ounces per acre. The maximum combined total amount of this product that may be applied during flowering (R2 stage soybean) is 44 fluid ounces per acre.

#### **Preharvest**

USE INSTRUCTIONS: Up to 22 fluid ounces of this product per acre may be applied to Enlist E3 soybean for weed control prior to harvest after pods have set and lost all green color. Take care to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Allow a minimum of 14 days between application and harvest of soybean grain or feeding of soybean grain, forage or hay.

#### **Post-Harvest**

USE INSTRUCTIONS: This product may be applied for weed control after harvest of Enlist E3 soybean. Higher rates might be needed for control of large weeds that were growing in the field at the time of harvest.

RESTRICTIONS: Application of this product must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

---

Read the "Limit of Warranty and Liability" on the [INSERT BRAND NAME] product labeling before using. These terms apply to this supplemental labeling. If these terms are not acceptable, return the product unopened at once.

---

[INSERT BRAND NAME] and Roundup Ready are [registered] trademarks of Bayer CropScience LP. Enlist E3 is a [registered] trademark of Corteva Company.

©[YEAR]

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

[INSERT DATE]

[PRINT PLATE NUMBER]

# SUPPLEMENTAL LABELING

This supplemental label expires on May 1, 2025 and must not be used or distributed after this date.


READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Primary Brand Name:

**Roundup Original® 2K Herbicide**

Alternate Brand Names:

Honcho® K6 Herbicide  
Roundup Original® MAX Herbicide  
RT Master® II Herbicide



EPA Reg. No. 524-539

---

## USE WITH GT27™ SOYBEAN

---

Keep out of reach of children

### CAUTION

In case of an emergency involving this product, call collect, day or night, (314) 694-4000.

---

#### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

Read the label affixed to the [INSERT BRAND NAME] container before applying.

Use of [INSERT BRAND NAME] according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the [INSERT BRAND NAME] container.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, EXCEPT AS DIRECTED FOR USE ON ROUNDUP READY® AND OTHER GLYPHOSATE-TOLERANT CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

See "PRODUCT INFORMATION" and "MIXING" sections of the [INSERT BRAND NAME] labeling for essential product information.

---

TYPES OF APPLICATION: Preplant; At-Planting; Preemergence; Postemergence (In-crop); Preharvest; Post-Harvest



<b>Maximum Application Rates</b>	
Combined total per year for all applications	5.3 quarts per acre
Total for all Preplant, At-Planting, Preemergence applications	3.3 quarts per acre
Total for all In-crop applications from cracking through flowering (R2 stage soybean)	64 fluid ounces per acre
Maximum Preharvest application rate	22 fluid ounces per acre

### **Preplant, At-Planting, Preemergence**

USE INSTRUCTIONS: This product may be applied before, during or after planting GT27 soybean.

TANK MIXTURES: This product may be tank-mixed with 2,4-D and applied prior to planting only. This product may be tank-mixed with one or more products containing the active ingredients listed below and applied prior to crop emergence. Refer to the individual label of all products used in the tank mixture for approved uses and application rates. Read and follow all label directions. It is the responsibility of the pesticide user to ensure that the intended use is included on the label of all products added to a tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

---

acetochlor; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; saflufenicil; quizalofop-p-ethyl; saflufenacil; sulfentrazone; thifensulfuron; tribenuron methyl; trifluralin

---

RESTRICTIONS: Maximum quantity of this product that may be applied for all preplant, at-planting and preemergence applications combined is 3.3 quarts per acre per season.

### **Postemergence (In-crop)**

USE INSTRUCTIONS: This product may be used to control annual grasses and broadleaf weeds in GT27 soybean from emergence (cracking) through flowering (R2 stage soybean). R2 stage soybean ends when a pod 5 millimeters (3/16 inch) long appears at one of the four uppermost nodes on the main stem with a fully developed leaf (R3 stage). Refer to the "ANNUAL WEEDS RATE SECTION" of the [INSERT BRAND NAME] label for application rates for specific annual weeds.

An initial application of 22 fluid ounces of this product per acre will control or suppress most 2 to 8-inch tall weeds, which are normally found approximately 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be applied up to 44 fluid ounces per acre as a single, in-crop application for control of annual weeds and where dense weed populations exist.

Application of 22 to 44 fluid ounces of this product per acre (single or multiple applications) will control or suppress perennial weeds, including bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marehail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For maximum performance, allow perennial weed species to achieve at least 6 inches of growth before applying this product.

Under adverse growing conditions, including drought, hail or wind damage, or a poor soybean stand that slows or delays canopy closure, a sequential application of this product might be necessary to control late flushes of weeds. IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE GT27 SOYBEAN CROP. To control giant ragweed, apply 22 fluid ounces of this product per acre when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

PRECAUTIONS: In-crop applications made alone or with the addition of other crop chemical products could result in a crop response. Contact your seed trait provider for more information.

RESTRICTIONS: The combined total application of this product from crop emergence through harvest must not exceed 64 fluid ounces per acre. The maximum rate for any single in-crop application is 44 fluid ounces per acre. The maximum combined total amount of this product that may be applied during flowering (R2 stage soybean) is 44 fluid ounces per acre.

#### **Preharvest**

USE INSTRUCTIONS: Up to 22 fluid ounces of this product per acre may be applied to GT27 soybean for weed control prior to harvest after pods have set and lost all green color. Take care to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Allow a minimum of 14 days between application and harvest of soybean grain or feeding of soybean grain, forage or hay.

#### **Post-Harvest**

USE INSTRUCTIONS: This product may be applied for weed control after harvest of GT27 soybean. Higher rates might be needed for control of large weeds that were growing in the field at the time of harvest.

RESTRICTIONS: Application of this product must be made a minimum of 30 days prior to the planting of any crop not listed on this label.

---

Read the "Limit of Warranty and Liability" on the [INSERT BRAND NAME] product labeling before using. These terms apply to this supplemental labeling. If these terms are not acceptable, return the product unopened at once.

---

[INSERT BRAND NAME] and Roundup Ready are [registered] trademarks of Bayer CropScience LP. GT27 is a [registered] trademark of BASF Company.

©[YEAR]

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167

[INSERT DATE]

[PRINT PLATE NUMBER]