



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AND

OFFICE OF
PREVENTION, PESTICIDES
TOXIC SUBSTANCES

JUL 1 - 2009

Ms. Dawn Fee-White
Monsanto Company
1300 I Street, NW—Suite 450 East
Washington, DC 20005

Dear Ms. Fee-White:

Subject: Roundup ProMax Herbicide (Revise Master Label)
EPA Registration No. 524-579
Application Dated May 13, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided you make the following changes before you release the product for shipment.

1. On page 2, revise the fourth paragraph to read "Not all products **listed** on this label are registered for use in California."
2. On page 3 and 17, revise the Heading for Section 8.0 to read "Site and Use **Instructions**".
3. On page 10, under Cultural Considerations, revised the section by replacing the word "recommended" with the word "specified".
4. On page 12, under Tank Mixing, revise the first sentence of the second paragraph to read "When this label **lists** a tank mixture with a generic active ingredient..."
5. On page 13, under Colorants and Dyes, revise the last sentence to read "Use colorants or dyes according to the manufacturer's **directions**."
6. On page 27, under Utility Sites, revise the first sentence of the fourth paragraph to read "This product **also may be used** in preparing or establishing wildlife openings within these sites."
7. On page 43, Supplemental B, revise the first note to read "The **labeled** rate for this use will limit the level of control of certain species of weeds."

2/49

Page 2

EPA Registration No. 524-579

Submit one copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supersede all previously approved ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

James A. Tompkins
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)


MASTER LABEL FOR EPA REG. NO. 524-579

Registered Brand Names:

Roundup PROMAX Herbicide

Table of Contents for Master Label

I.	Label for Industrial, Turf and Ornamental Uses	3 - 38
II.	Supplemental Labeling	39 - 48



[INSERT BRAND NAME]

Herbicide

The complete broad-spectrum postemergence professional herbicide for non-crop, industrial, turf and ornamental weed control.

Complete Directions for Use

EPA Reg. No. 524-579

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product.

Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the **LIMIT OF WARRANTY AND LIABILITY** statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. REPACKAGING OF THIS PRODUCT FOR DISTRIBUTION OR SALE MAY BE CONDUCTED ONLY UNDER THE TERMS OF A WRITTEN CONTRACT WITH MONSANTO .

Non-Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

PRODUCT INFORMATION

- 1 1.0 INGREDIENTS**
- 2 2.0 IMPORTANT PHONE NUMBERS**
- 3 3.0 PRECAUTIONARY STATEMENTS**
 - 3.1 Hazards to Humans and Domestic Animals
 - 3.2 Environmental Hazards
 - 3.3 Physical or Chemical Hazards

DIRECTIONS FOR USE

- 4 4.0 STORAGE AND DISPOSAL**
- 5 5.0 GENERAL INFORMATION**

- 5.1 Weed Resistance Management
- 5.2 Management Recommendations for Glyphosate-Resistant Weed Biotypes
- 6 6.0 MIXING**
 - 6.1 Mixing with Water
 - 6.2 Tank Mixing
 - 6.3 Tank Mixing Procedure
 - 6.4 Mixing for Hand-Held Sprayers
 - 6.5 Colorants or Dyes
- 7 7.0 APPLICATION EQUIPMENT AND TECHNIQUES**
 - 7.1 Aerial Equipment
 - 7.2 Ground Broadcast Equipment
 - 7.3 Hand-Held or Backpack Equipment
 - 7.4 Selective Equipment
 - 7.5 Injection Systems
 - 7.6 CDA Equipment
- 8 8.0 SITE AND USE RECOMMENDATIONS**
 - 8.1 Cut Stump
 - 8.2 Forestry Site Preparation
 - 8.3 General Areas and Industrial Sites
 - 8.4 Turfgrass
 - 8.5 Habitat Management
 - 8.6 Hollow-Stem Injection
 - 8.7 Injection and Frill (Woody Brush and Trees)
 - 8.8 Non-Food Tree, Shrub or Vine Production Sites
 - 8.9 Parks, Recreational and Residential Areas
 - 8.10 Railroads
 - 8.11 Roadsides
 - 8.12 Rangelands
 - 8.13 Utility Sites
 - 8.14 Grass Seed or Sod Production
 - 8.15 Pastures
- 9 9.0 WEEDS CONTROLLED**
 - 9.1 Annual Weeds
 - 9.2 Perennial Weeds
 - 9.3 Woody Brush and Trees
- 10 10.0 LIMIT OF WARRANTY AND LIABILITY**

PRODUCT INFORMATION

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its potassium salt.....48.7%

OTHER INGREDIENTS:.....51.3%

100.0%

6/49

*Contains 660 grams per liter or 5.5 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its potassium salt. Equivalent to 540 grams per liter or 4.5 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No's. 5,668,085 and 6,365,551. Other patents pending. No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE, 1-800-332-3111.

IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT,

(314)-694-4000.

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES MODERATE EYE IRRITATION

Avoid contact with eyes or clothing. Avoid breathing vapor or spray mist.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JUL 1 - 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
524-579

FIRST AID: Call a poison control center or doctor for treatment advice.	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses if present after the first 5 minutes then continue rinsing eye.
<ul style="list-style-type: none"> • Have the product container or label with you when calling a poison control center or doctor, or going for treatment. • You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information. • This product is identified as [INSERT BRAND NAME], EPA Registration No. 524-579. 	

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product can be mixed, stored and applied using only stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto supplemental labeling. Supplemental labeling can be found on the Internet at www.agrian.com, www.cdms.net or www.greenbook.net, or obtained from your Authorized Monsanto Retailer or Monsanto Company Representative.

The following paragraph is reserved and will only appear on final printed labeling for products under this registration if and when required by EPA:

ENDANGERED SPECIES PROTECTION REQUIREMENTS: This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, shoes plus socks and chemical-resistant gloves made of any waterproof material.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

4.0 STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage and disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

[Alternate PESTICIDE DISPOSAL statement for transport vehicles only: To avoid wastes, empty as much product from this transport vehicle as possible for repackaging or use in accordance with label directions. If wastes cannot be avoided, offer remaining product or rinsate to a waste disposal facility or pesticide disposal program. All disposal must be in accordance with applicable federal, state and local regulations and procedures.]

CONTAINER HANDLING AND DISPOSAL:

See container label for container handling and disposal instructions and for refilling limitations.

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID CONTAINERS OF LESS THAN 1-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill this container.

[Alternate container statement: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to

temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer this container for recycling, if available. If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[*Alternate container disposal statement:* Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-ROUNDUP (1-800-768-6387). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 2.5-GALLON CONTAINERS AND OTHER CONTAINERS OF GREATER THAN 1-GALLON BUT EQUAL TO OR LESS THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.

[*Alternate container statement:* Nonrefillable container. Do not reuse or refill this container.]

Triple rinse or pressure rinse (or equivalent) this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-ROUNDUP (1-800-768-6387). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[*Alternate container disposal statement:* Then offer this container for recycling, if available. If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR NONREFILLABLE RIGID PLASTIC 30-GALLON CONTAINERS AND OTHER CONTAINERS OF GREATER THAN 5-GALLON CAPACITY]

Nonrefillable container. Do not reuse or refill this container.

[Alternate container statement: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state.]

Triple rinse or pressure rinse (or equivalent) this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-ROUNDUP (1-800-768-6387). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

[Alternate container disposal statement: Then offer this container for recycling, if available. If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.]

[Optional container label statement: Return Properly Rinsed Container to Monsanto for Recycling Contact: 1-800-ROUNDUP (1-800-768-6387)]

[CONTAINER HANDLING AND DISPOSAL STATEMENT AND REFILLING LIMITATION FOR ALL REFILLABLE CONTAINERS, EXCEPT TRANSPORT CONTAINERS]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning this container before refilling is the responsibility of the refiller. Cleaning this container before final disposal is the responsibility of the person disposing of the container.

To clean this container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer this container for

12
/49

recycling, if available. To obtain information about recycling refillable containers, contact Monsanto Company at 1-800-ROUNDUP (1-800-768-6387).

[Optional container label statement: Return Properly Rinsed Container to Monsanto for Recycling, Contact 1-800-ROUNDUP (1-800-768-6387)]

5.0 GENERAL INFORMATION

Product Description: This product is a postemergence, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant and no additional surfactant is needed or recommended.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of aboveground growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds, effects may not be visible for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate weed control.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Plants arising from unattached underground rhizomes or rootstocks of perennials that have not yet emerged at the time of application will not be affected by this herbicide and will continue to grow.

Maximum Application Rates: The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as tank mixtures, on a basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalents) does not exceed the maximum allowed. The combined total of all treatments must not exceed 7 quarts of this product (8 pounds of glyphosate acid) per acre per year. See the **INGREDIENTS** section of this label for necessary product information.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or have other unintended consequences.

5.1 Weed Resistance Management

GROUP	9	HERBICIDE
-------	---	-----------

Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain plants naturally resistant to Group 9 herbicides. Weed species resistant to Group 9 herbicides may be effectively managed utilizing another herbicide from a different Group or using other cultural or mechanical practices.

To minimize the occurrence of glyphosate-resistant biotypes observe the following general weed management recommendations:

- Scout your application site before and after herbicides applications.
- Control weeds early when they are relatively small.
- Incorporate other herbicides and cultural or mechanical practices as part of your weed control system where appropriate.
- Utilize the labeled rate for the most difficult weed in the site. Avoid tank-mixtures with other herbicides that reduce this product's efficacy (through antagonism) or tank mixture recommendations which encourage rates of this product below the labeled rate.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from site to site to minimize spread of weed seed.
- Use new commercial seed as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product on a particular weed to your Monsanto representative, local retailer, or county extension agent.

5.2 Management Recommendations for Glyphosate-Resistant Weed Biotypes

NOTE: Appropriate testing is critical in order to confirm weed resistance to glyphosate. Call 1-800-ROUNDUP (1-800-768-6387) or contact your Monsanto representative to determine if resistance in any particular weed biotype has been confirmed in your area, or visit on the Internet www.weedresistancemanagement.com or www.weedscience.org.

Directions for the control of biotypes confirmed to be resistant to glyphosate are made available on separately published supplemental labeling or Fact Sheets for this product and may be obtained from your local retailer or Monsanto representative.

Since the occurrence of new glyphosate-resistant weeds cannot be determined until after product use and scientific confirmation, Monsanto Company is not responsible for any losses that may result from the failure of this product to control glyphosate-resistant weed biotypes.

The following good weed management practices are recommended to reduce the spread of confirmed glyphosate-resistant biotypes:

- If a naturally occurring resistant biotype is present at your site, this product may be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.

- Cultural and mechanical control practices may also be used as appropriate.
- Scout treated sites after herbicide applications and control escapes, including resistant biotypes, before they set seed.
- Thoroughly clean equipment before leaving sites known to contain resistant biotypes.

6.0 MIXING

Spray solutions of this product can be mixed, stored and applied using only clean stainless steel, fiberglass, plastic or plastic-lined steel containers. Clean sprayer parts immediately after using this product by thoroughly flushing with water.

Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by State or local regulations.

6.1 Mixing with Water

NOTE: PRODUCT PERFORMANCE MAY BE SIGNIFICANTLY REDUCED IF WATER CONTAINING SOIL SEDIMENT IS USED AS CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

This product mixes readily with water. Mix spray solutions of this product as follows: Begin filling the mixing tank or spray tank with clean water. Add the proper amount of this product near the end of the filling process and mix gently. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foaming, mix gently, terminate by-pass and return lines at the bottom of the tank and, if necessary, use an anti-foam or defoaming agent.

6.2 Tank Mixing

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive precautionary statements for each product in the mixture.

When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, dicamba, diuron or pendimethalin the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank mixture. Refer to all individual product labels, supplemental labeling and fact sheets for all products in the tank mixture, and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions and use according to the most restrictive precautionary statements for each product in the tank mixture.

Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressed in this label. Mixing this product with herbicides or other materials not on this label may result in reduced performance.

6.3 Tank-Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the specified amount of this product.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50-mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance. Ensure that the specific tank mixture product is registered for application at the desired site.

Refer to the **Tank Mixing** and **GENERAL INFORMATION** sections for additional precautions.

6.4 Mixing for Hand-Held Sprayers

Prepare the desired spray volume by mixing the amount of this product indicated in the following table in water:

Spray Solution

Desired Volume	Amount of [INSERT BRAND NAME]					
	0.4%	0.7%	1.0%	1.5%	4%	7%
1 gal	0.5 oz	1 oz	1.3 oz	2 oz	5 oz	9 oz
25 gal	0.8 pt	0.7 qt	1 qt	1.5 qt	4 qt	7 qt
100 gal	1.6 qt	2.8 qt	1 gal	1.5 gal	4 gal	7 gal

2 tablespoons = 1 fluid ounce

For use in backpack, knapsack or pump-up sprayers, add the appropriate amount of this product, mixed with water in a larger container and then filling sprayer with the mixed solution.

6.5 Colorants or Dyes

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce product performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

APPLY SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

AVOID DRIFT. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Use the labeled rates of this herbicide in 3 to 25 gallons of water per acre unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by Monsanto for this product.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray

volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aircraft Maintenance; Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE. Maintaining an organic coating (paint) that meets aerospace specification MIL-C-38413 may prevent corrosion.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial application.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversions" sections of this label).

Controlling Droplet Size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind speed, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, setup equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not be made during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify temperature inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Apply this product when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

7.2 Ground Broadcast Equipment

Apply the labeled rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by Monsanto for this product. As density of weeds increases, spray volume should be increased within the specified range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check spray pattern for uniform distribution of spray droplets.

7.3 Hand-Held or Backpack Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For low-volume directed spray applications, spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure

adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

7.4 Selective Equipment

This product may be diluted in water and applied through shielded applicators, hooded sprayers, wiper applicators or sponge bars to weeds listed on this label growing in any non-crop site specified on this label.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION, AS SERIOUS INJURY OR DEATH OF THE DESIRABLE VEGETATION IS LIKELY TO OCCUR.

Application equipment used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation is likely to result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators

Wiper applicators are devices that physically wipe the appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Application equipment used over the top of desirable vegetation should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust the height of the applicator to ensure adequate contact with weeds. Weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations or when weed height varies dramatically. In these instances, repeat treatments may be necessary.

Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to provide adequate wiper saturation with the herbicide solution. Better results may be obtained when two applications are made in opposite directions.

Droplets, mist, foam or splatter of the herbicide solution settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of the wiper applicator.

Do not use wiper applicators when weeds are wet.

Mix only the amount of this product to be used during a 1-day period, as reduced product performance may result from the use of solutions held in storage. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators—Solutions ranging from 25 to 70 percent of this product in water may be used.

For Panel Applicators and pressure-feed systems—Solutions ranging from 25 to 100 percent of this product in water may be used.

When applied as directed, this product **CONTROLS** the following weeds:

- | | |
|-----------------|------------------|
| Corn, volunteer | Sicklepod |
| Panicum, Texas | Spanishneedles |
| Rye, common | Starbur, bristly |
| Shattercane | |

When applied as directed, this product **SUPPRESSES** the following weeds:

- | | |
|------------------------|-----------------|
| Beggarweed, Florida | Ragweed, common |
| Bermudagrass | Ragweed, giant |
| Dogbane, hemp | Smutgrass |
| Dogfennel | Sunflower |
| Guineagrass | Thistle, Canada |
| Johnsongrass | Thistle, musk |
| Milkweed | Vaseygrass |
| Nightshade, silverleaf | Velvetleaf |
| Pigweed, redroot | |

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically instructed in this label.

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount directed in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 SITE AND USE RECOMMENDATIONS

This product may be used in non-crop areas such as airports, apartment complexes, commercial sites, ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, public areas, railroads, rangeland, recreation areas, residential areas, rights-of-way, roadsides, schools, shadehouses, sports complexes, storage areas, substations, turfgrass areas, utility sites, warehouse areas and wildlife management areas.

This product may also be used in non-food crop sites such as Christmas tree farms, plant nurseries, sod or turf seed farms.

Unless otherwise specified, applications may be made to control any weeds listed in the **WEEDS CONTROLLED** section of this label.

8.1 Cut Stump

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100- percent solution of this product to the freshly cut surface **immediately after** cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Saltcedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow
Pine, Austrian	

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2 Forestry Site Preparation

This product is labeled for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product is also labeled for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

Use this product for site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

Use the higher rates of this product within the specified range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the specified range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the specified range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any labeled rate of this product may be used in a tank mix with the following products for forestry site preparation.

- Arsenal Applicators Concentrate
- Chopper
- Escort or Escort XP
- Garlon 3A
- Garlon 4

Landmark XP

Oust XP

Westar

For control of herbaceous weeds, use the lower labeled tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher labeled rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release unless otherwise specified on this label, or in separate supplemental labeling or fact sheets published by Monsanto for this product.

8.3 General Areas and Industrial Sites

General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general non-crop and non-food crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in these sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

TANK MIXTURES: This product may be tank mixed with the following products, provided that the specific product is labeled for application at the use site. Refer to the individual product labels for approved sites and application rates.

Arsenal	atrazine ¹	Certainty
Barricade 65WG	Crossbow L	dicamba ¹
diuron ¹	Endurance	Escort or Escort XP
Gallery 75DF	Garlon 3A	Garlon 4
Goal 2XL	Karmex DF	Krovar I DF
Landmark II MP	Landmark MP	Landmark XP
Milestone		Oust XP
Outrider®	Pendulum 3.3 EC	Pendulum WDG
pendimethalin ¹	Plateau	Poast
Princep DF	Princep 4L	Ronstar 50WP
Sahara	simazine ¹	Surflan AS
Surflan WDG	Telar	Transline
Vanquish	Velpar DF	Velpar L
2,4-D ¹		

¹ Tank mixtures with products containing this single active ingredient may be made provided the specific product is labeled for application at the use site.

This product plus dicamba tank mixtures may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 22 to 44 fluid ounces of this product plus 2 to 4 ounces of Oust XP per acre.

Bahiagrass	Johnsongrass
Bermudagrass	Poorjoe

122
49

Broomsedge
Dallisgrass
Dock, curly
Dogfennel
Fescue, tall

Quackgrass
Vaseygrass
Vervain, blue

Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 5 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 4 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing – Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

8.4 Turfgrass

Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring green-up.

Apply 5 to 44 fluid ounces of this product in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 11 fluid ounces per acre may result in injury or delayed green-up in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust XP in highly maintained turfgrass areas.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. DO NOT apply more than 11 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust XP in highly maintained turfgrass areas. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

PRECAUTIONS, RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts. If application rates total 2 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove livestock before application and wait 8 weeks after application before grazing or harvesting.

8.5 Habitat Management

Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.6 Hollow Stem Injection

This product may be applied through hand-held injection devices that deliver recommended amounts of this product into targeted hollow-stem plants growing in any site specified on this label.

For control of the following hollow-stem plants, follow the use instructions below:

Castorbean, *Ricinus communis*

Inject 4 ml per plant of this product into the lower portion of the main stem.

Hemlock, Poison, *Conium maculatum*

Inject one leaf cane per plant 10 to 12 inches above the root crown with 5 ml of a 5% v/v solution of this product.

Hogweed, Giant, *Hercleum mantegazzianum*

Inject one leaf cane per plant 12 inches above the root crown with 5 ml of a 5% v/v solution of this product.

Horsetail, Field, *Equisetum arvense*

Inject one segment above the root crown with 0.5 ml per stem of this product. Use a small syringe that calibrates to this rate.

Knotweed, Bohemian, *Polygonum bohemicum*

Inject 5 ml per stem of this product between the second and third internode.

Knotweed, Giant, *Polygonum sachalinense*

Inject 5 ml per stem of this product between second and third internode.

Knotweed, Japanese, *Polygonum cuspidatum*

Inject 5 ml per stem of this product between second and third internode.

Reed, Giant *Arundo donax*

Inject 6 ml per stem of this product between second and third internode.

Thistle, Canada, *Cirsium arvense*

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.5 ml per stem of this product is injected into the stem.

Note: The combined total for all treatments must not exceed 7 quarts of this product per acre. At 5 ml per stem, 7 quarts should treat approximately 1300 stems per acre.

8.7 Injection and Frill (Woody Brush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 1 ml of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50- to 100-percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow run-off to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100-percent concentration of this product. For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:

<u>Control</u>	<u>Partial Control</u>
Oak	Black gum
Poplar	Dogwood
Sweetgum	Hickory
Sycamore	Maple, red

8.8 Non-Food Tree, Shrub, or Vine Production Sites

This product may be used for general weed control prior to the planting of and around established ornamentals, or any woody tree, shrub or vine species, including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew, growing in plant nurseries, on Christmas tree farms, or on other non-food tree production sites.

UNLESS OTHERWISE DIRECTED, THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be taken to avoid contact of spray, drift or mist with foliage or bark of desirable ornamental species.

This product may also be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off until after the application has dried.

TYPES OF APPLICATION: Site Preparation, Post-directed, Trim-and-edge, Wiper Application

Site Preparation

This product may be used prior to planting any tree, shrub or vine, including Christmas tree species, in a nursery or production setting.

Post-Directed, Trim-and-Edge

This product may be used as a post-directed spray around established woody ornamental species, or to trim and edge around trees, buildings, sidewalks, roads, potted plants and other objects in a production setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

Wiper Application

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs or vines. See the **SELECTIVE EQUIPMENT** section of this label for further information about the proper use of wiper applicators.

8.9 Parks, Recreational and Residential Areas

All of the instructions in the **General Areas and Industrial Sites** section may be made in park and recreational areas.

This product may be used in parks, recreational and residential areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around trees, fences, paths, around buildings, sidewalks and other objects in these areas. This product may be used for spot treatment of unwanted vegetation. This product may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

8.10 Railroads

All of the instructions in the **General Areas and Industrial Sites** section may be made to railroads.

Bare ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

TANK MIXTURES: This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments provided the specific product used is labeled for use on these sites. Refer to the individual product labels for approved sites and application rates:

Arsenal	atrazine ¹
dicamba ¹	diuron ¹
Escort	Escort XP
Garlon 3A	Garlon 4
Hyvar X	Hyvar X-L
Krovar I DF	
Oust XP	Outrider
Sahara DG	simazine ¹
Spike 80DF	Telar DF
Transline	Vanquish
Velpar DF	Velpar L
2,4-D ¹	

¹ Tank mixtures with products containing this single generic active ingredient may be made provided the specific product is labeled for application at the use site.

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 2.5 to 7 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.7- to 1.5-percent solution of this product when using high-volume spray-to-wet applications. Apply a 4- to 7-percent solution of this product when using low volume directed sprays for spot treatment.

TANK MIXTURES: This product may be mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments as well as for enhanced control of woody brush and trees, provided the specific product used is labeled for use on these sites. Refer to the individual product labels for approved sites and application rates:

Arsenal	atrazine ¹
dicamba ¹	diuron ¹
Escort	Escort XP
Garlon 3A	Garlon 4
Hyvar X	Hyvar X-L
Sahara DG	simazine ¹
Spike 80 DF	Vanquish
Transline	Velpar L
Velpar DF	2,4-D ¹

¹ Tank mixtures with products containing this single generic active ingredient may be made provided the specific product is labeled for application at the use site.

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 11 to 32 fluid ounces of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpetcreeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust XP. If tank-mixed, use no more than 11 to 32 fluid ounces of this product with 1 to 2 ounces of Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestem, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpetcreeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may occur.

8.11 Roadsides

All of the applications in the **General Areas and Industrial Sites** section may be made on roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails and Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

TANK MIXTURES: This product may be tank-mixed with the following products, for shoulder, guardrail, spot and bare ground treatments provided that the specific product used is labeled for use on these sites. Refer to the individual product labels for approved sites and application rates:

atrazine ¹	Oust XP
Clarity	Outrider
Crossbow L	Pendulum 3.3 EC
dicamba ¹	Pendulum WDG
diuron ¹	Plateau
Endurance	Princep DF
Escort	Princep 4L
Escort XP	Ronstar 50WP
Gallery 75 DF	Sahara
Krovar I DF	simazine ¹
Landmark II MP	Surflan
Landmark MP	Telar
Landmark XP	Vanquish
	2,4-D ¹

¹ Tank mixtures with products containing this single generic active ingredient may be made provided the specific product is labeled for application at the use site.

Release of Bermudagrass or Bahiagrass

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring green-up. This product may also be tank-mixed with Outrider, or Oust XP for residual control. Tank mixtures of this product with Oust XP may delay green-up.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 5 to 44 fluid ounces of this product in a tank mix with 0.75 to 1.33 ounces of Outrider herbicide per acre. Read and follow all label directions for Outrider herbicide.

Apply 6 to 44 fluid ounces of this product per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust XP. Apply the labeled rates in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in green-up and minimize injury, add no more than 1 ounce of Oust XP per acre on bermudagrass and no more than 0.5 ounce of Oust XP per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 11 to 32 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

- | | |
|------------------|----------------|
| Bahiagrass | Johnsongrass |
| Bluestem, silver | Trumpetcreeper |
| Fescue, tall | Vaseygrass |

This product may be tank-mixed with Outrider herbicide for control or partial control of Johnsongrass and other weeds listed on the Outrider label. Use 5 to 22 fluid ounces of this product with 0.75 to 1.33 ounces of Outrider per acre. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank-mixed with Oust XP. If tank-mixed, use no more than 11 to 22 fluid ounces of this product with 1 to 2 ounces of Oust XP per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust XP label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

- | | |
|------------------|----------------|
| Bahiagrass | Fescue, tall |
| Bluestem, silver | Johnsongrass |
| Broomsedge | Poorjoe |
| Dallisgrass | Trumpetcreeper |
| Dock, curly | Vaseygrass |
| Dogfennel | Vervain, blue |

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not recommended, since severe injury may occur.

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 1.5 to 3 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

This product may be used for control or partial control of Johnsongrass and other weeds listed on the Outrider herbicide label. Apply 4 fluid ounces of this product with 0.75 to 2.0 ounces of Outrider per acre. Use the higher rates for control of perennial weeds and annual weeds greater than 6 inches in height. Use only on well established bahiagrass.

A tank mixture of this product plus Oust XP may be used. Apply 4 fluid ounces of this product plus 0.25 ounce of Oust XP per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.12 Rangelands

This product will control or suppress many annual weeds growing in perennial cool- and warm-season grass rangelands, pastures, and industrial sites. Preventing weed seed production is critical to the successful control of annual grassy weeds invading these perennial grass sites. Follow-up applications in sequential years should eliminate most of the viable seeds. Grazing of treated areas should be delayed to

encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

Bromus: This product may be used to control or suppress downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*), cheatgrass (*Bromus secalinus*), cereal rye and jointed goatgrass found in rangelands pastures and industrial sites. Apply 5 to 11 fluid ounces of this product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead: To control or suppress medusahead, apply 11 fluid ounces of this product per acre at the 3-leaf stage when plants are actively growing. Delaying applications beyond this stage will result in reduced or unacceptable control. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses. Applications may be made in the fall or spring.

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

Spot Treatment, Wiper Application

This product may be applied in rangeland, pastures or industrial sites as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 2 quarts of this product per acre or less, the entire site or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 2 quarts of this product per acre, no more than 10 percent of the total site may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

8.13 Utility Sites

Use this product along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

Use this product in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product is also recommended for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

TANK MIXTURES: This product may be tank mixed with the following products for use in utility sites, provided that the specific product used is labeled for use on these sites. Refer to the individual product labels for approved sites and application rates.

For control of herbaceous weeds, use the lower tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher tank mixture rates.

Arsenal	atrazine ¹
Barricade 65WG	dicamba ¹

diuron ¹	Endurance
Escort	Escort XP
Garlon 3A ²	Garlon 4 ³
Krenite	Krovar I DF
	Oust XP
Outrider	pendimethalin ¹
Plateau	
Ronstar 50WP	Sahara
simazine ¹	Surflan AS
Surflan WDG	Telar DF
Transline	Vanquish
Velpar DF	Velpar L
2,4-D ¹	

¹ Tank mixtures with product containing this generic active ingredient may be made provided the specific product is labeled for application at the use site.

² Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray incompatibility problems.

³ For side trimming treatments, this product can be used alone or in a tank mixture with Garlon 4.

8.14 Grass Seed or Sod Production

Use this product in grass seed and sod production for preplant, at-planting, preemergence, removal of established stands, renovation, site preparation, shielded spraying, wiper application, spot treatment, and creating rows in annual ryegrass.

Preplant, At-Planting, Preemergence, Removal of Established Stands, Renovation, Site Preparation

This product controls most existing vegetation for purposes of renovating turf or forage grass seed areas or for establishing turfgrass grown for sod. It may also be used to destroy remaining undesired grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting, or for renovation purposes. Applications must be made prior to crop emergence in order to avoid crop injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the herbicide spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates up to 3.3 quarts per acre may be used to totally remove established stands of tough to kill grass species.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 2 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Shielded Sprayers

Apply 22 to 64 fluid ounces of this product in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aids in shielded sprayer applications. Best results are obtained when the grass seed plants are small enough to easily pass by the protective shields.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Wiper Application

This product may be applied over the top of desirable grasses using wiper applicators for the control of tall weeds.

Contact of this product in any manner with desirable vegetation may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

Spot Treatment

Apply a 1-percent solution of this product using hand-held spray equipment to control weeds within established vegetation prior to heading of grasses grown for seed. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

The grass sprayed in the treated area will be killed. Take care not to spray or allow spray to drift outside the target area in order to avoid unwanted grass injury or destruction.

Creating Rows in Annual Ryegrass

Apply 11 to 22 fluid ounces of this product per acre. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height. Use the higher rate within the labeled range when ryegrass is greater than 6 inches in height.

Set nozzle heights to allow the establishment of the desired row spacing. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

Take care not to spray or allow spray to drift outside target area in order to avoid unwanted grass destruction.

Grower assumes all responsibility for losses resulting from misapplication of this product.

8.15 Pastures

This product may be applied to any pasture grass (other than food crops in the *Gramineae* family), including bahiagrass, bermudagrass, bluegrass, brome, fescue, guineagrass, kikuyugrass, orchardgrass, pangola grass, ryegrass, timothy, and wheatgrass. Application can be made as a spot treatment, wiper application, preplant, preemergence, pasture renovation, or postemergent broadcast.

Preplant, Preemergence, Pasture Renovation

This product may be applied for weed control prior to planting or emergence of forage grasses. This product may also be applied to control perennial pasture species listed on this label prior to replanting.

If application rates total 2 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 2 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

Spot Treatment, Wiper Application

This product may be applied in pastures as a spot treatment, or over the top of desirable grasses using wiper applicators to control tall weeds. Applications may be repeated in the same area at 30-day intervals.

For spot treatments or wiper application methods using rates of 2 quarts of this product per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper applications are made using rates above 2 quarts of this product per acre, no more than 10 percent of the total pasture may be

treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting for feed.

Postemergent Weed Control (Broadcast Treatments)

This product may be applied to pastures to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 8 to 11 fluid ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. Use of higher application rates will cause stand reductions. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 2 quarts of this product per acre per year onto pasture grasses except for renovation uses as described previously in this section.

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the specified range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for rates to control annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 4 to 7 quarts per acre for enhanced results.

9.1 Annual Weeds

Use 22 fluid ounces of this product per acre if weeds are less than 6 inches in height or runner length and 1.0 to 2.7 quarts of this product per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions. Use the higher rate for tough-to-control species regardless of the weed size at application. Treat tough-to-control weeds early when they are relatively small. This product may be tank mixed provided the tank-mix product is labeled for application at the target site. Refer to the individual tank mix product labels for approved sites and application rates.

For spray-to-wet applications, apply a 0.4-percent solution of this product to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 0.7- to 1.5-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds.

For low volume directed spray applications, use a 4- to 7-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Weed Species

- Anoda, spurred
- Barley
- Barnyardgrass
- Bassia, fivehook
- Bittercress
- Black nightshade
- Bluegrass, annual
- Bluegrass, bulbous
- Brome, downy
- Brome, Japanese

Browntop panicum
Buttercup
Carolina foxtail
Carolina geranium
Castorbean
Cheatgrass
Cheeseweed (*Malva parviflora*)
Chervil
Chickweed
Cocklebur
Copperleaf, hophornbeam
Corn
Corn speedwell
Crabgrass
Dwarf dandelion
Eastern mannagrass
Eclipta
Fall panicum
Falsedandelion
Falseflax, smallseed
Fiddleneck
Field pennycress
Filaree
Fleabane, annual
Fleabane, hairy (*Conyza bonariensis*)
Fleabane, rough
Florida pusley
Foxtail
Goatgrass, jointed
Goosegrass
Grain sorghum (milo)
Groundsel, common
Hemp sesbania
Henbit
Horseweed/Marestail (*Conyza canadensis*)
Itchgrass
Johnsongrass, seedling
Junglerice
Knotweed
Kochia
Lamb's-quarters
Little barley
London rocket
Mayweed
Medusahead
Morningglory (*Ipomoea spp*)
Mustard, blue
Mustard, tansy
Mustard, tumble
Mustard, wild
Oats
Pigweed
Plains/Tickseed coreopsis
Prickly lettuce

Puncturevine
Purslane, common
Ragweed, common
Ragweed, giant
Red rice
Russian thistle
Rye
Ryegrass
Sandbur, field
Shattercane
Shepherd's-purse
Sicklepod
Signalgrass, broadleaf
Smartweed, ladythumb
Smartweed, Pennsylvania
Sowthistle, annual
Spanishneedles
Speedwell, purslane
Sprangletop
Spurge, annual
Spurge, prostrate
Spurge, spotted
Spurry, umbrella
Starthistle, yellow
Stinkgrass
Sunflower
Teaweed/Prickly sida
Texas panicum
Velvetleaf
Virginia copperleaf
Virginia pepperweed
Wheat
Wild oats
Witchgrass
Woolly cupgrass
Yellow rocket

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the specified range.

Ensure thorough coverage when using spray-to-wet treatments with hand-held equipment. For best results, use a 1.5-percent solution on harder-to-control perennials such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

For low volume directed spray applications, use a 4- to 7-percent solution of this product. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Allow 7 or more days after application before tillage.

35
49

Weed Species	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1 - 1.5	1.5
Alligatorweed*	3	1
Anise (fennel)	1.3 - 2.7	1 - 1.5
Bahiagrass	2 - 3.3	1.5
Beachgrass, European (<i>Ammophila arenaria</i>)	--	3.5
Bentgrass*	1	1.5
Bermudagrass	3.3	1.5
Bermudagrass, water (knotgrass)	1	1.5
Bindweed, field	2.7 - 3.3	1.5
Bluegrass, Kentucky	1.5	1.5
Blueweed, Texas	2.7 - 3.3	1.5
Brackenfern	2 - 3	1
Bromegrass, smooth	1.5	1.5
Bursage, woolly-leaf	--	1.5
Canarygrass, reed	1.5 - 2	1.5
Cattail	2 - 3.3	1.5
Clover; red, white	2 - 3.3	1.5
Cogongrass	2 - 3.3	1.5
Dallisgrass	2 - 3.3	1.5
Dandelion	2 - 3.3	1.5
Dock, curly	2 - 3.3	1.5
Dogbane, hemp	2.5	1.5
Fescue (except tall)	3	1.5
Fescue, tall	2	1.5
German ivy	1.3 - 2.7	1 - 1.5
Guineagrass	2	1
Horsenettle	2 - 3.3	1.5
Horseradish	3	1.5
Iceplant	1.3	1.5 - 2
Jerusalem artichoke	2 - 3.3	1.5
Johnsongrass	1.3 - 2	1
Kikuyugrass	1.5 - 2	1.5
Knapweed	3	1.5
Lantana	--	1
Lespedeza	2 - 3.3	1.5
Milkweed, common	2	1.5
Muhly, wirestem	1.5	1.5
Mullein, common	2 - 3.3	1.5
Napiergrass	2 - 3.3	1.5
Nightshade, silverleaf	1.5	1.5
Nutsedge; purple, yellow	2	1 - 1.5
Orchardgrass	1.5	1.5
Pampasgrass	2 - 3.3	1 - 1.5
Paragrass	2 - 3.3	1.5

Pepperweed, perennial	2.7	1.5
Phragmites*	2 - 3.3	1 - 1.5
Poison hemlock	1.3 - 2.7	1 - 1.5
Pokeweed, common	1	1.5
Quackgrass	1.3 - 2	1.5
Redvine*	1.5	1.5
Reed, giant	2.7 - 3.3	1.5
Ryegrass, perennial	1.5 - 2	1
Smartweed, swamp	2 - 3.3	1.5
Sowthistle, perennial	1.5 - 2	1.5
Spurge, leafy*	--	1.5
Starthistle, yellow	1.5	1.5
Sweet potato, wild*	--	1.5
Thistle, artichoke	1.3 - 2	1 - 1.5
Thistle, Canada	1.5 - 2	1.5
Timothy	1.5 - 2	1.5
Torpedograss*	2.7 - 3.3	1.5
Trumpetcreeper*	1.5 - 2	1.5
Vaseygrass	2 - 3.3	1.5
Velvetgrass	2 - 3.3	1.5
Wheatgrass, western	1.5 - 2	1.5

* Partial control

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed on this label, or in separate supplemental labeling or Fact Sheets published by Monsanto Company for this product. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

For best results when using hand-held equipment, use a 1.5-percent solution on harder-to-control woody brush and trees.

For low volume directed-spray applications, apply a 4- to 7-percent solution of this product. Spray coverage should be uniform with at least 50-percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple spouts.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Broadcast Rate (QT/A)	Hand-Held Spray-to-Wet % Solution
Alder	2 - 3	1

Ash*	1.5 - 3.3	1 - 1.5
Aspen, quaking	1.5 - 2	1
Bearclover (Bearnat)*	1.5 - 3.3	1 - 1.5
Beech*	1.5 - 3.3	1 - 1.5
Birch	1.5 - 2	1
Blackberry	2 - 3	1
Blackgum	1.5 - 3.3	1 - 1.5
Bracken	1.5 - 3.3	1 - 1.5
Broom; French, Scotch	1.3 - 3.3	1 - 1.5
Buckwheat, California*	1.3 - 2.5	1 - 1.5
Cascara*	1.5 - 3.3	1 - 1.5
Catsclaw*	--	1
Ceanothus*	1.5 - 3.3	1 - 1.5
Chamise*	1.3 - 3.3	1
Cherry; bitter, black, pin	1.5 - 2	1
Coyote brush	2 - 2.7	1 - 1.5
Deerweed	1.3 - 3.3	1
Dogwood*	1.5 - 3.3	1 - 1.5
Elderberry	1.5 - 2	1
Elm*	1.5 - 3.3	1 - 1.5
Eucalyptus	--	1.5
Gorse*	1.5 - 3.3	1 - 1.5
Hasardia*	1.3 - 2.5	1 - 1.5
Hawthorn	1.5 - 2	1
Hazel	1.5 - 2	1
Hickory*	1.5 - 3.3	1 - 1.5
Honeysuckle	2 - 3	1
Hornbeam, American*	1.5 - 3.3	1 - 1.5
Kudzu	2.5 - 3	1.5
Locust, black*	1.5 - 2.5	1 - 1.5
Madrone resprouts*	--	1.5
Manzanita*	1.5 - 3.3	1 - 1.5
Maple, red	1.5 - 3	1
Maple, sugar	--	1
Monkey flower*	1.3 - 2.7	1 - 1.5
Oak; black, white*	1.5 - 3	1 - 1.5
Oak, post	2 - 3	1
Oak; northern, pin	1.3 - 2.7	1
Oak, Scrub*	1.3 - 2.7	1
Oak; southern red	1.5 - 2	1
Peppertree, Brazilian (Florida holly)*	1.3 - 3.3	1 - 1.5
Persimmon*	1.5 - 3.3	1 - 1.5
Pine	1.5 - 3.3	1 - 1.5
Poison ivy	2.5 - 3.3	1.5
Poison oak	2.5 - 3.3	1.5
Poplar, yellow*	1.5 - 3.3	1 - 1.5

Redbud, eastern	1.5 - 3.3	1 - 1.5
Rose, multiflora	1.5	1
Russian olive*	1.5 - 3.3	1 - 1.5
Sage, black	1.3 - 2.7	1
Sage, white*	1.5 - 2.7	1 - 1.5
Sage brush, California	1.3 - 2.7	1
Salmonberry	1.5 - 2	1
Saltcedar*	1.5 - 3.3	1 - 1.5
Sassafras*	1.5 - 3.3	1 - 1.5
Sourwood*	1.5 - 3.3	1 - 1.5
Sumac; laurel, poison, smooth, sugarbush, winged*	1.5 - 3	1 - 1.5
Sweetgum	1.5 - 2	1
Swordfern*	1.5 - 3.3	1 - 1.5
Tallowtree, Chinese	--	1
Tan oak resprouts*	--	1.5
Thimbleberry	1.5	1
Tobacco, tree*	1.5 - 2.5	1 - 1.5
Toyon*	--	1.5
Trumpetcreeper	1.5 - 2	1
Vine maple*	1.5 - 3.3	1 - 1.5
Virginia creeper	1.5 - 3.3	1 - 1.5
Waxmyrtle, southern*	1.5 - 3.3	1 - 1.5
Willow	2 - 3	1
Yerbasenta*	--	1.5

* Partial control

10.0 LIMIT OF WARRANTY AND LIABILITY

Monsanto Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

To the extent consistent with applicable law, buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the fullest extent permitted by law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company to the extent consistent with applicable law, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

39
/ 49

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

[INSERT BRAND NAME], Outrider, Monsanto and Vine symbol are trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

This product is protected by U.S. Patent No's. 5,668,085 and 6,365,441. Other patents pending. No license granted under any non-U.S. patent(s).

EPA Reg. No. 524-579

In case of an emergency involving this product, or for medical assistance,
Call Collect, day or night, (314) 694-4000.

© [DATE]
Packed For:
MONSANTO COMPANY
800 N. Lindbergh Blvd.
ST. LOUIS, MISSOURI, 63167 USA

APPENDIX 1 – Consolidated List of Label Claims

Rainfast in 15 Minutes
Rainfast in 30 Minutes
Rainfast in 45 Minutes
Rainfast in 60 Minutes
Rainproof in 15 Minutes
Rainproof in 30 Minutes
Rainproof in 45 Minutes
Rainproof in 60 Minutes
15-Minute Rainfastness
30-Minute Rainfastness
45-Minute Rainfastness
60-Minute Rainfastness
First New Formula in over 30 years
First New Formula in 33 years
Treats 50% more area
50% more concentrated
Brand Advantage
Performance Guarantee
Performance Guaranteed
Guaranteed Performance
Faster Symptoms
Faster physical symptoms
Faster results
Faster control of weeds
3 Times Faster
4 Times Faster
5 Times Faster
3 Times More to the Root
4 Times More to the Root
5 Times More to the Root
See Results 3x Faster
See Results 4x Faster
See Results 5x Faster
Works 3x Faster
Works 4x Faster

41
49

Works 5x Faster
More Consistent performance
NEW

42
/ 49

IV. SUPPLEMENTAL LABELING FOR INDUSTRIAL, TURF, & ORNAMENTAL USES

Table of Contents: Industrial, Turf, and Ornamental Supplemental labeling

	Name	1 st Approval Date
	TEMPLATE FOR SUPPLEMENTAL LABELS AND STANDARD LANGUAGE	20 June 2007
A	FOR USE FOR SELECTIVE WEED CONTROL ON [INSERT BRAND NAME] TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.	20 June 2007
B	AERIAL APPLICATIONS IN CALIFORNIA	20 June 2007
C	TANK MIXES FOR IMPROVED CONTROL OF BENTGRASS (Agrostis spp.)	20 June 2007
D	TANK MIXTURES FOR TOUGH TO CONTROL WEEDS	11/1/07
E	ALTERNATE INGREDIENT STATEMENT	11/1/07

43
/ 49

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using [INSERT BRAND NAME] as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

[INSERT BRAND NAME]

Herbicide

EPA Reg. No. 524-579

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

[INSERT BRAND NAME] is a registered trademark of Monsanto Technology LLC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for [INSERT BRAND NAME] for essential product performance information.

[INSERT SPECIFIC DIRECTIONS FOR USE FROM FOLLOWING SECTIONS HERE]

Read the "Limit of Warranty and Liability" in the label booklet for [INSERT BRAND NAME] before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [DATE] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

A. LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY, INCLUDING FRESNO COUNTY, CALIFORNIA

DIRECTIONS FOR USE

All labeled treatments may be made by aerial equipment where appropriate, provided that the applicator complies with the precautions and restrictions specified on this supplemental labeling and in the product label booklet. Refer to Aerial Equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the product label for additional information. Refer to the individual use site section of the product label, or to other supplemental labeling or technical fact sheets published separately for this product by Monsanto, for specific use instructions.

AVOID DRIFT—DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

When applied as directed, under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

When tank-mixing this product with 2,4-D, only 2,4-D amine formulations may be used for aerial application in California. Tank mixtures with 2,4-D amine formulations may be applied by air in California for fallow and reduced tillage systems, and for alfalfa and pasture renovation applications only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

ADDITIONAL INFORMATION FOR FRESNO COUNTY, CALIFORNIA

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

- North: Fresno County line
- South: Fresno County line
- East: State Highway 99
- West: Fresno County line

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Recommendations

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Applications at Night—Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

To report known or suspected misuse of this product, call 1-800-332-3111.

For additional information on the proper aerial application of this product, call 916-784-1718.

B. FOR USE FOR SELECTIVE WEED CONTROL ON [INSERT BRAND NAME] TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS

DIRECTIONS FOR USE

[INSERT BRAND NAME] Tolerant Tall Fescue Selections For Seed Production

Use this product on **[INSERT BRAND NAME]** tolerant tall and fine fescue grown for seed production only.

This product may be applied at rates of 2.5 to 9 fluid ounces per acre as a postemergence spray on **[INSERT BRAND NAME]** tolerant tall fescue selections. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage of weeds.

When applied postemergence, this product will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, pennycress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the **[INSERT BRAND NAME]** label booklet for a complete list of weeds controlled or suppressed.

NOTE: The recommended rate for this use will limit the level of control of certain species of weeds.

NOTE: Some crop discoloration and yellowing may occur at higher rates of application with **INSERT BRAND NAME]** tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

Timing Of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy and in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

NOTE: Only two applications per crop growth cycle may be made to any one site. If two applications are required, only one Fall and one Spring application may be made during one 12 month cycle.

C. TANK MIXTURES FOR IMPROVED CONTROL OF BENTGRASS (*Agrostis spp.*)

DIRECTIONS FOR USE

[INSERT BRAND NAME] herbicide may be tank mixed with the products listed below for improved control of bentgrass (*Agrostis spp.*). When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of each product used. Refer to each individual product label for approved use sites.

ENVOY®	FUSILADE® II
FUSION®	VANTAGE®

Dry ammonium sulfate, at 1 to 2 percent by weight, may be added to the spray solution for the above applications. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

Broadcast Treatment

- Apply 1.6 to 2.2 quarts per acre of this product plus 34 fluid ounces per acre of **Envoy** in 20 to 40 gallons per acre of spray solution.
- Apply 1.6 to 2.2 quarts per acre of this product plus 24 fluid ounces per acre of **Fusilade II** in 20 to 40 gallons per acre of spray solution.
- Apply 1.6 to 2.2 quarts per acre of this product plus 60 fluid ounces per acre of **Vantage** in 20 to 40 gallons per acre of spray solution.
- Apply 1.6 to 2.2 quarts per acre of this product plus 9 fluid ounces per acre of **Fusion** in 20 to 40 gallons per acre of spray solution.

Re-treatment may be needed in the event of incomplete control.

Spot Treatment

- Mix 1.5 fluid ounces of this product with 1.3 fluid ounces of **Envoy** in one gallon of water and spray-to-wet.
- Mix 1.5 fluid ounces of this product with 0.75 fluid ounces of **Fusilade II** in one gallon of water and spray-to-wet.
- Mix 1.5 fluid ounces of this product with 3 fluid ounces of **Vantage** in one gallon of water and spray-to-wet.
- Mix 1.5 fluid ounces of this product with 0.25 fluid ounces of **Fusion** in one gallon of water and spray-to-wet.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

D. TANK MIXTURES FOR TOUGH TO CONTROL WEEDS

[INSERT INTO SUPPLEMENTAL LABEL TEMPLATE AS INDICATED]

DIRECTIONS FOR USE

Use in areas such as airports, apartment complexes, Christmas tree farms, commercial sites, ditch banks, dry ditches, dry canals, fencerows, golf courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, recreational areas, residential areas, rights-of-way, roadsides, schools, sod or turf seed farms, sports complexes, storage areas, substations, turfgrass areas, utility sites, warehouse areas, and wildlife management areas.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for [INSERT BRAND NAME] for essential product performance information.

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "WEEDS CONTROLLED" section of the [INSERT BRAND NAME] label booklet for specific rates. For tough to control species, where dense stands occur, or where conditions for control are not ideal, 4 to 7 quarts per acre of this product can be used for improved results.

TANK MIXTURES

This product provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of the weeds listed in the individual product labels.

The following list of products may be tank mixed with this product, provided that the specific product is registered for application to the target site. Any labeled rate of this product may be used in a tank mixture with these products. Refer to these product labels for approved sites and application rates.

- | | |
|-----------------------|----------------------------|
| 2,4-D ¹ | Krovar I DF |
| Arsenal | Landmark II MP |
| atrazine ¹ | Landmark MP |
| Barricade 65WG | Landmark XP |
| Crossbow L | Milestone |
| dicamba ¹ | |
| diuron ¹ | Oust XP |
| Endurance | Outrider |
| Escort | Overdrive |
| Escort XP | pendimethalin ¹ |
| Gallery 75 DF | Plateau |
| Garlon 3A | Plateau DG |
| Garlon 4 | Poast |
| Goal 2XL | Quicksilver |
| Hyvar X | Ronstar 50 WSP |
| Hyvar X-L | Sahara DG |
| Krenite | simazine ¹ |

Spike 80DF
Surflan AS
Surflan WDG
Telar DF

Transline
Velpar DF
Velpar L

¹ Tank mixtures with products containing this generic active ingredient may be made provided the specific product is registered for this use.

Refer to the individual product labels for specific sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mix products are well mixed with the spray solution before adding this product.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

When used in combination as directed by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Monsanto product in such combination use.

Outrider is a trademark of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

E. ALTERNATE INGREDIENT STATEMENT

[INSERT INTO SUPPLEMENTAL LABEL TEMPLATE AS INDICATED]

INGREDIENTS:

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its potassium salt..... 48.7%

OTHER INGREDIENTS (including 8.8% surfactant):.....51.3%

100.0%

*Contains 660 grams per liter or 5.5 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its potassium salt. Equivalent to 540 grams per liter or 4.5 pounds per U.S. gallon of the acid, glyphosate.

Product Description: This product is a postemergence, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing 8.8 percent surfactant and no additional surfactant is needed or recommended.

LIST OF CHANGES THIS SUBMISSION

- Update Storage and Disposal Section for compliance with the Container Containment Rule and PR Notice 2007-4.
- Incorporate hollow stem injection supplemental label approved November 1, 2007
- General updates for clarification and consistency with previously approved glyphosate product labels
- Deleted references to Oust to only reference Oust XP as Oust is no longer marketed

524-579

7/1/2009

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
TOXIC SUBSTANCES

AND

JUL 1 - 2009

Ms. Dawn Fee-White
Monsanto Company,
1300 I (Eye) Street, N.W., Suite 450 East
Washington, DC 2005

Dear Ms. Fee-White:

Subject: Roundup Pro Max Herbicide (Foliar and Broadcast Treatment of Japanese Knotweed and Oriental Bittersweet)
EPA Registration No. 524-579
Application Dated April 2, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable. A stamped copy of labeling is enclosed for your records.

Sincerely,

James A. Tompkins for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **ROUNDUP PROMAX™ HERBICIDE** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PROMAX Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

ROUNDUP PROMAX™ HERBICIDE LOGO

For Foliar and Broadcast Treatment of Japanese Knotweed & Oriental Bittersweet


EPA Reg. No. 524-579

Roundup PROMAX is a trademark of Monsanto Technology LLC.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION", "MIXING AND APPLICATION INSTRUCTIONS" and "AQUATIC AND OTHER NON-CROP SITES" sections of the label booklet for Roundup PROMAX herbicide for essential product performance information. See the "WEEDS CONTROLLED" section of the label booklet for Roundup PROMAX herbicide for optimum stage of treatment of weed for best results.

JAPANESE KNOTWEED

For control of Japanese knotweed (*Polygonum cuspidatum*), this product may be applied as a 2.0% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

For broadcast applications, apply 3.25 quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

ORIENTAL BITTERSWEET

For control of Oriental bittersweet (*Celastrus orbiculatus*), this product may be applied as a 1.5% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

For broadcast applications, apply 2 quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

WARRANTY AND LIABILITY

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup PROMAX herbicide before buying or using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© 2009 MONSANTO COMPANY
800 N. Lindbergh Blvd.
St. Louis, Missouri, 63167 U.S.A.

[Print Plate] [Approval date]