524-579

08-17-2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

8-17:09

Dawn Fee-White Monsanto Company 1300 I (Eye) Street, NW Suite 450 East Washington, DC 20005

Subject: EPA Reg. 524-579 / Supplemental Label

Dear Ms. Fee-White:

The Supplemental Label is acceptable. The supplemental label must be incorporated to the Master Label within 18 months from the date of this letter

If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

Jim Tompkins

Jim Tompkins Product Manager 25 Herbicide Branch Registration Division (7505P)

READ THE ENTIRE LABEL FOR ROUNDUP PROMAX™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PROMAX Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

ROUNDUP PROMAX™ HERBICIDE LOGO

For Foliar and Broadcast Treatment of Japanese Knotweed & Oriental Bittersweet

EPA Reg. No. 524-579

.

Roundup PROMAX is a trademark of Monsanto Technology LLC.

()

Keep out of reach of children.

CAUTION!

In EPA Letter Dated: 3 - 1 - 0 - 0 - 0Under the Federal Insecticide Fungicide, and Rodenticide A. as amended, for the pesticide

registered under EPA Rep 344

د ٥٥ درتهد

00066

with COMMENTS

524-579

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION", <u>and</u> "MIXING AND APPLICATION INSTRUCTIONS" and "AQUATIC AND OTHER NON-CROP SITES" sections of the label booklet for Roundup PROMAX herbicide for essential product performance information. See the "WEEDS CONTROLLED" section of the label booklet for Roundup PROMAX herbicide for optimum stage of treatment of weed for best results.

JAPANESE KNOTWEED

For control of Japanese knotweed (*Polygonum cuspidatum*), this product may be applied as a 2.0% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

deleted AQUATIC AND OTHER NON-CROP SITES as not applicable to this product. comested rate error on knotweed

For broadcast applications, apply 3.25<u>2.75</u> quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

ORIENTAL BITTERSWEET

For control of Oriental bittersweet (*Celastrus orbiculatus*), this product may be applied as a 1.5% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

For broadcast applications, apply 2 quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

WARRANTY AND LIABILITY

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup PROMAX herbicide before buying or using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© 2009 MONSANTO COMPANY 800 N. Lindbergh Blvd. St. Louis, Missouri, 63167 U.S.A.

[Print Plate]