

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

JAN -4 2012

Daniel J. Jenkins Monsanto Company 1300 I Street, NW Suite 450 East Washington, DC 20005

Subject:

Supplemental Label: Pre-plant, At-Planting, and Pre-emergence Applications in

Soybeans and Cotton Warrant Herbicide EPA Reg. No. 524-591

Date Submitted: November 9, 2011

Dear Mr. Jenkins:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable.

At your next label printing, or within eighteen (18) months of the date of this letter, whichever comes first, you must incorporate this supplemental labeling into the main product labeling. A stamped copy of your label is enclosed for your records. If you have any questions, please contact Maggie Rudick at (703) 347-0257 or rudick.maggie@epa.gov.

Sincerely,

Kable Bo Davis Product Manager 25 Herbicide Branch

Registration Division (7505P)

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL AFFIXED TO THE CONTAINER FOR WARRANT HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. Use of [Insert Brand Name] according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container and in this supplemental labeling for [Insert Brand Name].

It is a violation of federal law to use this product in any manner inconsistent with its labeling. The labeling must be in the possession of the user at the time of herbicide application.

This supplemental label expires on January 4, 2015 and must not be used or distributed after this date.

Primary Brand Name: MON63410 Herbicide Alternate Brand Name: WARRANT™ Herbicide JAN -4 2012

Under the Federal Insections.

Functions, and Hoderatology designed to the positions of the positions.

DPA Neg. No.

EPA Reg. No. 524-591

SHAKE WELL BEFORE USING

Preplant, At-Planting, and Preemergence Applications in Soybeans and Cotton

Keep Out of Reach of Children.

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

See the "Product Information" and "Mixing, Spraying, and Handling Instructions" sections of the main label booklet for [Insert Brand Name] for essential product information.

Read the "Limit of Warranty and Liability" in the main label booklet for [Insert Brand Name] before using. These terms apply to this supplemental label and if these terms are not acceptable, return the product unopened at once.

Approved Application System Ground: Broadcast boom

When applied preplant, at-planting, or preemergence in soybeans or cotton this product will provide preemergence control or reduced competition of the annual weeds listed in the "WEEDS CONTROLLED" section of the label booklet for [Insert Brand Name]. If weeds are emerged at the time of application, apply a labeled postemergence herbicide to control the emerged weeds. Use of a residual herbicide for the control of weeds not listed on the [Insert Brand Name] label is recommended. Observe all directions for use, precautions, and restrictions on the labeling of the postemergence herbicide or residual herbicide.

SOYBEANS

Approved Application Method
Preplant, At-planting or Preemergence Surface

[Insert Brand Name] may be applied preplant, at-planting or preemergence to **soybeans** at 1.25 to 2 quarts per acre (qts/A) according to the rate table below. The optimum rate of application is 1.5 qts/A. Apply broadcast to the soil surface according to the rate table listed below. Mechanical incorporation is not recommended. This product will not control emerged weeds.

Application of this product with other postemergence or soil applied herbicides may increase the potential for crop injury.

Application of this product followed by conditions that do not favor adequate crop growth or which cause stress (cold, wet soils), or under waterlogged conditions from excessive irrigation or rainfall, may result in crop injury.

DO NOT exceed 4 quarts per acre per season of [Insert Brand Name] when making a second application, including a postemergence application to soybeans. Read and follow the use instructions for postemergence applications to soybeans in the main label booklet for [Insert Brand Name].

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE*							
	Less than 1.5% Organic Matter (quarts)			1.5% or More Organic Matter (quarts)				
Coarse	1.25	to	1.6	1.25	to	1.7		
Medium	1.25	to	1.7	1.25	to	1.9		
Fine	1.25	to	1.9	1.25	to	2.0		

^{*} Use the higher rate in the range for areas of heavy weed infestation.

COTTON

Approved Application Method

Preplant, At-planting or Preemergence Surface

[Insert Brand Name] may be applied preplant, at-planting or preemergence to **cotton** at 1.25 to 2 quarts per acre (qts/A) according to the rate table below. The optimum rate of application is 1.5 qts/A. Apply broadcast to the soil surface according to the rate table listed below. Mechanical incorporation is not recommended. This product will not control emerged weeds.

Application of this product with other postemergence or soil applied herbicides may increase the potential for crop injury.

Application of this product followed by conditions that do not favor adequate crop growth or which cause stress (cold, wet soils), or under waterlogged conditions from excessive irrigation or rainfall, may result in crop injury.

DO NOT exceed 4 quarts per acre per season of [Insert Brand Name] when making a second application, including a postemergence application to cotton. Read and follow the use instructions for postemergence applications to cotton in the main label booklet for [Insert Brand Name].

SOIL TEXTURAL GROUP Coarse	BROADCAST RATE PER ACRE*								
	Less that	n 1.5% Or (quarts	ganic Matter)	1.5% or More Organic Matter (quarts)					
	1.25	to	1.6	1.25	to	1.7			
Medium	1.25	to	1.7	1.25	to	1.9			
Fine	1.25	to	1.9	1.25	to	2.0			

^{*} Use the higher rate in the range for areas of heavy weed infestation.

© 2012 MONSANTO COMPANY

[™] WARRANT is a trademark of Monsanto Technology LLC.