

ACCEPTED

527-40

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROUPELANTICIDE ACT
FOR ECONOMIC POISON REGISTRI
ED UNDER NO. 527-40. SUBJECT
TO ATTACHED COMMENTS.

WARE-O-CIDE SUPER STRENGTH

U.S.D.A. Reg. No. 527-40

Ware-O-Cide Super Strength is a highly effective residual insecticide formulated specifically for use in Furniture Warehouses as an aid in the control of accessible stages of clothes moths, silverfish, carpet beetles, roaches, flies, bedbugs and certain other insects.

Active Ingredients 100%

Refined Mineral Oil	95.93%
Technical Chlordane*	3.00%
Piperonyl Butoxide, Technical**	0.42%
N-Octyl Bicycloheptene Dicarboximide	0.25%
Pyrethrins	0.25%
Methyl Salicylate	0.15%

*Equivalent to 1.20% octachloro-4,7-methanotetrahydroindane and 1.20% related compounds.

**Equivalent to 0.34% (butylcarbityl) (6-propylpiperonyl) ether and 0.08% related compounds.

DIRECTIONS

Thoroughly clean area by removing accumulations of dust and lint from cracks, crevices and around stored articles. After wrapping or covering furniture, paint (fine spray or coarse spray) directly into cracks, crevices and all places where insects may propagate, and on walls and other surfaces where insects congregate. Spray (coarse spray) at frequent intervals and when necessary in crevices where all over-stuffed furniture, rugs and other textiles are properly wrapped or covered.

Do not apply in any manner not specified on the label. Not for household use.

CAUTION: Avoid contact with skin or eyes. If brought into repeated or prolonged contact with skin this solution can cause toxic symptoms. Wash at once with soap and water. Harmful if swallowed. Avoid inhalation of spray—use of respirator or mask is advised. Do not use on household pets or other animals. Avoid contamination of feed or foodstuffs. Do not smoke while spraying. Combustible mixture—do not spray into or near an open flame.

KEEP OUT OF REACH OF CHILDREN

CONTENTS

GALLONS

fungicide

SUGGESTED SPREADER SETTINGS

to ensure proper distribution calibrate spreader before application

60 Lbs (27.2 Kg) treats 22,000 sq ft (1/2 Acre/2045 sq m) at NORMAL RATE

60 Lbs (27.2 Kg) treats 11,000 sq ft (1/4 Acre/1022 sq m) at DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF SPREADER	SPREADER SETTINGS	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type) Scotts Rotaries	3 mph	Overlap wheels	6	8
(R-7, PR-1, PT-10)	3 mph	8 feet	H	L
Lely—WTR Model	4.5 mph	18 feet	5H	6 1/2H
Lely—HR (PTO Model)	450 rpm @ 4.5 mph	18 feet	5H	6 1/2H
Gandy 8 ft A Series	4.5 mph	Overlap wheels	19	25
42" Model	2.5 mph	Overlap wheels	38	46

FOR USE ON BENTGRASS PUTTING GREENS AND OTHER TURFGRASS AREAS

Apply to either moist or dry foliage.

DOLLARSPOT* (*Sclerotinia homoeocarpa*) and BROWNPATCH (*Rhizoctonia solani*)—to prevent fungus activity treat in late spring at **NORMAL RATE**, repeat at 2 or 3 week intervals.

During prolonged periods of high temperature and humidity, the above intervals may have to be shortened for brownpatch prevention.

To control unexpected attacks of either fungus, treat immediately when damage is first noticed. Use **DOUBLE RATE**. Follow with preventative program outlined above.

*If treatment is not effective following use of this product as recommended, a tolerant strain of fungi may be present. Contact your ProTurf Tech Rep, consideration should be given to prompt use of other suitable ProTurf fungicides.

FUSARIUM PATCH (*Fusarium nivale*)—to control fungus activity, apply at **DOUBLE RATE** at first sign of disease. Repeat at this rate if needed.

ProTurf Division, O M Scott & Sons, Marysville, Ohio 43040



**STORE THIS PRODUCT
IN A COOL PLACE**
HOW TO STORE OPENED BAGS



hor-
lical

ring

ff is
pply

oc