

# MILL-O-CIDE. SUPER STRENGTH

**ACCEPTED**  
JUN 01 1979  
Under the Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 527-81

CONTACT INSECTICIDE  
FOR FOOD ESTABLISHMENTS  
EPA Reg. No. 527-81-AA

Mill-O-Cide Super Strength is a high kill contact insecticide for wholesale grocers, supermarkets, bakeries, dairies, bottling plants, packing plants, breweries, mills and other establishments where food is processed or stored. When used as directed, this product will kill accessible weevils, flour moths, silverfish, roaches, flies, gnats and certain other insects on contact.

Active Ingredients 100%	
Refined Mineral Oil .....	98.50%
Piperonyl Butoxide, Technical* .....	0.80%
Pyrethrins .....	0.30%
N-Octyl Bicycloheptene Dicarboximide .....	0.20%
Methyl Salicylate .....	0.20%

\*Equivalent to 0.64% (butylcarbityl) (6-propylpiperonyl) ether and 0.16% of related compounds.

### DIRECTIONS

Area to be sprayed should be thoroughly cleaned by removing accumulations from all cracks, shelves, beams, and on and around packages of stored food products. Close doors and windows and keep closed for 10 to 15 minutes after spraying. Do not remain in treated areas. Ventilate the area after treatment is completed. Sweep up and dispose of fallen insects.

**Where To Spray:** Spray Mill-O-Cide Super Strength into the air in the form of a fine mist or fog. Spray under and around all equipment and racks, into all cracks, crevices and corners, and on floors, walls and ceilings.

**When To Spray:** Mill-O-Cide Super Strength should be sprayed once or twice a week as a regular procedure, and daily in cases of severe infestations.

**Spraying Equipment:** For best coverage and penetration use a Midland Mill-O-Jet or Tri-Jet Electric Sprayer or Midland Thermo-Fogger.

### CAUTION: KEEP OUT OF REACH OF CHILDREN.

**HARMFUL IF SWALLOWED:** Avoid breathing of spray mist. Mill-O-Cide Super Strength is a combustible mixture and should not be sprayed into or near an open flame. All pilot lights should be off while spraying. Foods should be removed & covered during treatment. All food processing surfaces should be covered during treatment, or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation. Do not reuse empty containers. Dispose of by perforating, crushing, or burying.

EPA EST. 1588-NY-1      **NET CONTENTS**      **GALLONS**      EPA EST. 1588-IL-1

FORM 1102-B

**Midland LABORATORIES, INC.**            SINCE 1903

P.O. BOX 5196      •      DES MOINES, IOWA 50306