

MILL-O-CIDE® "100"

CONTACT INSECTICIDE
FOR FOOD ESTABLISHMENTS

EPA Reg. No. 527-92

Mill-O-Cide "100" is a high kill contact insecticide for use by wholesale grocers, supermarkets, mills, dairies, bottling plants, packing plants, breweries, bakeries and other establishments where food is processed or stored. When used as directed, this product will kill roaches, flies, silverfish, gnats, confused flour beetles, cadelles, accessible granary and rice weevils, and Mediterranean flour moths.

Active Ingredients 100%

Refined Mineral Oil	99.28%
Piperonyl Butoxide, Technical*	0.40%
N-Octyl Bicycloheptene Dicarboximide	0.20%
Pyrethrins	0.12%

*Equivalent to 0.32% (butylcarbityl) (6-propylpiperonyl) ether and 0.08% of related compounds.

D I R E C T I O N S

Area to be sprayed should be thoroughly cleaned by removing accumulations from all cracks, beams, shelves and on and around packages and stored food products. Close doors and windows of area to be sprayed and keep closed for 10 to 15 minutes after spraying. Sweep up and dispose of fallen insects.

Where To Spray—Spray Mill-O-Cide "100" into the air in the form of a fine mist or fog. Spray under and around all equipment and racks, into all cracks, crevices and corners, and on walls, ceilings and floors.

When To Spray—Mill-O-Cide "100" should be sprayed once a week as a regular procedure. During the summer months spraying should be conducted at least twice weekly. In cases of severe infestation daily spraying is recommended.

Spraying Equipment—For best coverage and penetration use a Midland Tri-Jet Electric Sprayer or Midland Thermo-Fogger.

Coverage: For flying insects use 2 oz. per 1,000 cubic feet. For crawling insects wet surface or 1 gallon per 1,000 square feet.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF SWALLOWED. Mill-O-Cide Super Strength "100" is a combustible mixture and should not be sprayed into or near an open flame. All pilot lights should be off while spraying. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment, or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation. Do not reuse empty containers. Dispose of by perforating, crushing, or burying.

NET CONTENTS GALLONS

Midland LABORATORIES INC
DUBUQUE IOWA 52001



SINCE
1903

292

ACCEPTED
527-92
SEP ~~28~~³⁰ 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROUENICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
ED UNDER NO. [REDACTED] SUBJECT
TO ATTACHED COMMENTS.

RECEIVED
4 SEP 1974
REGISTRATION DIVISION
PESTICIDES, EPA