


NET CONTENTS 1 GALLON

BAYPAX

RESIDUAL INSECTICIDE

Contains BAYGON®

For Industrial and Institutional Use



BAYPAX RESIDUAL INSECTICIDE is a highly effective ready-to-use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and long residual activity.

ACTIVE INGREDIENTS:

o-Isopropoxyphenyl methylcarbamate*	1.00%
Petroleum Distillates	83.75%

INERT INGREDIENTS:	15.25%
	100.00%

* U.S. Patent No 3,111,539

BAYGON® is the registered trademark of Farbenfabriken Bayer A.G., Chemagro Corporation Licensee.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water.

Do not apply where runoff is likely to occur.
Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

DO NOT REUSE EMPTY CONTAINER. DESTROY BY PERFORATING OR CRUSHING, AND BURYING IN A SAFE PLACE.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

CAUTION: Keep out of reach of children.

See other precautions on the back panel.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Millipedes, Mosquitoes, Punkies, Sandflies, Silverfish, Sowbugs and Spiders

Exposed stages of Cadelle, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Sawtoothed Grain Beetles

Manufactured by

BIOCERTA CORPORATION
NEW YORK, N.Y. 10016

DIRECTIONS

GENERAL. Apply with a [redacted] sprayer that can deliver a pin stream of insecticide.

Spray surfaces only until wet. Avoid excessive runoff. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

SILVERFISH, FIREBRATS [redacted]—Apply to localized areas infested by these insects.

BROWN DOG TICKS—Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, [redacted] AND SPIDERS—Thoroughly spray the interior and exterior perimeter of infested premises. For CLOVER MITES, give attention around window frames.

FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES AND SANDFLIES (outdoors only)—Spray infested areas thoroughly, including outside surfaces of screens, doors, window frames, foundations, patios, and other areas where insects may enter.

APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS (COCKROACHES, ANTS, SILVERFISH, FIREBRATS, AND TO AID IN THE CONTROL OF CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTH GRAIN BEETLES)

Limited to Crack and Crevice Treatment Only—Apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Applications of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

COCKROACHES—Thoroughly treat the infested area. Spray baseboard, cracks and crevices, around window and door frames, under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

ANTS—Apply to trails and places where ants enter premises, as around window and door frames. Spray hills and runways outdoors.

CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES (Exposed stages)—Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

SILVERFISH, FIREBRATS [redacted]—Apply to localized areas infested by these insects.

RESTRICTIONS: Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

CAUTION: MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. AVOID BREATHING SPRAY MIST. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WARM WATER. WASH HANDS, ARMS, AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE RE-USE. WASH AFTER HANDLING.

IF ILLNESS OCCURS, GET PROMPT MEDICAL AID

TO PHYSICIAN: Atropine sulfate is antidotal

