

Reg # 538-9

PM-23

1/92

Scotts  
Lawn Weed Control

Treats 5,000 sq ft

46805  
1

SCOTT'S POLICY  
No-Quibble  
Guarantee  
SEE BACK FOR DETAILS

Scotts

# Lawn Weed Control

Controls dandelions,  
buckhorn, chickweed,  
knotweed, plantain,  
white clover and  
other weeds.

(see complete list on the back panel)

Not for use on St Augustinegrass, carpetgrass or  
dichondra lawns.

In accordance with EPA Notice 02-2.

Based on Scott's Labeling: 3/18/92

NET WEIGHT 5 LBS (2.27 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling.

Environmental Hazards: Do not contaminate water used for irrigation or domestic purposes. Avoid application on or near garden plants and flowers. Keep away from sensitive areas of shrub and the branch spread of trees. Draft or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bays, marshes and ponds). Do not contaminate water when cleaning of equipment or outdoors.

STORAGE: Store in a clean dry place. Recycle opened bag by folding top down and air drying.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENTS

2,4-Dichlorophenoxyacetic acid 2.95%

Dicamba (2-Methyl-4

chlorophenoxyacetic acid) 0.25%

INERT INGREDIENTS 66.80%

Total 100.00%

Product of USA

EPA Reg. No. 538-9

The O.M. Scott & Son Company

Marion, Ohio 43040

EPA # 538-091

Scotts

Lawn Weed Control

No-Quibble  
Guarantee

Treats 5,000 sq ft  
Lawn Weed Control

Scotts

46805  
1

46805  
1

Easy Open



Treats 5,000 sq ft

# Scotts Lawn Weed Control

## Controls dandelions and these lawn weeds

black medic	chickweed	knitweed	plantain	white clover
aster	cutwells	blaze	marigold	scarlet pimpernel
bindweed	chicory	dog foot	mayweed	salmon
blueberry	cleome	ground ivy	mint	shepherd
burdock	crucifer	groundsel	poppy	shepherd
catnip	cutweed	hairy	poppy	shepherd
chickweed	cutweed	hairy	poppy	shepherd
bull thistle	cutweed	hairy	poppy	shepherd
buttercup	cutweed	hairy	poppy	shepherd
buttercup	cutweed	hairy	poppy	shepherd
carpetweed	cutweed	hairy	poppy	shepherd

### Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- DO use when weeds are up and growing actively. On bare lawns only, wait until after the second mowing of the year but not before mid-March. In the fall do not treat after the first of November.
- DO apply only when lawn is wet from dew, rain or thorough watering (very important so that the Scotts Lawn Weed Control particles will stick to the weed leaves instead of falling to the soil).
- DO withhold watering for at least 24 hours after application (to avoid washing particles off the weed leaves).
- DO apply on a calm day.
- DO apply at Scotts Spreader (Standard, Automatic or Precision Flow Control Model) setting **5** (treats 5,000 sq ft)
- on bermgrass and bermudagrass lawns use setting **3 1/2** (treats 10,000 sq ft)
- DO slightly overlap spreader's wheel tracks to avoid "stripes" of untreated lawn.
- DO NOT use when rain is expected within 24 hours.
- DO NOT use when grass is tall it touches bottom of spreader (mow lawn first in such situations).
- DO NOT use on a newly seeded, sodded or sprigged lawn until the new grass has been mowed at least four times. If seeding in late spring, delay weed control application until early fall. If seeding in the late fall, delay weed control application until spring.
- DO NOT plant grass seed right after using Lawn Weed Control. Wait at least two weeks, and after a heavy watering or rain.
- DO NOT use on St. Augustinegrass, carpetgrass, dichondra lawns or lippia; nor on flowers, shrubs, fruit, vegetable plants.
- DO NOT apply on the rootzone areas of shrubs or under the branch spread of trees.
- DO NOT apply with a rotary-type spreader.

### What to expect

This product does not work overnight. It is designed to provide slow, steady controlled action. Stems and leaves of weeds may begin to twist and curl at a few days, but a month or more may be required for complete control, especially when weather is dry or cool. It is not necessary to rake out dead weeds. They will eventually disintegrate and disappear.

Only weeds growing at time of application will be eliminated. This product does not prevent weed seeds from producing new plants, whether such seeds are on the surface or buried in the soil.


If particles become washed off weed foliage within 24 hours after application, control may be less effective. If there is no indication that it is working after 2 or 3 weeks of good growing weather another treatment may be needed.

A few weeds may be resistant to control. If this happens, please forward specimens so we can identify them and advise you further.

### Tips on spreading with Scotts spreaders

Scotts Drop:

- First apply 2 header strips
- Operate spreader the longest way of the lawn
- Overlap wheels to insure uniform feeding
- Shut off spreader when you have reached the header strips or stopped
- Mow lawn first if grass is tall enough to touch the bottom of spreader



### Questions?

If you have questions you can talk directly with a Scotts Lawn Counselor. Simply dial the toll-free number 1-800-543-TURF (in Ohio, 1-800-762-4010).

**Scotts Policy**  
**No-Quibble Guarantee**

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

the lawn people  
Marysville, Ohio 43041

