

GUSSETS:

FRONT:

NET WEIGHT 24 LBS

(Scotts (R))

ProTurf

K-O-G TM

weed control

<b>ACCEPTED</b>
<b>JUN 8 1973</b>
<small>UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER EC. 136.22 SUBJECT TO ATTACHED COMMENTS.</small>

8773

8773

For Professional Use Only

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Do not contaminate domestic or irrigation waters. Do not store near fertilizers, insecticides, fungicides, seeds or foodstuffs. Avoid application on garden plants, flowers, and rootzone areas of trees and shrubs.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

Active ingredients: Dicamba (2-Methoxy-3, 6-dichlorobenzoic acid)	0.95%
Inert ingredients:	99.05%
Total	100.00%

US Pat No 3,083,089      EPA Reg No 538-25-AA  
ProTurf Division, O M Scott & Sons

Product of USA  
Marysville, Ohio 43040  
Oakland, Calif. 94604



K-O-G weed control

K-O-G weed control

(Scotts)  
ProTurf

(Scotts)  
ProTurf


8773

LACK:

(Scotts)  
ProTurf

K-O-G TM

weed control



USE THESE SPREADER SETTINGS  
to ensure proper distribution

FOR PUTTING GREENS

	SCOTT'S (drop type)	GANDY (42" Model)
QUARTER RATE treats 44,000 sq ft . . . . . to control clover and chickweed in bentgrass. Apply to moist foliage whenever weeds are actively growing. Repeat in three weeks if needed.	3½	—

FOR OTHER TURFGRASS AREAS

HALF RATE treats 22,000 sq ft . . . . . to control clover (except on putting greens)	5	28
NORMAL RATE treats 11,000 sq ft . . . . . to control knotweed, chickweed (common and mouse-ear), sheep sorrel, black medic	7	35
DOUBLE RATE treats 5,500 sq ft . . . . . to control wild onion and garlic, curly dock, ground ivy, Canada thistle, cudweed, yarrow, horseweed. (For control of wild onion, garlic, and cudweed - see below.)	10	45

FOR USE ON TURFGRASS ONLY

Apply whenever weeds are actively growing. Foliage may  
be either wet or dry.

For control of wild onion, wild garlic and cudweed start  
treatments in either spring or fall and repeat at 6 month  
intervals for a total of 2 or 3 applications as needed.

Avoid application in rootzone areas of trees and shrubs.

Do not apply when ground is frozen.

Delay planting grass seed until a heavy rain or watering  
and not sooner than two weeks after treatment.

On new plantings delay treatment until after the second  
mowing.

Not for use on St. Augustinegrass or Dichondra.

ProTurf Division, O M Scott & Sons    Marysville, Ohio 43040  
Oakland, Calif. 94609