


8714

Weedgrass  
Preventer

8714

Weedgrass  
PreventerWeedgrass  
PreventerWeedgrass  
PreventerWeedgrass  
PreventerWeedgrass  
Preventer

• prevents annual grassy weeds

8714

Net Weight 37 lbs (16.78 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazardous to Humans and Domestic Animals. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Do not contaminate feed or livestock or water used for agricultural or domestic purposes.

Environmental Hazard. This product is toxic to fish. Do not apply directly to water. Do not apply where runoff may occur. Do not apply when weather conditions favor drift from treated areas. Do not apply where there is a risk of drift to water. Avoid contact with water.

ACTIVE INGREDIENT  
S-ODIUM CHLORIDE 91.5%NET WEIGHTS  
37 lbs (16.78 kg)

EPA Reg. No. 538-26

Produced by The O. M. Scott & Sons Company  
Marysville, Ohio 43041

8714

For Professional Use Only

8714

Weedgrass  
Preventer

8714

Weedgrass  
Preventer

## Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Controls these grassy weeds as they sprout, without harm to desirable turf:

barnyardgrass	green foxtail
crabgrass	yellow foxtail
goosegrass	Poa annua

For use on established turfgrass, dichondra and putting greens

For most effective control of Poa annua (annual bluegrass) apply in late summer or early fall, before germination, and repeat yearly.

To control other grassy weeds listed above, apply in fall, winter or spring prior to germination.

Turf may be either moist or dry.

## Precautions

Do not use on putting greens composed of about 50% or more of Poa annua.

On newly planted areas, wait until the grass has been mowed at least twice and has made good coverage. On newly planted dichondra no waiting is necessary.

Where bermudagrass has been thinned out 25% or more by winter damage, diseases, or insects, with-

hold application until desired filling-in and rooting of the stolons is achieved.

After application avoid any severe aerating or raking which might disturb the surface and break the chemical barrier.

Delay seeding, sprigging or sodding for 4 "growing" months after application.

Applications should not exceed twice yearly.

## Storage and disposal

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bags by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

ProTurf Division  
The O. M. Scott & Sons Company  
Marysville, Ohio 43041

## Suggested Spreader Settings

to provide proper distribution of the spreader before application  
37 lbs (16.78 kg) treats 11,000 sq ft of normal rate.

SREADER	Ground or PTO Drive	Rate of Coverage	Spreading Width
Scott Drop	3 mph	6 feet	6
Scott Rotary	3 mph	8 feet	8
Lely-Webb	4.5 mph	16 feet	11
WFR Mowers			
Lely HST (PTO Mount)	4.50 mph	16 feet	11