



ProTurf Broad Spectrum Fungicide (XXXX)

FRONT:

5/2/83 JD/hs  
5/12/83 JD/hs

(Scotts (R) )  
ProTurf  
Brand

Broad Spectrum Fungicide


- . for use on winter turf diseases
- . prevents and controls pink and gray snow mold

NET WEIGHT 29 1/8 LBS (13.21 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS



Hazardsto Humans and Domestic Animals: CAUTION, causes eye injury. Avoid contact with eyes, skin, or clothing. In case of contact, immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. Harmful if swallowed. Avoid inhalation.

Environmental Hazards: Do not apply within twenty-five feet of any water body where fish are taken for human consumption. Toxic to fish and wildlife. Do not apply directly to water. Apply this product only as specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENTS:

Phenylmercuric acetate . . . . .	0.69%
Thiram (Tetramethylthiuram disulfide) . . . . .	4.65%

INERT INGREDIENTS: . . . . .	94.66%
TOTAL	100.00%

EPA Reg No 538-27

EPA Est 538-OH-1

Product of USA

ProTurf Division

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1983. The O M Scott & Sons Company. All rights reserved.

For Professional Use Only

XXXX

Broad Spectrum Fungicide

Broad Spectrum Fungicide

XXXX

(Scotts (R) )  
ProTurf  
Brand

Broad Spectrum Fungicide

---

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product shall be used only by or under the direct supervision of golf course superintendents.

For use on winter turf diseases.

Prevents or arrests fungus activity causing pink snow mold (Fusarium nivale) and gray snow mold (Typhula spp.).

For use on greens, tees and aprons

For the prevention of snow mold, apply at the NORMAL RATE in November or December prior to the first snow. Apply two repeat treatments at approximately 4-6 week intervals whenever the ground is free of snow.

To control or arrest severe infections or when heavy disease is anticipated, use the DOUBLE RATE.

Foliage may be either moist or dry when making application. On golf greens, rinse surface lightly after application to avoid pick up on shoes, golf balls or maintenance equipment.

Do not contaminate feed or foodstuffs.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bags by folding top down and securing. Protect from water and extreme heat.

PESTICIDE DISPOSAL: Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to Federal, State, or local procedures.

CONTAINER DISPOSAL: Completely empty bags by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

ProTurf Division

The O M Scott & Sons Company

Marysville, Ohio 43041

---

Con't

BACK Con't:

SUGGESTED SPREADER SETTINGS

to provide proper distribution calibrate spreader before application

29 1/8 lbs (13.21 kg) treats 11,000 ft<sup>2</sup> (1/4 acre/1022 m<sup>2</sup> at NORMAL RATE)

29 1/8 lbs (13.21 kg) treats 5,500 ft<sup>2</sup> (1/8 acre/ 511 m<sup>2</sup> at DOUBLE RATE)

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTING</u>	
			<u>NORMAL RATE</u>	<u>DOUBLE RATE</u>
Scotts (Drop Type)	3 mph	overlap wheels		
Scott Rotaries	3 mph			
Lely-WTR & WFR Models	4.5 mph			
Lely-HR (PTO Model) @	450 rpm 4.5 mph			