

FRONT:

**ACCEPTED**

MAR 20 1969

U.S. DEPARTMENT OF AGRICULTURE  
NATIONAL WEED CONTROL BOARD  
REGISTRATION NO. 538-32

TREATS  
5,000 sq ft/\$0.00  
NET WEIGHT 13 3/4 LBS

(Scotts)

SUPER  
CLOUT

Kills lawns of 36 kinds of weeds

<del>crabgrass</del>	<del>dandelions</del>	<del>chickweeds</del>
<del>foxtail</del>	<del>plantain</del>	<del>knotweed</del>
<del>black medic</del>	<del>dollarweed</del>	<del>matchweed</del>
<del>buckhorn</del>	<del>English daisy</del>	<del>Mexican clover</del>
<del>bur clover</del>	<del>evening primrose</del>	<del>penstemon</del>
<del>buttonweed</del>	<del>filaree</del>	<del>pepperweed</del>
<del>clover</del>	<del>foxglove</del>	<del>pigweed</del>
<del>cudweed</del>	<del>ground ivy</del>	<del>purple hellebore</del>
<del>curly dock</del>	<del>heal-all</del>	<del>ragweed</del>
<del>daisy fleabane</del>	<del>henbit</del>	<del>sheep sorrel</del>
<del>dallisgrass</del>	<del>lambquarters</del>	<del>shepherd's purse</del>
<del>dichondra</del>	<del>prickly lettuce</del>	<del>yellow rocket</del>

CAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin, eyes or clothing. After use, wash all exposed areas thoroughly. Clear water may be used as eye wash. Keep away from feed or food products. Do not contaminate water used for irrigation or domestic purposes. Avoid application on garden plants or flowers. Keep away from rootzone areas of shrubs and the branch spread of trees. Do not store near fertilizers, seeds, insecticides or fungicides.

Active ingredients: Monoammonium acid methanearsonate	4.50%
Maximum total water soluble arsenic	0.16%
2,4-Dichlorophenoxyacetic acid	1.10%
Dicamba (2-Methoxy-3, 6-dichlorobenzoic acid)	0.18%
Inert ingredients relative to herbicidal activity	94.16%
<b>Total</b>	<b>100.00%</b>

US Pat No 3,083,089  
O M SCOTT & SONS

USDA Reg No 538-32

Product of USA  
MANSFIELD, OHIO 44904


**BACK:**

Use any time weeds are growing actively. If crabgrass is involved, wait until it has grown enough to be plainly visible.

On new planted lawns wait until after the fourth mowing.

Apply to moist foliage. Choose a time when rain is not likely for 24 hours. Do not water in for 24 hours. Good turf grasses will color up again as watering is resumed.

Scotts promise to you: Satisfaction  
or your money back



Pull down tab for complete directions and helpful tips

SCOTT'S SPREADER  
SETTING . . . 6 1/2

TREATS  
5,000 sq ft/\$0.00

(Scotts)

SUPER  
CLOUT

Rids lawns of 36 kinds of weeds

<u>crabgrass</u>	<u>dandelions</u>	<u>chickweeds</u>
<u>foxtail</u>	<u>plantain</u>	<u>knotweed</u>
black medic	dollarweed	matchweed
buckhorn	English daisy	Mexican clover
bur clover	evening primrose	pennywort
buttonweed	filaree	pepperweed
clover	fog fruit	pigweeds
cudweed	ground ivy	purslane
curly dock	heal-all	ragweed
daisy fleabane	henbit	sheep sorrel
dallisgrass	lambquarters	shepherds purse
dichondra	prick'y lettuce	yellow rocket

Not for lawns of creeping bentgrass, St Augustine, centipede, bahia, carpetgrass, dichondra, lippia



SCOTT'S  
SUPER CLOUD

How to use

SUPER CLOUD

Controls 36 kinds of lawn weeds including weedy grasses, rosette and vining weeds

When to apply

Use any time weeds are growing actively.

Whenever crabgrass is involved, wait until it has grown enough to be plainly visible in the lawn (June - July).

On newly planted lawns wait until after the fourth mowing.

Growth should not be so tall as to touch bottom of spreader. Apply when lawn foliage is moist from dew, rain or sprinkling but when rain is not likely for 24 hours. Do not water in for at least 24 hours.

How to apply

Use Scotts Spreader at setting . . . . . 6½

Overlap spreader wheel tracks slightly so no strips are missed. Shut off spreader as you stop or turn around. (spreader picture)

Localized infestations may be spot or patch treated.

What to expect


Leaves and stems of rosette and vining weeds begin to twist in a few days but a month may pass before complete control is achieved. Grassy weeds first pale, then slowly turn yellow and brown.

Good turf grasses will be temporarily discolored but will recover with watering.

It is not necessary (or even desirable) to rake out dead weeds before seeding. They will eventually disintegrate and disappear if left alone.

Extremely heavy patches of mature crabgrass may not be completely wiped out by a single treatment. However, they will be sufficiently nullified to permit sowing grass seed without need for retreatment.

34



Precautions

Not for use on lawns of creeping bentgr., dichondra, lippia, St Augustine, bahia, centipede, rpetgrass or any lawns maintained like putting greens.

Avoid application on flowers, fruit or vegetable plants. Keep away from the rootzone areas of shrubs and the branch spread of trees.

Delay planting grass seed until after a heavy rain or watering but not sooner than 2 weeks after application.

Rain within 24 hours after application reduces effectiveness by washing granules off weed leaves. A repeat treatment may then be needed but wait at least one week.

**BACK:**

Use any time weeds are growing actively. If crabgrass is involved, wait until it has grown enough to be plainly visible.

On new planted lawns wait until after the fourth mowing.

Apply to moist foliage. Choose a time when rain is not likely for 24 hours. Do not water in for 24 hours. Good turf grasses will color up again as watering is resumed.

Scotts promise to you: Satisfaction  
or your money back

Put down tab for complete directions and helpful tips

SCOTTS SPREADER  
SETTING . . . 6½

TREATS  
5,000 sq ft/\$0.00

(Scotts)

SUPER  
CLOUT

Kills lawns of 36 kinds of weeds

crabgrass

foxtail

black medic  
buckhorn  
bur clover  
buttonweed  
clover  
cudweed  
curly dock  
daisy fleabane  
dallisgrass  
dichondra

dandelions

plantain

dollarweed  
English daisy  
evening primrose  
filaree  
fog fruit  
ground ivy  
heal-all  
henbit  
lambquarters  
prickly lettuce

chickweeds

knotweed

matchweed  
Mexican clover  
pennywort  
pepperweed  
pigweeds  
purslane  
ragweed  
sheep sorrel  
shepherds purse  
yellow rocket

Not for lawns of creeping bentgrass, St Augustine, centipede, bahia, carpetgrass, dichondra, lippia

How to use

**SUPER CLOUT**

Controls 36 kinds of lawn weeds including weedy grasses, rosette and vining weeds

---

When to apply

Use any time weeds are growing actively.

Whenever crabgrass is involved, wait until it has grown enough to be plainly visible in the lawn (June - July).

On newly planted lawns wait until after the fourth mowing.

Growth should not be so tall as to touch bottom of spreader. Apply when lawn foliage is moist from dew, rain or sprinkling but when rain is not likely for 24 hours. Do not water in for at least 24 hours.

How to apply .

Use Scotts Spreader at setting . . . . . 6½

Overlap spreader wheel tracks slightly so no strips are missed. Shut off spreader as you stop or turn around. (spreader picture)

Localized infestations may be spot or patch treated.

What to expect

Leaves and stems of rosette and vining weeds begin to twist in a few days but a month may pass before complete control is achieved. Grassy weeds first pale, then slowly turn yellow and brown.

Good turf grasses will be temporarily discolored but will recover with watering.

It is not necessary (or even desirable) to rake out dead weeds before seeding. They will eventually disintegrate and disappear if left alone.

Extremely heavy patches of mature crabgrass may not be completely wiped out by a single treatment. However, they will be sufficiently nullified to permit sowing grass seed without need for retreatment.

00-211

00-711

Precautions

Not for use on lawns of creeping bentgrass, dichondra, lippia, St Augustine, bahia, centipede, carpetgrass or any lawns maintained like putting greens.

Avoid application on flowers, fruit or vegetable plants. Keep away from the rootzone areas of shrubs and the branch spread of trees.

Delay planting grass seed until after a heavy rain or watering but not sooner than 2 weeks after application.

Rain within 24 hours after application reduces effectiveness by washing granules off weed leaves. A repeat treatment may then be needed but wait at least one week.