

FFII

Fertilizer and fungicide

8522

FFII

Fertilizer and fungicide

For use on Kentucky Bluegrass and St Augustinegrass
Do not apply on bentgrasses

Fertilizes grass. Prevents and controls stripe smut, leaf-spot, rust, brownpatch, dollarspot and snowmold. Intensifies and extends green color of grass

USE THESE SPREADER SETTINGS to ensure proper distribution

35 LBS TREATS 5,500 SQ FT (1/8 ACRE) AT NORMAL RATE
35 LBS TREATS 2,750 SQ FT (1/16 ACRE) AT DOUBLE RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTING	
			NORMAL RATE	DOUBLE RATE
Scotts (Drop Type)	3 mph	overlap wheels	9	13 1/2
Scotts Rotary	3 mph	8 feet	N	
Scotts 1000—8 ft	4.5 mph	overlap wheels	7 1/2	11
Lely-WTR & WFR (PTO) Models	4.5 mph	16 feet	6 III	
Lely-HR (PTO Model)	450 rpm @ 4.5 mph	16 feet	6 III	
Gandy 8 ft A Series	4.5 mph	overlap wheels	26	33

Model

To control active fungus on bluegrass

Stripe smut, leafspot, rust, dollarspot—apply at normal rate as soon as symptoms are first noticed

For areas heavily infected—apply at double rate or repeat normal rate in one month

To prevent fungus activity on bluegrass

Stripe smut and leafspot—apply normal rate spring and fall each year


Winter disease such as snowmold—apply prior to first expected snow

To prevent brownpatch fungus on St Augustinegrass

—apply normal rate in late winter or early spring and again in early fall (Sept-Oct). If extreme weather makes it necessary, treatment can be repeated any time after one month

Slight but temporary yellowing of grass can be expected following treatment. Rinse foliage following applications made when temperature is 80° or above

ProTurf Division, O M Scott & Sons Marysville, Ohio 43040
Oakland, Calif 94604



8522-40

Fertilizer and fungicide

Fertilizer and fungicide

Model

14-3-3

FF II

Fertilizer and fungicide
for bluegrass and St Augustinegrass

NOT FOR USE ON BENTGRASS

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid contact with skin, may cause irritation. Wash thoroughly after using. Avoid contamination of food.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not graze treated areas. Do not feed clippings to livestock.

Do not reuse container. Destroy when empty.

Active ingredient: Pentachloronitrobenzene
Inert ingredients: Relative to fungicidal activity

EPA Reg No 538.40 ZA

Net Weight 35 lbs ProTurf FF II 14.3.3

Total nitrogen (N)

0.8% ammoniacal nitrogen from ammoniated phosphate
8.5% water soluble nitrogen from urea and methylene ureas
4.7% water insoluble nitrogen from methylene ureas

Available phosphoric acid (P₂O₅)

from ammoniated phosphate

Soluble potash (K₂O)

from muriate of potash

Potential acidity equivalent to 300 lbs calcium carbonate per ton

US Pat No 3,076,700

Other Pat. pend

ProTurf Division, O M Scott & Sons

*In Michigan reads "minimum guaranteed analysis"

1: 40%

84.60%

Total: 100.00%

guaranteed analysis*

14%

3%

3%

Product of USA

Marysville, Ohio 42040

Oakland, Calif 94604

8522

8522

8522

