

FRONT:

controls weeds...fertilizes grass
(Scotts (R)) Bonus type B

Treats 2,500 sq ft/NET WEIGHT 18 LBS
\$5.95

31FO

ACCEPTED
OCT 3 1969
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 538-44

NEW!
Polyform
process
same benefits
in less weight

(Scotts (R))
Bonus (R)
BRAND
WEED CONTROL PLUS FERTILIZER
type B

for use only on ~~talia,~~
~~bermuda, zoysia, fescue~~
or bluegrass lawns

Controls 21 kinds of lawn
weeds. See list on other side

Gives long lasting feeding.
Ideal for iron-deficient soils

DO NOT USE ON ST AUGUSTINE,
DICHONDRA OR GROUND COVERS

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with the eyes, skin and clothing. Wash thoroughly after handling. Avoid application on flowers, fruit or vegetable plants. Keep away from rootzone areas of shrubs and the branch spread of trees. Do not store near fertilizer, seeds, insecticides or fungicides. Do not contaminate water used for irrigational or domestic purposes.

Active ingredients: 2,4-Dichlorophenoxyacetic acid	0.40%
Dicamba (2-Methoxy-3, 6-dichlorobenzoic acid)	0.07%
Inert ingredients (relative to herbicidal activity):	99.53%
Total	100.00%

US Pat No 3,231,363	USDA Reg No 538-44	Product of USA
Net Weight 18 lbs	Bonus type-B 25-5-3	guaranteed analysis
Total nitrogen (N), not less than		25%
nitrate nitrogen, not less than	0%	
ammonical nitrogen, not less than	1.6%	
water soluble organic nitrogen, not less than	15.0%	
water insoluble nitrogen, not less than	8.4%	
Available phosphoric acid (P ₂ O ₅), not less than		5%
insoluble phosphoric acid, not less than	0%	
Water soluble potash (K ₂ O), not less than		3%
Total available primary plant food		33%
Chlorine maximum		0.37%
Derived from: ammoniated phosphate, methylene ureas, urea, potassium sulfate		
Iron (Fe) (equivalent to Fe ₂ O ₃ -1.2%)		0.90%
Derived from ferrous ammonium sulfate		
Potential acidity equivalent to 1600 lbs calcium carbonate per ton.		
Florida Reg No 643-1108	Filed 1966	
O M Scott & Sons (Scotts (R)) Bonus type-B	controls weeds...fertilizes grass	

Other patent pending
Marysville, Ohio 43040
31FO

64

BACK:

controls weeds...fertilizes grass
(Scotts (R)) Bonus Type B

Treats 2,500 sq ft/\$5.95

31FO

NEW!
Polyform process
same benefits in less weight

(Scotts (R))
Bonus (R)
BRAND
WEED CONTROL PLUS FERTILIZER
type B

for use only on bahia, bermuda,
zoysia, fescue or bluegrass lawns

When to use
(SEE COMPLET DIRECTIONS IN POCKET BELOW)

To control weeds and fertilize grass, apply anytime after
mid-March when weeds are growing actively, up to the end
of October.

On newly seeded, sodded or sprigged lawns wait until new
grass has been mowed 3 or 4 times. After treatment, delay
any seeding until after heavy watering or rain—but never
sooner than a month.

Controls these weeds:

aster	daisy	pepperweed
black medic	treabana	pigweed
but clover	dandelion	plantain
butenweed	dichondra	prickly lettuce
centaury	dollarweed	primrose
chickweeds	Florida pusley	evening
cudweed	(Mexican clover)	purple
(rabbit tobacco)	matchweed	sorrel
	(creeping Charlie)	white clover

NOT A CONTROL FOR OXALIS OR GRASS-TYPE WEEDS SUCH AS CRABGRASS

Scotts No-Quibble Guarantee
If for any reason you are not satisfied with results after applying this
product with a Scotts Spreader, you are entitled to get your money back.
Simply send us evidence of purchase and we will mail you a refund check
promptly.

SCOTTS...the grass people
Marysville, Ohio 43040

Use Scotts Spreader (Standard or Automatic) at setting 7½
to ensure proper distribution

PULL DOWN
Directions
inside

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Avoid application on flowers, fruit or vegetable plants. Keep away from rootzone areas of shrubs and the branch spread of trees. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water used for irrigational or domestic purposes. Contains 2,4-Dichlorophenoxyacetic acid 0.4% (8.0 lbs per ton); Dicamba (2-Methoxy-3,6-dichlorobenzoic acid) 0.07% (1.4 lbs per ton). Apply 18 lbs per 2,500 sq ft on moist foliage after mid-March when weeds are growing actively.



(Scotts (R)) Bonus type-B
controls weeds...fertilizes grass

31F0

31500
40240

66

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Avoid application on flowers, fruit or vegetable plants. Keep away from rootzone areas of shrubs and the branch spread of trees. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water used for irrigation or domestic purposes. Contains 2,4-Dichlorophenoxyacetic acid 0.47 (8.0 lbs per ton); Dicamba (2-Methoxy-3,6-dichlorobenzoic acid) 0.07 (1.4 lbs per ton). Apply 18 lbs per 2,500 sq ft on moist foliage after mid-March when weeds are growing actively.

(Scotts (R)) Bonus type-B
controls weeds...fertilizes grass

31F0

GUIDE TO SUCCESSFUL USE

Bonus (R) TYPE B
BRAND

Weed control plus fertilizer for bahia,
bermuda, zoysia, fescue, bluegrass lawns

When to apply

Use on a calm day, any time after mid-March
when weeds are growing actively, up to the
end of October.

Apply when lawn is moist from dew, rain or
sprinkling but when rain is not likely for
24 hours.

Do not water for 24 hours after application.

On newly seeded, sodded or sprigged lawns
wait until the new grass has been mowed
3 or 4 times.



Spreader setting

To ensure proper distribution, use a Scotts Spreader (Standard or Automatic) at setting 7½

How to apply

As shown, the material is dropped in a swath only as wide as the hopper. To ensure complete coverage, overlap wheel tracks so you avoid "stripes" of untreated lawn.

(Spreader illus.)

The spreader should be "open" for application only as you push it forward. You will put on serious overapplication if you pull it backwards. Be sure to shut off spreader as you come to a stop and as you turn around.

What to expect

Stems and leaves of susceptible weeds begin to twist in a few days but a month of good growing weather may be required before control is complete. Action is slower if soil is dry or weather cool.

Rain, sprinkling, mowing or heavy traffic within 24 hours reduces weed control effectiveness by knocking granules off leaves. Repeat treatment may then be needed but wait at least one week.



Dead weeds will eventually disintegrate and disappear. It is not necessary (nor even desirable) to rake them out.

This product does not control oxalis or grass-type weeds such as crabgrass.

Precautions

Not for use on lawns of creeping bentgrass, St Augustine, dichondra, lippia or similar ground covers.

Avoid application on flowers, fruit or vegetable plants. Keep away from rootzone areas of shrubs and the branch spread of trees.

Sweep material off walks, driveways or similar surfaces to avoid discoloration.

Soft, lush grass may be temporarily discolored if temperature is consistently 90 or above.

After treatment, delay any seeding until after heavy watering or rain—but never sooner than a month.

Patience please!

We ask that you give this fine product time to do its job. It was designed deliberately to provide slow, steady, controlled action. The beneficial results may not be immediately evident to you. However, your patience will be well rewarded.

(Scotts)...the grass people
MARYSVILLE, OHIO 43040