ACCOR 116 × 54 . FOR FCONUMIC FO. TO ATTACHED COMMUNIA

Front:

</r>
</r>

Ц,

A)

iy

 $|\mathbf{x}'|$

NET WEIGHT 21 LBS

(Scotts (R))

THE GRASS PLOPLE

ProTurf (R) BRAND

Broad Spectrum

WEEDICIDE

patented TRIONIZED (R) bonding ensures sustained action

CAUTION: Keep out of reach of children. Avoid contact with skin, eyes or clothing. Avoid application on garden plants, flowers, shrubs and root zone areas of trees and shrubs. Do not store near fertilizers, seeds insecticides, or fungicides. Do not contaminate water used for irrigational or domestic purposes.

Active ingredients: 2,4-dichlorophenoxyacetic acid	2.95%
dicamba (2-methoxy-3, 6-dichlorobenzoic acid)	1.00%
Inert ingredients	96.05%
Total	100.00%

US Pat Nc 3,083,089

 $\mathcal{A}^{(k)}$

USDA Reg No 538-XX

Product of USA

 \rightarrow

0

ß

1284

Active ingredients: 2,4-dichlorophenoxyacetic acid 2.95%; dicamba (2-methoxy-3, 6-dichlorobenzoic acid) 1.00%. Inert ingredients 96.05%

CAUTION: Keep out of reach of children. Avoid contact with skin, eyes or clothing. Avoid application on garden plants, flowers, shrubs and root zone areas of trees and shrubs. Do not store near fertilizers, seeds insecticides or fungicides. Do not contaminate water used for irrigational or domestic purposes.



TO OPEN PULL WHITE TAPE

1.7

ProTurf BRAND

Broad Spectrum WEEDICIDE

(Scotts)

FOR PUTTING OREENS

Treats 40,000 sq ft -- Scotts Spreader setting 3½

Controls clover (Trifolium) and chickweed (Cerastium and Stellaria) in bentgrass or bermuda greens. Apply to moist or dry foliage whenever weeds are actively growing.

FOR OTHER TURFGRASS AREAS 19WAS

Treats 20,000 sq ft - Scotts Spreader setting

Back:

.

υ

۰۰ د ۱۰ د د

ి., ი Controls:

knotweed (Polygonum aviculare) sheep somel (Rumex acetosella) chickweeds (Cerastium vulgatum and Stellaria media) dandelion (Taraxacum officinale)

5

١

Also such wedds as: black medic bucknorn clover curlu dock English daisy filmee ground ty heal and hemoit horseweed pennywort peppargrass shepherds purse yellow rocket

Apply to moist foliage. Repeat in one month if needed.

FOR USE ON TURFGRASSES ONLY

Use when weeds are actively growing.

. •

Apply to foliage moist from dew, rain or sprinkling. Growth should not be so tall as to touch bottom of spreader.

Rain, sprinkling, mowing or heavy traffic within 24 hours after application knocks granules off weeds, necessitates treating again.

0

PRECAUTIONS

Do not apply when ground is frozen.

On new plantings delay treatment until after the third or fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 2 weeks after treating.

Soft, lush grass may become discolored temporarily in periods of high heat and humidity.

Not for use on St Augustinegrass or dichondra.

O M SCOTT & SONS

•

is' coc

`∂` 11 11 *' 8 0 ¹⁾

.

MARYSVILLE, OHIO

د. ب

14

1284

••

• 0